







CHEMISTRY (QUESTION BANK)**11.ALCOHOLS, PHENOLS AND ETHERS**

Single Correct Answer Type

- An organic compound 'X' on treatment with pyridinium chloro chromate in dichloromethane gives compound 'Y'. Compound 'Y' reacts with I₂ and alkali to form triiodomethane. The compound 'X' is
a) C₂H₅OH b) CH₃CHO c) CH₃COCH₃ d) CH₃COOH
- Ethyl alcohol is industrially prepared from the ethylene by:
a) Permanganate oxidation
b) Catalytic reduction
c) Absorbing in sulphuric acid followed by hydrolysis
d) Fermentation
- CH₂ClCH₂OH is stronger acid than CH₃CH₂OH because:
a) +IE of Cl disperses - ve charge on O -atom to produce more stable anion
b) -IE of Cl disperses - ve charge on O -atom to produce more stable anion
c) +IE of Cl increases - ve charge on O -atom to alcohol
d) None of the above
- Alcohol (CH₃)₂CHCH₂OH cannot be obtained by




- Lucas reagent is used to distinguish among primary, secondary and tertiary:
a) Alkyl halides b) Alcohols c) Aliphatic amines d) Aromatic amines
- Ketone upon treatment with Grignard reagent gives
a) Primary alcohol b) Secondary alcohol c) Tertiary alcohol d) Aldehyde
- The starting material for the preparation of CH₃I in one step reaction is:
a) CH₃OH b) C₂H₅OH c) CH₃CHO d) CH₃COCH₃
- From methyl alcohol we get:
a) Neoprene rubber
b) Perspex rubber
c) Bakelite a hard plastic
d) Sponge rubber
- Which one of the following will most readily be dehydrated in acidic condition?







- Tert*-butyl methyl ether on heating with anhydrous HI in ether gives
a) CH₃OH + (CH₃)₃Cl b) CH₃I + (CH₃)₃COH c) CH₃I + (CH₃)₃Cl d) None of the above
- Diethyl ether is decomposed on heating with:
a) NaOH b) Water c) KMnO₄ d) HI
- Ether fire can be extinguished by:
a) Sand b) Pyrene c) CO₂ d) All of these









13. Diethyl ether on reaction with CO in specific conditions forms:
 a) Acetic acid b) Carbon dioxide c) Ethyl propanoate d) Acetyl chloride
14. Most viscous among the following is:
 a) Propan-1-ol b) Propan-2-ol c) Propane-1, 2-diol d) Propane-1,2,3-triol
15. In the fermentation of sugar molasses, the percentage of ethanol formed is:
 a) 10 % b) 40 % c) 95 % d) 70 %
16. A liquid was mixed with ethanol and a drop of concentrated H_2SO_4 was added. A compound with a fruity smell was formed. The liquid was:
 a) HCHO b) CH_3COCH_3 c) CH_3COOH d) CH_3OH
17. Ethyl alcohol reacts with following to form a compound of fruity smell:
 a) PCl_5 b) $K_2Cr_2O_7 + H_2SO_4$ c) CH_3COOH d) CH_3COCH_3
18. Carbolic acid is
 a) HCOOH b) CH_3COOH c) C_6H_5COOH d) C_6H_5OH








A and *B* are



20. 2-methyl-2-butanol on treatment with HCl gives predominantly
 a) 2-chloro-3-methylbutane b) 2,2-dimethylpentane
 c) 2-chloro-2-methylbutane d) 1-chloro-2-methylbutane
21. In Williamson's synthesis ethoxy ethane is prepared by
 a) Passing ethanol over heated alumina
 b) Heating sodium ethoxide with ethyl bromide
 c) Treating ethyl alcohol with excess of H_2SO_4 at 430-440 K
 d) Heating ethanol with dry Ag_2O
22. Which of the following reacts fastest with a mixture of anhydrous $ZnCl_2$ and conc. HCl?
 a) Trimethyl carbinol
 b) Ethanol
 c) Propanol
 d) Methanol
23. Ethers are made free from peroxide linkage on distilling impure sample with:
 a) Conc. HNO_3 b) Conc. H_2SO_4 c) Conc. HCl d) None of these
24. Which of the property given below is not associated with glycerol?
 a) Formation of water and CO_2 on reduction
 b) Formation of tartronic acid on oxidation
 c) Formation of acrolein on dehydration
 d) Formation of allyl iodide with PI_3
25. The products obtained when anisole is heated in a sealed tube with HI are



27. The product in the given reaction is:




28. When CH_3MgI is made to react with acetone and the addition product formed is hydrolysed, we get:

- a) A primary alcohol b) A secondary alcohol c) A tertiary alcohol d) An aldehyde

29. The factor adversely affecting the process of fermentation is:

- a) Low concentration of sugar
b) High concentration of sugars
c) Presence of ammonium salts
d) Presence of air

30. The correct order of ease of dehydration of following is







- a) I > II > III b) III > II > I c) I > III > II d) III > I > II

31. The correct order of boiling point for primary (1°), secondary (2°) and tertiary (3°) alcohols is

- a) $1^\circ > 2^\circ > 3^\circ$ b) $3^\circ > 2^\circ > 1^\circ$ c) $2^\circ > 1^\circ > 3^\circ$ d) $2^\circ > 3^\circ > 1^\circ$

32. Which substance will not react with ϕNCl (ϕ = Phenyl) to give dye?






33. Phenol can be distinguished from ethanol by the following reagents except


- a) Sodium b) NaOH/I_2
c) Neutral FeCl_3 d) $\text{Br}_2/\text{H}_2\text{O}$

34. The compound which does not react with sodium is:





35. Ethylene glycol reacts with excess of PCl_5 to give
- a) $\text{CH}_3\text{CHOHCH}_3$ b) $\text{CH}_3-\text{O}-\text{CH}_3$ c) CH_3COOH d) $\text{C}_2\text{H}_5\text{OH}$
- a) 1, 1-dichloroethane b) 1, 2-dichloroethane
c) 1, 1, 1-trichloroethane d) 2, 2-dichloroethane
36. Alcohol is sometimes used in:
- a) Baking powder b) Paints c) Thermometers d) Weighing
37. Phenolphthalein is formed by condensation of phthalic anhydride and ϕOH . Which structure shows colour in basic medium?




d) All of the above







- a) $\text{C}_6\text{H}_5\text{OC}_2\text{H}_5$ b) $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$ c) $\text{C}_6\text{H}_5\text{OC}_6\text{H}_5$ d) $\text{C}_6\text{H}_5\text{I}$
39. The major product in the reaction of $\text{PhCH}_2\text{CH}(\text{OH})\text{CH}(\text{CH}_3)_2$ with concentrated H_2SO_4 is



40. Which is not an alcohol?
- a) $\text{CH}_2=\text{CHCH}_2\text{OH}$ b) $\text{CH}_2\text{OHCH}_2\text{OH}$ c) $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$ d) $\text{C}_6\text{H}_5\text{OH}$




The product A is







42. Glycerol catches fire on mixing with:
- a) KMnO_4 b) $\text{K}_2\text{Cr}_2\text{O}_7$ c) HNO_3 d) None of these
43. The end product of the reaction,

58.



Product is



59. Amongst the following, HBr reacts fastest with

a) Propane-1-ol

c) 2-methyl propane-1-ol

b) Propane-2-ol

d) 2-methyl propane-2-ol

60. Physical properties of:

a) Alcohols lie between alkanes and H₂Ob) H₂O lie between alcohols and alkenesc) Alkenes lie between alcohols and H₂O

d) None of the above

61. Which of the following ethers form peroxide readily?

a) Me—O—Me

b) Et—O—Et

c) iPr—O—iPr

d) Me—O—Et


62. Association of alcohol molecules takes place because of:

a) Electrovalent bond

b) Ionic bond

c) Covalent bond

d) Hydrogen bond

63. The reaction, $2\text{CH}_3\text{CH}_2\text{OH} \xrightarrow[413\text{ K}]{\text{H}^+} \text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ is believed to occur through the formation of

d) Both (b) and (c)

64. Ethyl iodide on treatment with dry Ag₂O will yield:





a) Ethyl alcohol

b) Diethyl ether

c) Ethyl methyl ether

d) Ethylene

65. Which of the following compounds is weakest acid?



66. Fusel oil is a mixture of:

a) Alcohols

b) Ethers

c) Ethers and alcohols




d) Alcohols and acetone

67. When benzene sulphonic acid and *p*-nitrophenol are treated with NaHCO₃, the gases released respectively area) SO₂, NO₂b) SO₂, NOc) SO₂, CO₂d) CO₂, CO₂

68. Which is correctly matched?

	Alcohol	α -H	β -H	Colour in Victor Meyer test
A.	X	3	0	Colourless
B.	Y	1	6	Blue
C.	Z	0	9	Red






- a) A and B
b) B and C
c) Only C
d) Only B
69. Lucas reagent is
a) Conc. HCl and anhydrous $ZnCl_2$
b) Conc. HNO_3 and hydrous $ZnCl_2$
c) Conc. HCl and hydrous $ZnCl_2$
d) Conc. HNO_3 and anhydrous $ZnCl_2$
70. An aldehyde on treatment with Zn/HCl yields:
a) 1° alcohol
b) 2° alcohol
c) 3° alcohol
d) None of these
71. In the reaction,


$$A \xrightarrow[H_2SO_4]{K_2Cr_2O_7} \text{acetone} \xrightarrow{\text{Oxidation}} \text{acetic acid, A is}$$
 a) 1-propanol
b) 2-butanol
c) 2-propanol
d) Ethanol
72. When glycerol is treated with excess of HI, it produces:
a) 2-iodopropane
b) Allyl iodide
c) Propene
d) Glycerol tri-iodide
73. The product obtained by the reaction of HBr with phenol is
 a) 
 b) 
 c) 
 d) There is no reaction
74. An ether is more volatile than an alcohol having the same molecular formula. This is due to
a) Dipolar character of ethers
b) Alcohols having resonance structures
c) Intermolecular hydrogen bonding in ethers
d) Intermolecular hydrogen bonding in alcohols
75. Glycol condenses with ketones to give:
a) Cyclic acetals
b) Cyclic ketals
c) Acetaldehyde
d) Oxalic acid
76. In the following reaction sequence

$$R-OH \xrightarrow{P+I_2} R-I \xrightarrow{AgNO_2} RNO_2 \xrightarrow{HNO_2} \text{no reaction}$$
 The alcohol is a
a) Primary alcohol
b) Secondary alcohol
c) Tertiary alcohol
d) Phenol
77. The explosive nitroglycerine is:
a) A soap
b) A salt
c) An ester
d) A complex compound
78. The compound $CH_3CH_2CH_2Br$ is converted into $CH_3CH_2CH_2OH$ by:
a) Dehydration
b) Hydrogenation
c) Elimination
d) Substitution
79. Consider the following reaction,

$$\text{ethanol} \xrightarrow{PBr_3} X \xrightarrow{\text{alc. KOH}} Y \xrightarrow[\text{(ii) } H_2O, \text{ heat}]{\text{(i) } H_2SO_4 \text{ at room temperature}} Z;$$
 The product Z is:
a) CH_3CH_2OH
b) $CH_2=CH_2$
c) $CH_3CH_2-O-CH_2-CH_3$

- d) $\text{CH}_3\text{—CH}_2\text{—O—SO}_3\text{H}$
80. Glycerol reacts with potassium bisulphate to produce
 a) Allyl iodide b) Allyl sulphate c) Acryl aldehyde d) Glycerol trisulphate
81. To prepare an ether by Williamson's synthesis, the reactants needed are
 a) Ethyl alcohol and tert butyl alcohol
 b) Sodium ethoxide and tert butyl bromide
 c) Sodium tertiary butoxide and ethyl bromide
 d) Sodium ethoxide and sodium tert butoxide
82. Fenton's reagent is:
 a) $\text{H}_2\text{O} + \text{FeSO}_4$ b) $\text{H}_2\text{O}_2 + \text{FeSO}_4$ c) $\text{H}_2\text{O}_2 + \text{ZnSO}_4$ d) $\text{NaOH} + \text{FeSO}_4$
83. Which of the following is simple ether?
 a) $\text{C}_6\text{H}_5\text{OCH}_3$ b) $\text{CH}_3\text{OC}_2\text{H}_5$ c) $n\text{PrOEt}$ d) MeOMe
84. The number of methoxy groups in a compound can be determined by treating it with:
 a) HI and AgNO_3 b) Sodium carbonate c) Sodium hydroxide d) Acetic acid
85. When $\text{C}_2\text{H}_5\text{OH}$ is mixed with ammonia and passed over heated alumina, the compound formed is:
 a) $\text{C}_2\text{H}_5\text{NH}_2$ b) C_2H_4 c) $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$ d) CH_3OCH_3
86. If there be a compound of the formula $\text{CH}_3\text{C}(\text{OH})_3$, which one of the following compounds would be obtained from it without treatment with any reagent?
 a) Methanol b) Ethanol c) Acetic acid d) Formaldehyde
87. In Lucas test an alcohol reacts immediately and gives insoluble chloride. The alcohol is
 a) CH_3OH b) $\text{CH}_3\text{CH}_2\text{OH}$ c) $(\text{CH}_3)_2\text{CHOH}$ d) $(\text{CH}_3)_3\text{COH}$
88. $(\text{CH}_3)_3\text{CONa}$ on reaction with CH_3Br will give:
 a) $(\text{CH}_3)_3\text{COC}(\text{CH}_3)_3$ b) CH_3OCH_3 c) $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ d) $(\text{CH}_3)_3\text{COCH}_3$
89. Which one has highest boiling point?
 a) Ethane b) Butane c) Butan-1-ol d) Pentane
90. Glyoxal is:
 a) $\text{CH}_2\text{OH—CHO}$ b) $\text{CH}_2=\text{OH}$ c) CHO—CHO d) $\text{CH}_2=\text{CHCHO}$
91. Methylated spirit is:
 a) Methanol containing some pyridine
 b) Ethanol containing some methanol
 c) Pure methanol
 d) 95% methanol
92. Dehydrogenation of 2-butanol gives:
 a) 2-butene b) Butanone c) Butyraldehyde d) 1-butene
93. The density of glycerol is higher than propanol due to
 a) Van der Waals' attraction b) Hydrogen bonding
 c) Ionic bonding d) More number of covalent bonds
94. Ethyl acetate is treated with double the molar quantity of $\text{C}_2\text{H}_5\text{MgBr}$ and the reaction mixture is hydrolysed with water. The product is:
 a) $\text{C}_2\text{H}_5\text{OH}$ b) $(\text{C}_2\text{H}_5)_2\text{CHOH}$ c) $\text{C}_2\text{H}_5\text{—}\begin{array}{c} \text{CH}_3 \\ | \\ \text{COH} \\ | \\ \text{C}_2\text{H}_5 \end{array}$ d) $\text{CH}_3\text{COOC}_2\text{H}_5$
95. The correct order of decreasing acidity of nitrophenols will be
 a) *m*-nitrophenol > *p*-nitrophenol > *o*-nitrophenol
 b) *o*-nitrophenol > *m*-nitrophenol > *p*-nitrophenol
 c) *p*-nitrophenol > *m*-nitrophenol > *o*-nitrophenol
 d) *p*-nitrophenol > *o*-nitrophenol > *m*-nitrophenol
96. The reaction of $\text{CH}_3\text{OC}_2\text{H}_5$ with HI gives:

- a) CH_3I only b) $\text{C}_2\text{H}_5\text{OH}$ only c) $\text{CH}_3\text{I} + \text{C}_2\text{H}_5\text{OH}$ d) $\text{C}_2\text{H}_5\text{I} + \text{CH}_3\text{OH}$
97. Glycerol has:
- 3 primary alcoholic groups
 - 3 secondary alcoholic groups
 - 1 primary alcoholic group and 2 secondary alcoholic groups
 - 2 primary alcoholic groups and 1 secondary alcoholic group
98. An ether is more volatile than an alcohol having the same molecule formula. This is due to
- Intermolecular hydrogen bonding in alcohols
 - Dipolar character of ethers
 - Alcohols having resonance structures
 - Intermolecular hydrogen bonding in ether
99. When phenol is heated with phthalic anhydride and H_2SO_4 , it produces
- Phenol red
 - Methyl orange
 - Salicylic acid
 - Phenolphthalein
100. When ethyl alcohol is dissolved in water, it is accompanied with:
- Absorption of heat and contraction in volume
 - Evolution of heat and contraction in volume
 - Absorption of heat and increase in volume
 - Evolution of heat and increase in volume
101. The products obtained when benzyl phenyl ether is heated with HI in the mole ratio 1:1 are
- Phenol
 - Benzyl alcohol
 - Benzyl iodide
 - Iodobenzene
- 1 and 3 only
 - 3 and 4 only
 - 1 and 4 only
 - 2 and 4 only
102. Which of the following is an example of elimination reaction?
- Chlorination of CH_4
 - Dehydration of $\text{C}_2\text{H}_5\text{OH}$
 - Nitration of benzene
 - Hydroxylation of C_2H_4
103. Glycerol on oxidation with conc. HNO_3 mainly yields:
- Glyceric acid
 - Tartronic acid
 - Mesoxalic acid
 - Both (a) and (b)
104. During fermentation little H_2SO_4 is added:
- To get acidic medium
 - To hydrolyse the glucose solution
 - To prevent the growth of undesirable bacteria
 - Which acts as dehydrating agent
105. The principal organic product in the reaction is:
- 
- 
 - 
 - 
 - 
106. Dialkyl sulphides are known as:
- Sulphonal
 - Mercaptan
 - Thioethers
 - Thioesters
107. Acrolein is obtained when glycerol is dehydrated with:
- KHSO_4
 - P_2O_5
 - Conc. H_2SO_4
 - All of these
108. In the following reaction, X and Y respectively are




- a) $\text{CH}_3\text{OH}, \text{C}_2\text{H}_5\text{OH}$ b) $\text{CH}_3\text{CHO}, \text{CH}_3\text{OH}$ c) $\text{CH}_3\text{CO}_2\text{H}, \text{C}_2\text{H}_5\text{OH}$ d) $\text{C}_2\text{H}_4, \text{CH}_3\text{CO}_2\text{H}$

109. The compound which gives turbidity immediately with Lucas reagent at room temperature is

- a) Butan-1-ol b) Butan-2-ol
c) 2-methyl propan-2-ol d) 2-methyl propan-1-ol

110. Which of the following will not react with NaOH?



- b) $\text{C}_2\text{H}_5\text{OH}$ c) CH_3CONH_2 d) $\text{CH}(\text{CN})_3$

111. The alcohol manufactured from water gas is

- a) CH_3OH b) $\text{C}_2\text{H}_5\text{OH}$ c) $\text{CH}_3\text{CH}_2\text{COOH}$ d) $(\text{CH}_3)_2\text{CHOH}$

112. The -OH group of an alcohol or the -COOH group of a carboxylic acid can be replaced by -Cl using

- a) Phosphorus pentachloride b) Hypochlorous acid
c) Chlorine d) Hydrochloric acid

113. Methanol cannot be dried with anhydrous CaCl_2 because

- a) CaCl_2 dissolves in it b) It is not good dehydrating agent
c) It forms a solid $\text{CaCl}_2 \cdot 4\text{CH}_3\text{OH}$ d) It reacts with CH_3OH

114. Sodium ethoxide has reacted with ethanoyl chloride. The compound that is produced in the above reaction is:


- a) Diethyl ether b) 2-Butanone c) Ethyl chloride d) Ethyl ethanoate

115. Which method is employed to convert alkyl halide into alcohol?

- a) Substitution b) Addition c) Dehydration d) Rearrangement

116. Lucas test is associated with

- a) Aldehydes b) Phenols c) Carboxylic acids d) Alcohols





In the above sequence, C is

- a) *o*-bromophenol b) *p*-bromophenol
c) *m*-bromophenol d) 2, 4, 6-tribromophenol

118. The boiling points of thio-ethers are...than those of ether.

- a) Lesser b) Equal c) Higher d) None of these



In the above sequence D is

- a) $\text{CH}_3\text{COOC}_2\text{H}_5$ b) $\text{CH}_3\text{COOCH}_3$ c) $\text{C}_2\text{H}_5\text{COOC}_2\text{H}_5$ d) $(\text{C}_2\text{H}_5)_2\text{O} \rightarrow \text{BF}_3$

120. The toxicity order for $\text{CH}_3\text{OH}, \text{C}_2\text{H}_5\text{OH}$ and $\text{C}_3\text{H}_7\text{OH}$ is:


- a) $\text{C}_2\text{H}_5\text{OH} < \text{CH}_3\text{OH} < \text{C}_3\text{H}_7\text{OH}$
b) $\text{C}_3\text{H}_7\text{OH} < \text{C}_2\text{H}_5\text{OH} < \text{CH}_3\text{OH}$
c) $\text{C}_2\text{H}_5\text{OH} < \text{C}_3\text{H}_7\text{OH} < \text{CH}_3\text{OH}$
d) $\text{CH}_3\text{OH} < \text{C}_2\text{H}_5\text{OH} < \text{C}_3\text{H}_7\text{OH}$

121. The alcohol that forms fats with fatty acids is:

- a) Glycerol b) Ethanol c) Methanol d) Glycol


122. The reduction,

- c) $\text{CH}_3 - \text{CH}_2 - \text{I}$ d) None of the above
139. A mixture of alcohol and ether is called:
 a) Natalite b) Power alcohol c) Peroxide d) None of these



In this reaction, the end product C is

- a) Salicylaldehyde b) Salicylic acid c) Phenyl acetate d) Aspirin
141. In fermentation by zymase, alcohol and CO_2 are obtained from
 a) Invert sugar b) Glucose c) Fructose d) All of these
142. Oxidation of allyl alcohol, $(\text{CH}_2=\text{CH}-\text{CH}_2\text{OH})$ gives a mixture of oxalic acid and formic acid. If this oxidation is done in presence of bromine. One would expect only:
 a) Oxalic acid b) Formic acid c) Succinic acid d) Acrylic acid
143. In the given transformation, which of the following is the most appropriate reagent?







- a) $\text{Zn} - \text{Hg}/\text{HCl}$ b) Na, Liq. NH_3 c) NaBH_4 d) $\text{NH}_2\text{NH}_2, \text{OH}$





144. Glycerol is highly viscous. It is due to the fact that:
 a) It is highly polar
 b) It forms extensive H-bonding
 c) It shows intramolecular H-bonding
 d) It has high b.p.
145. The best method to prepare cyclohexene from cyclohexanol is by using
 a) Conc. $\text{HCl} + \text{ZnCl}_2$ b) Conc. H_3PO_4 c) HBr d) Conc. HCl
146. Phenol on treatment with diethyl sulphate in presence of NaOH gives
 a) Phenetole b) Anisole c) Diphenyl ether d) Diethyl ether
147. Vapours of an alcohol were passed over hot reduced copper. It gave an olefin. The alcohol is:
 a) Primary b) Secondary c) Tertiary d) None of these
148. Propane, $\text{CH}_3 - \text{CH} = \text{CH}_2$ can be converted into 1-propanol by oxidation. Which set of reagents among the following is ideal to effect the conversion?
 a) H_2O b) $\text{B}_2\text{H}_6, \text{H}_2\text{O}_2$ c) H_2SO_4 d) None of these
149. $\text{C}_4\text{H}_{10}\text{O}$ gives white precipitate within 5 min with concentrated hydrochloric acid in the presence of anhydrous zinc chloride.

Alcohol can be



150. Propan-2-ol on reacting with Cl_2 produces:
 a) Trichloroethanal b) Trichloroacetone c) Acetone d) None of these
151. Which of the following compounds is resistant to nucleophilic attack by hydroxyl ions?
 a) Acetamide b) Methyl acetate c) Diethyl ether d) Acetonitrile
152. Ethers are quite stable towards:
 a) Oxidizing agents b) Reducing agents c) Na metal d) All of these
153. The function of ZnCl_2 in Lucas test for alcohols is




- a) To act as acid catalyst and react with HCl to form H_2ZnCl_4
 b) To act as base catalyst and react with NaOH to form $Na_2Zn(OH)_4$
 c) To act as amphoteric catalyst
 d) To act as neutral catalyst
154. When ethyl alcohol is heated with conc. H_2SO_4 , the product obtained is
 a) $CH_3COOC_2H_5$ b) C_2H_2 c) C_2H_6 d) C_2H_4
155. Phenol is heated with phthalic anhydride in presence of conc H_2SO_4 . The product gives pink colour with alkali. The product is
 a) Phenolphthalein b) Bakelite c) Salicylic acid d) Fluorescein
156. The action of halogen acids on an ether, has the following order of reactivity:
 a) $HCl > HBr > HI$ b) $HI > HCl > HBr$ c) $HI > HBr > HCl$ d) $HCl > HI > HBr$
157. Rectified spirit contains:
 a) 75.0 % alcohol b) 85.5% alcohol c) 95.6% alcohol d) 100.0% alcohol
158. Phenyl magnesium bromide reacts with methanol to give a mixture of:
 a) Anisole and $Mg(OH)Br$
 b) Benzene and $Mg(OMe)Br$
 c) Toluene and $Mg(OH)Br$
 d) Phenol and $Mg(Me)Br$
159. Phenol $\xrightarrow{NaNO_2/H_2SO_4} B \xrightarrow{H_2O} C \xrightarrow{NaOH} D$
 Name of the reaction is
 a) Liebermann's reaction b) Phthalein fusion test
 c) Reimer-Tiemann reaction d) Schotten-Baumann reaction
160. The commonly used dehydrating agent in the preparation of an ester is:
 a) P_2O_5 b) Anhydride $CaCl_2$ c) Anhydride $AlCl_3$ d) Conc. H_2SO_4
161. Nobel's oil is:
 a) Fire extinguisher b) Insecticide c) Explosive d) Detergent
162. Phenol, *p*-methylphenol, *m*-nitrophenol and *p*-nitrophenol follows order of increasing strength as
 a) Phenol, *p*-methylphenol, *p*-nitrophenol, *m*-nitrophenol
 b) *p*-methylphenol, phenol, *m*-nitrophenol, *p*-nitrophenol
 c) *p*-methylphenol, *m*-nitrophenol, phenol, *p*-nitrophenol
 d) *m*-nitrophenol, *p*-nitrophenol, phenol, *p*-methylphenol
163. Ethylene glycol on oxidation with per-iodic acid gives:
 a) Oxalic acid b) Glyoxal c) Formaldehyde d) Glycollic acid
164. 
 a) $C_6H_5OC_2H_5$ b) $C_2H_5OC_2H_5$ c) $C_6H_5OC_6H_5$ d) C_6H_5I
165. The major product of the following reaction,
 $C_6H_5CH = CHCH_3 \xrightarrow[(ii) NaBH_4]{(i) Hg(OAc)_2, THF-H_2O}$ is
 a)  b) 
 c)  d) 
166. The structure of the compound that gives a tribromo derivative on treatment with bromine water is



167. Which of the following reagents may be used to distinguish between phenol and benzoic acid?


- a) Aqueous NaOH b) Tollen's reagent c) Molisch reagent d) Neutral FeCl₃

168. Which is obtained on treating phenol, with dilute HNO₃?



- d) None of these


169. Consider the following reaction,



Among the following, which one cannot be formed as a product under any conditions?

- a) Ethyl hydrogen sulphate b) Ethylene
c) Acetylene d) Diethyl ether

170. Dehydration of the following in increasing order is




- a) I < II < III < IV b) II < III < IV < I c) I < II < III < IV d) I < IV < II < III

171. Excess of glycol when dehydrated gives:

- a) Ethylene oxide b) Ethanol c) Acrolein d) 1,4-dioxan

172. In the reduction,



The catalyst used is:

- a) Ni b) Pd c) Pt d) All of these

173. Action of HNO₂ on CH₃NH₂ gives:

- a) CH₃OH b) CH₃ · O · CH₃ c) CH₃O—N=O d) Both (b) and (c)





174. Primary and secondary alcohols on action of reduced copper give:

- a) Aldehydes and ketones respectively
b) Ketones and aldehydes respectively
c) Only aldehydes
d) Only ketones

175. Diethyl ether absorbs oxygen to form:

- a) Red coloured sweet smelling compound
b) Acetic acid
c) Ether suboxide
d) Ether peroxide

176. (A) $\xrightarrow{HIO_4}$ cyclohexanone + HCHO. What is (A)?





177. Which of the following undergoes dehydration most readily?

- a) 1-phenyl-1-butanol b) 1-phenyl-2-butanol c) 2-phenyl-2-butanol d) 2-phenyl-1-butanol

178. Ether in contact with air for a long time form peroxides. The presence of peroxide in ether can be tested by adding Fe⁺² ion in it and then adding:

- a) KCNS b) SnCl₂ c) HgCl₂ d) KI

179. Cyclohexanol is a:
 a) Phenol b) Primary alcohol c) Sec. alcohol d) *tert.* Alcohol
180. Glycerol on oxidation with dil. HNO_3 gives:
 a) Tartronic acid b) Mesoxalic acid c) Oxalic acid d) Glyceric acid
181. Butan-2-ol is:
 a) Primary alcohol b) Secondary alcohol c) Tertiary alcohol d) None of these
182. Peppermint can be extracted from plant sources by using solvents like:
 a) NH_3 b) H_2O c) CH_3COOH d) $\text{C}_2\text{H}_5\text{OH}$
183. Chlorine reacts with ethanol to give:
 a) Ethyl chloride b) Chloroform c) Acetaldehyde d) Chloral
184. Molasses contains:
 a) 70 % sugar b) 50% sugar c) 60% sugar d) 10% sugar
185. Which of the following are known as mercaptans?
 a) Thio-alcohols b) Thio-ethers c) Thio-aldehydes d) Thio-acids
186. Which forms most stable hydrate?
 a) CH_3CHO b) $\text{C}_6\text{H}_5\text{CHO}$ c) CCl_3CHO d) CH_3COCH_3
187. An organic compound dissolved in dry benzene evolved hydrogen on treatment with sodium. It is:
 a) A ketone b) An aldehyde c) A tertiary amine d) An alcohol
188. Sodium ethoxide is obtained by the reaction of ethyl alcohol with:
 a) NaOH b) Na c) NaCl d) NaHCO_3
189. Which one of the following compounds will not react with CH_3MgBr ?
 a) Ethyl acetate b) Acetone c) Dimethyl ether d) Ethanol
190. The major organic product in the reaction,
 $\text{CH}_3-\text{O}-\text{CH}(\text{CH}_3)_2 + \text{HI} \rightarrow$ Product is:
 $\text{CH}_3\text{OC}(\text{CH}_3)_2$
 a) $\begin{array}{c} | \\ \text{I} \end{array}$ b) $\text{CH}_3\text{I} + (\text{CH}_3)_2\text{CHOH}$ c) $\text{CH}_3\text{OH} + (\text{CH}_3)_2\text{CHI}$ d) $\text{ICH}_2\text{OCH}(\text{CH}_3)_2$
191. Structure of diethyl ether can be confirmed by:
 a) Kolbe's synthesis
 b) Frankland's synthesis
 c) Wurtz's synthesis
 d) Williamson's synthesis
192. Glycerol on oxidation with bismuth nitrate mainly gives:
 a) Glyceric acid b) Tartronic acid c) Mesoxalic acid d) Oxalic acid
193. The end product of the following sequence is:
 $\text{CH}_3\text{Br} \xrightarrow{\text{KCN(alc.)}} (\text{A}) \xrightarrow{\text{H}_3\text{O}^+} (\text{B}) \xrightarrow[\text{Ether}]{\text{LiAlH}_4} (\text{C})$
 a) CH_3CHO b) $\text{CH}_3\text{CH}_2\text{OH}$ c) CH_3COCH_3 d) CH_4
194. Saponification means hydrolysis of an ester with:
 a) Enzyme b) CH_3COOH c) H_2SO_4 d) NaOH
195. Which of the following can work as dehydrating agent for alcohols?
 a) H_2SO_4 b) Al_2O_3 c) H_3PO_4 d) All of these
196. In $\text{CH}_3\text{CH}_2\text{OH}$ the bond which most readily undergoes heterolytic cleavage during its reaction with $\text{CH}_3\text{COOH}/\text{H}_2\text{SO}_4$ is:
 a) $\text{C}-\text{C}$ b) $\text{C}-\text{O}$ c) $\text{O}-\text{H}$ d) $\text{C}-\text{H}$
197. When ethyl alcohol vapours mixed with air, are passed over heated platinized asbestos, the compound formed is:
 a) Acetaldehyde b) Diethyl ether c) Acetone d) None of these
198. Which of the following reactions will not yield *p*-*tert* butylphenol?
 a) CH_3 b) Phenol + $(\text{CH}_3)_3\text{COH} \xrightarrow{\text{H}^+}$



199. One mole of an organic compound *A* with the formula $\text{C}_3\text{H}_8\text{O}$ reacts completely with two moles of HI to form *X* and *Y*. When *Y* is boiled with aqueous alkali it forms *Z*. *Z* answers the iodoform test. The compound *A* is


- a) Propan-2-ol b) Propan-1-ol c) Ethoxyethane d) Methoxyethane

200. Which one of the following alcohol is used as an antifreeze reagent for making explosives?

- a) Glycerol b) Glycol c) Ethanol d) Phenol

201. The IUPAC name of $\text{CH}_3\text{OCH}(\text{CH}_3)_2$ is:

- a) 1-methoxy propane
b) 3-methoxy propane
c) Methyl-isopropylether
d) 2-methoxy propane



is an example of

- a) 1, 2-addition of HCl followed by tautomerism b) 1, 2-addition followed by reduction
c) 1, 4-addition followed by tautomerism d) 1, 4-addition followed by oxidation

203. Absolute ethanol cannot be obtained by simple fractionation of a solution of ethanol and water because:

- a) Their boiling points are very near
b) Ethanol remains dissolved in water
c) They form a constant boiling mixture
d) Ethanol molecules are solvated

204. Etherates are

- a) Ethers b) Solution in ether
c) Complexes of ethers with Lewis acid d) Complexes of ethers with Lewis base



205. Glycerol is not used in:

- a) Cosmetics b) Matches c) Explosives d) Soaps

206. Which will not form a yellow precipitate on heating with an alkaline solution of iodine?

- a) $\text{CH}_3\text{CHOHCH}_3$ b) $\text{CH}_3\text{CH}_2\text{CHOHCH}_3$ c) CH_3OH d) $\text{CH}_3\text{CH}_2\text{OH}$

207. Which of the following is an alkoxide?

- a)  b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{ONa}$ c) $\text{CH}_2\text{OH} \cdot \text{CH}_2\text{OH}$ d) 

208. The acidic character of 1° , 2° , 3° alcohols, H_2O and $\text{RC} \equiv \text{CH}$ is of the order

- a) $\text{H}_2\text{O} > 1^\circ > 2^\circ > 3^\circ > \text{RC} \equiv \text{CH}$ b) $\text{RC} \equiv \text{CH} > 3^\circ > 2^\circ > 1^\circ > \text{H}_2\text{O}$
c) $1^\circ > 2^\circ > 3^\circ > \text{H}_2\text{O} > \text{RC} \equiv \text{CH}$ d) $3^\circ > 2^\circ > 1^\circ > \text{H}_2\text{O} > \text{RC} \equiv \text{CH}$

209. The enzyme which can catalyse the conversion of glucose to ethanol is:

- a) Zymase b) Diastase c) Maltase d) Invertase

210. Oxygen atom of ether is:

- a) Very active b) Replaceable c) Active d) Comparatively inert

211. Argol, a brown crust, formed during the fermentation of grape juice contains

- a) CO_2 b) Fused oil
c) Potassium hydrogen tartarate d) lye


212. Benzoylation of phenol in alkaline medium is known as

- a) Friedel-Crafts reaction
 b) Wurtz-Fittig reaction
 c) Schotten-Baumann reaction
 d)

213. The prospective fuel 'gashol' is a mixture of:

- a) Gaseous hydrocarbons and heavy water
 b) Petrol and phenol
 c) Petrol and ethanol
 d) Radioactive substances

214. Identify the product/s in the following reaction.



Products + H_3BO_3

- a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ b) $\text{CH}_3\text{CHOHCH}_3$ c) $\text{CH}_3\text{CH}_2\text{CHO}$ d) $\text{CH}_3\text{CH}_2\text{OH} + \text{CH}_3\text{OH}$

215. A fruity smell is obtained by the reaction of ethanol with

- a) CH_3COCH_3 b) PCl_5 c) CH_3COOH d) CH_3CHO







216. Which of the following reactions does not yield an ether?

- a) Sodium methoxide reacts with dimethyl sulphate
 b) Sodium ethoxide reacts with ethyl bromide
 c) Sodium ethoxide reacts with bromocyclopropane
 d) Ethanol reacts with CH_2N_2 in presence of HBF_4






217. An alcohol on alk. KMnO_4 oxidation gives first acetone and on further oxidation acetic acid. It is:

- a) Ethyl alcohol
 b) Isopropyl alcohol
 c) Primary alcohol
 d) None of these

218. Which is not the intermediate stage of following conversion?






219.



220. When diethyl ether is heated with an excess of PCl_5 , it yields

- a) Ethyl chloride b) Diethyl ether peroxide
 c) Ethanoyl chloride d) Perchlorodiethyl ether

221. Which of the following represents the Dow process for the manufacture of phenol?



d) None of the above

222. The organic compound present in tincture of iodine is:

- a) Alcohol b) CCl₄ c) Acetone d) CS₂

223. Phenol on heating with CCl₄ and aqueous KOH gives salicylic acid. This reaction is

- a) Friedel-Craft reaction b) Diels-Alder reaction
c) Reimer-Tiemann reaction d) Wittig reaction

224. The —OH group of methyl alcohol cannot be replaced by chlorine by the action of:

- a) Chlorine b) HCl c) PCl₃ d) PCl₅


225. The following substance can be used as a raw material for obtaining alcohol:

- a) Potatoes b) Molasses c) Maize d) All of these

226. On oxidation, an alcohol gives an aldehyde having the same number of carbon atoms as that of alcohol. The alcohol is:

- a) 1° alcohol
b) 2° alcohol
c) 3° alcohol
d) None of these

227. The end product of which of the following reaction is isomer of alcohols?




228. From amongst the following alcohols the one that would react fastest with conc. HCl and anhydrous ZnCl₂ is

- a) 2-butanol b) 2-methyl propan-2-ol c) 2-methylpropanol d) 1 butanol

229. Which of the following is least soluble in water?

- a) C₂H₅OH b) C₃H₇OH c) C₄H₉OH d) C₅H₁₁OH


230. The reaction given below is called:



- a) Kharasch effect b) Wurtz reaction c) Darzen's reaction d) Hunsdicker reaction


231. The compound with formula C₄H₁₀O yields a compound C₄H₈O on oxidation. The compound C₄H₁₀O is:

- a) An aldehyde b) An alcohol c) A ketone d) An anhydride





232. Reaction of  with RMgX followed

with hydrolysis produces :

- a) RCHOHR b) RCH₂CH₂OH c) RCHOHCH₃ d) RCH=CHOH



The product is



234. C_2H_5OH cannot be dried by anhydrous $CaCl_2$, because:

- C_2H_5OH is soluble in water
- Explosion takes place
- C_2H_5OH reacts with $CaCl_2$
- None of the above

235. Denatured spirit is mainly used as a:

- Good fuel
- Drug
- Solvent in preparing varnishes
- Material in the preparation of oil

236. The dehydration of 2-methyl butanol with conc. H_2SO_4 gives

- 2-methyl butane as major product
- Pentene
- 2-methyl but-2-ene as major product
- 2-methyl pent-2-ene

237. Ethers are not distilled to dryness for fear of explosion. This is due to formation of:

- Oxides
- Alcohol
- Ketones
- Peroxides

238. Tertiary alcohols (3°) having at least four carbon atoms upon drastic oxidation yield carboxylic acid with

- One carbon atom less
- Two carbon atoms less
- Three carbon atoms less
- All the above three options are correct

239. Lucas reagent is

- Anhydrous $AlCl_3$ with concentrated HCl
- Anhydrous $ZnCl_2$ and concentrated H_2SO_4
- Anhydrous $ZnCl_2$ and concentrated HCl
- Anhydrous $CaCl_2$ and concentrated HCl

240. The cleavage of an aryl-alkyl ether with cold HI gives

- Alkyl iodide and water
- Aryl iodide and water
- Alkyl iodide, aryl iodide and water
- Phenol and alkyl iodine


241. Phenol is heated with a solution of mixture of KBr and $KBrO_3$. The major product obtained in the above reaction is

- 2-bromophenol
- 3-bromophenol
- 4-bromophenol
- 2, 4, 6-tribromophenol

242. For the preparation ter-butylmethylether by Williamson's method the correct choice of reagents is:

- Methoxide and ter-butylbromide
- Methanol and 2-bromobutane
- 2-butanol and methylbromide
- Ter-butoxide and methylbromide

243. Consider the following reactions,



Y can be converted to X on heating with... at ... temperature.


- $Al_2O_3, 350^\circ C$
- $Cu, 300^\circ C$
- $\frac{Ca(OH)_2}{+ CaOCl_2, 60^\circ C}$
- $NaOH/I_2, 60^\circ C$

244. Which of the following methods cannot be used for the preparation of an ester?



- $RCOOH + R' OH + OH^-$
- $RCOCl + R' OH + Pyridine$
- $RCOOH + R' OH + H^+$
- $(RCO)_2O + R' OH + Pyridine$

245. Oxygen containing organic compound upon oxidation forms a carboxylic acid as the only organic product with its molecular mass higher by 14 units. The organic compound is


- a) An aldehyde b) A primary alcohol c) A secondary alcohol d) A ketone
246. A compound X with molecular formula C_3H_8O can be oxidised to a compound Y with the molecular formula $C_3H_6O_2$. X is most likely to be:
- a) Primary alcohol b) Secondary alcohol c) Aldehyde d) Ketone
247. $HOH_2C \cdot CH_2OH$ on heating with periodic acid gives:
- a) $2 \begin{array}{l} H \\ \diagdown \\ C=O \\ \diagup \\ H \end{array}$ b) $2 CO_2$ c) $2 HCOOH$ d) $\begin{array}{c} CHO \\ | \\ CHO \end{array}$
248. Reaction of tertiary butyl alcohol with hot Cu at $350^\circ C$ produces
- a) Butanol b) Butanal c) 2-butene d) Methylpropene
249. Ethyl chloride is converted into diethyl ether by
- a) Perkins reaction b) Grignard reagent
c) Wurtz reaction d) Williamson's synthesis
250. The product obtained by heating diethyl ether with HI is
- a) C_2H_5I b) C_2H_5OH c) $C_2H_5OH + C_2H_5I$ d) $C_2H_5 - C_2H_5$
251. The reaction,
 $C_2H_5ONa + C_2H_5I \rightarrow C_2H_5OC_2H_5 + NaI$ is known as
- a) Kolbe's synthesis b) Wurtz's synthesis
c) Williamson's synthesis d) Grignard's synthesis
252. Which one can differentiate between C_2H_5OH and CH_3OH ?
- a) H_2O b) $Na_2CO_3 + I_2$ c) NH_3 d) HCl
253. Ethylene oxide when, treated with Grignard reagent yields:
- a) Cyclopropyl alcohol b) Primary alcohol c) Secondary alcohol d) Tertiary alcohol
254. Among the following compounds which can be dehydrated very easily?
- a) $CH_3CH_2CH_2CH_2CH_2OH$ b) $\begin{array}{c} OH \\ | \\ CH_3CH_2CH_2CHCH_3 \end{array}$
c) $\begin{array}{c} CH_3 \\ | \\ CH_3CH_2CCH_2CH_3 \\ | \\ OH \end{array}$ d) $\begin{array}{c} CH_3CH_2CHCH_2CH_2OH \\ | \\ CH_3 \end{array}$
255. Catalytic dehydrogenation of a primary alcohol gives a
- a) Secondary alcohol b) Aldehyde c) Ketone d) Ester
256. Action of nitrous acid on ethyl amine gives:
- a) C_2H_6 b) C_2H_5OH c) NH_3 d) nitromethane
257. Which of the following compounds is most acidic?
- a) CH_4 b) C_2H_6 c) $CH \equiv CH$ d) C_2H_5OH
258. $2\text{-propanol} + NaBr \xrightarrow{\text{Reflux}} X$. What is X ?
- a) 2-bromopropane
b) Propane
c) Propene
d) Propanone
259. Which of the following reaction is/are feasible?



a)



b)



c) Both (a) and (b)

d) None of the above

260. Alcohols are neutral in character whereas thio-alcohols arein character.

a) Strongly acidic

b) Weakly acidic

c) Basic

d) Neutral

261. On boiling with concentrated hydrobromic acid, phenyl ethyl ether yields


a) Phenol and ethane

b) Phenol and ethyl bromide

c) Bromobenzene and ethanol


d) Bromobenzene and ethane

262. General formula of primary alcohol is:



d) All of these

263. The compound *B* formed in the following sequence of reactions,



a) Propyne

b) Propene

c) Propanal

d) Propane

264. Formation of diethyl ether from ethanol is based on a

a) Dehydration reaction

b) Dehydrogenation reaction

c) Hydrogenation reaction

d) Homolytic fission reaction

265. Two aromatic compounds having formula $\text{C}_7\text{H}_8\text{O}$ which are easily identifiable by FeCl_3 solution test (violet colouration) are


a) *o*-cresol and benzyl alcohol

b) *m*-cresol and *p*-cresol

c) *o*-cresol and *p*-cresol

d) Methyl phenyl ether and benzyl alcohol

266. In the reaction,



a) HCHO , HCOONH_4

b) HCOOH , HCOONH_4

c) HCOOH , HCONH_2

d) HCHO , HCONH_2

267. Acetic acid and methanol are obtained on a large scale by destructive distillation of:

a) Wood

b) Coal

c) Turpentine oil

d) CH_3COOH

268. Which of the following statement is incorrect?


a) Enzymes are in colloidal state

b) Enzymes are catalyst

c) Enzymes can catalyse any reaction

d) Urease is an enzyme

269. In the following sequence the product (C) is:




a) Alcohol

b) Ether

c) Alkene

d) None of these

270. In the reaction, the products formed are:



a) $(\text{CH}_3)_2\text{CHCH}_3 + \text{CH}_3\text{CH}_2\text{OH}$

b) $(\text{CH}_3)_2\text{CH} \cdot \text{CH}_2\text{OH} + \text{C}_2\text{H}_6$





c) $(\text{CH}_3)_2\text{CHCH}_2\text{OH} + \text{C}_2\text{H}_5\text{I}$

d) $(\text{CH}_3)_2\text{CH} \cdot \text{CH}_2\text{I} + \text{CH}_3\text{CH}_2\text{OH}$

271. When glycerol is treated with a mixture of excess of conc. HNO_3 and H_2SO_4 , the compound formed is:

- a) Glycerol mononitrate b) Glycerol dinitrate c) Glycerol trinitrate d) acrolein

272. Identify Z in the following series,



273. 2 mole of ethanol are burnt. The amount of CO_2 obtained will be:

- a) 132 g b) 44 g c) 176 g d) 88 g

274. In which case, methyl *t*-butyl ether is formed?

- a) $(\text{C}_2\text{H}_5)_3\text{CONa} + \text{CH}_3\text{Cl}$ b) $(\text{CH}_3)_3\text{CONa} + \text{CH}_3\text{Cl}$
 c) $(\text{CH}_3)_3\text{CONa} + \text{C}_2\text{H}_5\text{Cl}$ d) $(\text{CH}_3)_2\text{CHONa} + \text{CH}_3\text{Cl}$

275. Grignard reagent reacts with HCHO to produce

- a) Secondary alcohol
 b) Anhydride
 c) Acid
 d) Primary alcohol

276. Alcohol is not used in making:

- a) Chloral b) Chloroform c) Benzene d) Acetaldehyde

277. Among the alkenes which one produces tertiary butyl alcohol on acid hydration?

- a) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ b) $\text{CH}_3\text{CH}=\text{CH}-\text{CH}_3$ c) $(\text{CH}_3)_2\text{C}=\text{CH}_2$ d) $\text{CH}_3-\text{CH}=\text{CH}_2$

278. Diethyl ether is soluble in:

- a) Water b) Dilute HCl c) Conc. H_2SO_4 d) Conc. KOH

279. Salicyl aldehyde is obtained when phenol is heated with CHCl_3 and aqueous NaOH . This reaction is known by which name?

- a) Carbyl amine reaction b) Hofmann's reaction
 c) Reimer-Tiemann reaction d) Kolbe-Schmidt reaction

280. The conversion of *m*-nitrophenol to resorcinol involves respectively

- a) Hydrolysis, diazotization and reduction b) Diazotization, reduction and hydrolysis
 c) Hydrolysis, reduction and diazotization d) Reduction, diazotization and hydrolysis

281. In Williamson's synthesis

- a) An alkyl halide is treated with sodium alkoxide b) An alkyl halide is treated with sodium
 c) An alcohol is heated with conc. H_2SO_4 at 130°C d) None of the above





282. $\text{C}-\text{O}-\text{C}$ angle would be maximum in

- a) $\text{CH}_3-\text{O}-\text{CH}_3$ b) $\text{CH}_3-\text{O}-\text{C}_2\text{H}_5$
 c) $\text{C}_2\text{H}_5-\text{O}-\text{C}_2\text{H}_5$ d) $(\text{CH}_3)_2\text{CH}-\text{O}-\text{CH}(\text{CH}_3)_2$


283. Ethers are very good solvent for which type of compounds?

- a) Lewis base b) Acids c) Lewis acid d) None of these

284. In which molecule, cleavage by HIO_4 is not observed?



285. The products formed in the following reaction,




- a) $\text{C}_6\text{H}_5\text{OH}$ and CH_3I b) $\text{C}_6\text{H}_5\text{I}$ and CH_3OH c) $\text{C}_6\text{H}_5\text{CH}_3$ and HOI d) C_6H_6 and CH_3OI

286. Acid catalysed hydration of alkenes except ethene leads to the formation of
 a) Mixture of secondary and tertiary alcohols b) Mixture of primary and secondary alcohols
 c) Secondary or tertiary alcohol d) Primary alcohol
287. Which of the following compounds when heated with CO at 150°C and 500 atm pressure in presence of BF₃ forms ethyl propionate?
 a) C₂H₅OH b) CH₃OCH₃ c) C₂H₅OC₂H₅ d) CH₃OC₂H₅
288. Which among the following compounds will give a secondary alcohol on reacting with Grignard reagent followed by acid hydrolysis?
 I. HCHO
 II. C₂H₅CHO
 III. CH₃COCH₃
 IV. HCOOC₂H₅

Select the correct answer using the codes given below.




- a) II only b) III only c) I and IV d) II and IV
289. When phenolic ether is heated with HI, it yields
 a) Alkyl halide + aryl halide + water



c) Alcohol + aryl halide

d) None of the above

290. The red coloured compound formed during Victor-meyer's test for ethanol is:



d) None of these

291. Picric acid is a stronger acid than acetic acid and benzoic acid. It contains



a) —SO₃H group

b) Two —COOH groups

c) Phenolic group

d)

292. Which will not form yellow precipitate on heating with an alkaline solution of iodine?



293. The cleavage of an aryl-alkyl ether with hydrogen halide will give:


a) A molecule each of an alkyl halide and water

b) A molecule each of an aryl halide and water

c) A molecule each of an alkyl halide, aryl halide and water

d) A molecule each of phenol and an alkyl halide

294. HBr reacts with CH₂=CH—OCH₃ under anhydrous conditions at room temperature to give:



295. In ether the active group is:

a) Oxygen

b) C₂H₅

c) Hydroxyl

d) None of these

296. The correct order of solubility of 1°, 2° and 3° alcohol in water is:





a) 3° > 2° > 1°

b) 1° > 2° > 3°

c) 3° > 1° > 2°


d) None of these





297. Maximum dehydration takes place in that of



298. The dehydration of butane-1-ol gives

- a) 1-butene as the main product
b) 2-butene as the main product
c) Equal amounts of 1-butene and 2-butene
d) 2-methyl propene



- a) 
b) 
c) 
d) 

300. When an ether is treated with P_2S_5 we get:

- a) Thio-alcohol
b) Thio-ester
c) Thio-ether
d) Thio-aldehyde

301. Order of reactivity of halogen acids towards an alcohol is

- a) $HCl > HBr > HI$
b) $HBr > HI > HCl$
c) $HI > HBr > HCl$
d) $HI > HCl > HBr$

302. In which of the following reactions the product is an ether?

- a) $C_6H_6 + CH_3COCl$ /anhydrous $AlCl_3$
b) $C_2H_5Cl + aq. KOH$
c) $C_6H_6 + C_6H_5COCl$ /anhydrous $AlCl_3$
d) $C_2H_5Cl + C_2H_5ONa$

303. The b.p. of alcohols are....than corresponding thiols.

- a) More
b) Less
c) Same
d) Either of these

304. Oxidation of 2-propanol by $K_2Cr_2O_7$ and dilute H_2SO_4 leads to the formation of:

- a) Propanal
b) Propanoic acid
c) Methanoic acid
d) Propanone

305. When phenol is treated with excess of bromine water, it gives

- a) *m*-bromophenol
b) *o*- and *p*-bromophenols
c) 2, 4-dibromophenol
d) 2, 4, 6-tribromophenol

306. An aqueous solution of ethyl alcohol:

- a) Turns blue litmus red
b) Turns red litmus blue
c) Does not affect the litmus colour
d) Decolourises litmus

307. Enzymes are:

- a) Living organisms
b) Dead organisms
c) Complex nitrogenous substances produced from living cells
d) None of the above

308. Which of the following is used as anaesthetic?

- a) $CHCl_3$
b) C_2H_5OH
c) $C_2H_5OC_2H_5$
d) $CHCl_3$ and $C_2H_5OC_2H_5$

309. Picric acid is

- a) 2, 4, 6-tribromophenol
b) *Sym*-trinitrophenol
c) trinitrophenol
d) 2, 4, 6-trinitrotoluene

310. The correct order of reactivity of hydrogen halides with ethyl alcohol is

- a) $HF > HCl > HBr > HI$
b) $HCl > HBr > HF > HI$
c) $HBr > HCl > HI > HF$
d) $HI > HBr > HCl > HF$

311. Denatured alcohol is

- a) Ethanol + methanol
b) Rectified spirit + methanol + naphtha
c) Undistilled ethanol
d) Rectified spirit

312. Which of the following reacts with water?

- a) CHCl_3 b) CCl_4 c) CCl_3CHO d) $\text{CH}_2\text{ClCH}_2\text{Cl}$





313. Formic acid is obtained when:

- a) $(\text{CH}_3\text{COO})_2\text{Ca}$ is heated with conc. H_2SO_4
 b) Calcium formate is heated with calcium acetate
 c) Glycerol is heated with oxalic acid
 d) Acetaldehyde is oxidized with $\text{K}_2\text{Cr}_2\text{O}_7$ and conc. H_2SO_4

314. Primary, secondary and tertiary alcohols are distinguished from one another by

- a) Ninhydrin test b) Tollen's reagent c) Lucas test d) Wittig reaction

315. Ethyl ester $\xrightarrow[\text{excess}]{\text{CH}_3\text{MgBr}}$ P. The product P will be

- a)  b) 
 c)  d) 




316. Metal alkoxides contain:

- a) Metal-carbon bond b) Metal-oxygen bond c) Metal-methyl bond d) None of these

317. 3-methyl-2-butanol on treatment with HCl gives predominantly:


- a) 2-chloro-2-methylbutane
 b) 2-chloro-3-methylbutane
 c) 2,2-dimethylpentane
 d) None of the above

318. Which reaction will occur?

- a)  b) 
 c)  d) None of these

319. No reacts rapidly with:

- a) 1° alcohol b) 2° alcohol c) 3° alcohol d) None of these



This reaction is called

- a) Reimer-Tiemann reaction b) Lederer-Manasse reaction
 c) Sandmeyer reaction d) Kolbe's reaction

321. By which of the following procedures can ethyl n-propyl ether be obtained?

- a) $\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{HBr}} \text{I} \xrightarrow[\text{ether}]{\text{Mg}} \text{II} \xrightarrow{\text{H}_2\text{O}} \text{III} \xrightarrow{\text{Na}} \text{CH}_3\text{CH}_2\text{Br}$
 b) $\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{HBr}} \text{I} \xrightarrow[\text{ether}]{\text{Mg}} \text{II} \xrightarrow[2. \text{H}_3\text{O}^+]{1. \text{CH}_2\text{O}} \text{III} \xrightarrow{\text{Na}} \text{CH}_3\text{CH}_2\text{Br}$
 c) $\text{C}_2\text{H}_5\text{OH} + \text{H}_2\text{SO}_4 \xrightarrow{140^\circ\text{C}}$
 d) $\text{C}_2\text{H}_5\text{OH} + \text{Conc. H}_2\text{SO}_4 \xrightarrow{180^\circ\text{C}} \text{I} \xrightarrow{\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}}$


322. Which of the following statements is wrong in case of ethoxyethane?

- a) It is used as anaesthetic
 b) It is inflammable
 c) Its dipole moment is zero
 d) It is soluble in conc. H_2SO_4

323. Which of the following alcohols is made by fermentation?

- a) Methanol b) Ethanol c) Glycerol d) Propanol

324. Correct acidic order of the following compounds is



- a) I > II > III b) III > I > II c) II > III > I d) I > III > II

325. How many isomers of $C_5H_{11}OH$ will be primary alcohols?

- a) 5 b) 4 c) 2 d) 3

326. Glycerol is oxidised by bismuth nitrate to produce

- a) Oxalic acid b) Mesoxalic acid c) Glyceric acid d) Glyoxalic acid


327. The alcohol that produces turbidity immediately with $ZnCl_2/conc. HCl$ at room temperature





- a) 1-hydroxy butane b) 2-hydroxy butane
 c) 2-hydroxy-2-methyl propane d) 1-hydroxy-2-methyl propane

328. The formula for allyl alcohol is:

- a) $CH_3-CH=CHCl$ b) $CH_2=CHCH_2OH$ c) $CH_2ClCH_2CH_3$ d) None of these

329.



- a)  b)  c)  d) 

330. The compound that will react most readily with NaOH to form methanol is:

- a) $(CH_3)_4N^+I^-$ b) CH_3OCH_3 c) $(CH_3)_3S^+I^-$ d) $(CH_3)_3C \cdot Cl$

331. Ethylene reacts with 1% cold alkaline $KMnO_4$ to give:

- a) Oxalic acid b) Acetone c) Formaldehyde d) Glycol

332. In the Lucas test of alcohols, appearance of cloudiness is due to the formation of

- a) Aldehydes b) Ketones c) Acid chlorides d) Alkyl chlorides

333. Tertiary alcohol is obtained when Grignard reagent reacts with:

- a) Acetone b) Butanone c) Propanone d) All of these

334. On conversion into the Grignard reagent followed by treatment with absolute ethanol, how many isomeric alkyl chlorides would yield 2-methylbutane?

- a) 2
 b) 3
 c) 4
 d) 5

335. Ether on reacting with P_2S_5 form

- a) Diethyl sulphide b) Thioalcohol c) Thioether d) Thioaldehyde

336. The best reagent to convert pent-3-en-2-ol into pent-3-en-2-one is:

- a) Acidic $KMnO_4$
 b) Alkaline $K_2Cr_2O_7$
 c) Chromium anhydride in glacial acetic acid
 d) Pyridinium chlorochromate

337. For one mole of glycerol, how many mole of acetyl chloride are required for complete acetylation?

- a) One b) Two c) Three d) Four

338. In the reaction involving C—OH bond, in alcohols the order of reactivity is:

- a) $1^\circ > 2^\circ > 3^\circ$ b) $3^\circ > 2^\circ > 1^\circ$ c) $2^\circ > 3^\circ > 1^\circ$ d) None of these


339. Which is not correct?





- a) Phenol is more acidic than acetic acid. b) Ethanol is less acidic than phenol.
 c) Ethanol has higher boiling point than ethane. d) Ethane is non-linear molecule.

340. Under drastic conditions all the alcohols can be oxidized to carboxylic acids but the following alcohols give carboxylic acids having same number of carbon atoms:




- a) Primary b) Secondary c) Tertiary d) None of these

341. The product *A* is



- a)  b)  c)  d) 

342. Glycol is prepared industrially by the following reactions:



- d) None of the above

343. Scientific aspect of fermentation was first studied by:

- a) Pasteur b) Brot c) Buchner d) Liebig

344. Ethyl alcohol is also known as:

- a) Spirit of wine b) Methyl carbinol c) Grain alcohol d) All of these

345. Decreasing order of boiling points of *n*-pentanol (*A*), *n*-pentane (*B*), 3-pentanol (*C*) and 2,2-dimethyl propanol (*D*) is :

- a) *A, C, D, B* b) *B, D, C, A* c) *C, A, D, B* d) None of these

346. CH_3COOH reacts rapidly with:

- a) $\text{CH}_3\text{CH}_2\text{OH}$ b) $(\text{CH}_3)_2\text{CHOH}$ c) $(\text{CH}_3)_3\text{COH}$ d) All of these

347. Reaction of *t*-butyl bromide with sodium methoxide produces:

- a) Isobutane b) Isobutylene c) Sodium *t*-butoxide d) *t*-butyl methyl ether

348. Which of the following reactions can be used for the preparation of tert. butylmethyl ether?

- a) $\text{CH}_3\text{Br} + (\text{CH}_3)_3\text{CO}^- \text{Na}^+ \rightarrow$ b) $(\text{CH}_3)_3\text{CCl} + \text{CH}_3\text{O}^- \text{Na}^+ \rightarrow$
 c) $(\text{CH}_3)_3\text{OH} + \text{CH}_3\text{Cl} \rightarrow$ d) $(\text{CH}_3)_3\text{CCl} + \text{CH}_3\text{OH} \rightarrow$

349. Alcohols cannot be prepared from

- a) $\text{C}_2\text{H}_5\text{Br} + \text{aq. KOH} \rightarrow$ b) $(\text{CH}_3)_2\text{C} = \text{O} \xrightarrow{\text{LiAlH}_4}$ c) $\begin{array}{c} \text{O} \\ || \\ \text{CH}_3 - \text{C} - \text{OCH}_3 \end{array} \xrightarrow{\text{Na/EtOH}}$ d) $\text{CH}_3\text{CH}_2\text{Cl} \xrightarrow{\text{H}_2\text{O}}$




350. Alcohols of low molecular weight are:

- a) Soluble in water
 b) Soluble in water on heating
 c) Insoluble in all solvents
 d) Soluble in all solvents

351. $\text{CH}_3\text{CH} = \text{CH} - \text{CH} - \text{CH}_3$



$\xrightarrow[\text{reagent}]{\text{Jones}}$? product is



352. 23 g of sodium react with CH_3OH to give:


a) 1 mole of O_2

b) $1/2$ mole of H_2





c) 1 mole of H_2

d) None of these





353. In the reaction for dinitration



The major dinitrated product X is



354. In the following compounds the order of acidic strength is



(I)

(II)

(III)

(IV)

a) III > IV > I > II

b) I > IV > III > II

c) II > I > III > IV

d) IV > III > I > II

355. Diethyl ether may behave as:

a) Lewis acid

b) Lewis base

c) Oxidising agent

d) Reducing agent

356. For drying ether sodium metal can be used, but it cannot be used for drying ethyl alcohol because:

a) Na is very reactive

b) Ether reacts easily with Na

c) Ethyl alcohol reacts with sodium metal

d) None of the above

357. Saccharification is the process of conversion of:

a) Sugar solution into alcohol

b) Alcohol into starch

c) Starch into alcohol

d) Starch into alcohol






358. $\text{R}-\text{CH} = \text{CH}_2$ reacts with B_2H_6 in presence of H_2O_2 to give:


a) RCOCH_3

b) $\text{RCHOHCH}_2\text{OH}$

c) $\text{RCH}_2\text{CH}_2\text{OH}$

d) RCH_2CHO

359. Sodium phenoxide reacts with CO_2 at 400 K and 4.7 atm pressure to give
- a) Catechol b) Salicylaldehyde c) Sodium salicylate d) Benzoic acid
360. The reaction of *iso*-propylbenzene with oxygen in the presence of a catalytic amount of HBr followed by treatment with an acid gives phenol. The reaction proceeds through the intermediate formation of
- a) $\text{HO}\ddot{\text{O}}$ b) $\text{C}_6\text{H}_5-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}\cdot$ c) $\text{C}_6\text{H}_5-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{O}-\text{O}\cdot$ d) All of these
361. Product formed when HCHO is heated with KOH (aq):
- a) CH_4 b) CH_3CHO c) CH_3OH d) C_2H_2
362. Diacetone alcohol is obtained by the reaction of:
- a) Acetone and ethanol
b) Acetone and conc. H_2SO_4
c) Acetone and $\text{Ba}(\text{OH})_2$
d) Acetone and $\text{Al}(\text{OH})_3$
363. The general formula of ether is:
- a) $\text{R}-\text{CHO}$ b) $\text{R}-\text{CO}-\text{R}'$ c) $\text{R}-\text{O}-\text{R}'$ d) $\text{R}-\text{COOR}'$
364. The enzyme pepsin hydrolyses:
- a) Proteins to amino acids
b) Fats to fatty acids
c) Glucose to ethyl alcohol
d) Polysaccharides to monosaccharides
365. $\text{CH}_3\text{CH}_2\text{OH}$ convert into CH_3CHO in the presence of
- a) $\text{Na}_2\text{Cr}_2\text{O}_7$ and NaOH b) $\text{Na}_2\text{Cr}_2\text{O}_7$ and dil. H_2SO_4
c) NaOH d) Fe in presence of NaOH
366. Which of the following combinations can be used to synthesise ethanol?
- a) CH_3MgI and CH_3COCH_3 b) CH_3MgI and $\text{C}_2\text{H}_5\text{OH}$
c) CH_3MgI and $\text{CH}_3\text{COOC}_2\text{H}_5$ d) CH_3MgI and HCHO
- 367.
- 
- Product is
- a)  b)  c)  d) 
368. The boiling point of ethyl alcohol is much higher than that of dimethyl ether and $\text{C}_2\text{H}_5\text{SH}$, though both have the same molecular weight. The reason for this is:
- a) Ether is insoluble in water
b) Methyl groups are attached to oxygen in ether
c) Dipole moment of ethyl alcohol is less
d) Ethyl alcohol shows hydrogen bonding
369. Acetylene and formaldehyde interact in the presence of copper acetylide as a catalyst to furnish the compound:
- a) Butyne-1, 4-diol b) Butyne-2 c) Ethylene-1, 4-diol d) None of these

370. An unknown compound 'D' first oxidised to aldehyde and then acetic acid by a dilute solution of $K_2Cr_2O_7$ and H_2SO_4 . The compound 'D' is
 a) CH_3OH b) C_2H_5OH c) CH_3CH_2COOH d) CH_3CH_2CHO
371. Glycerol on oxidation with Fenton's reagent produces:
 a) Glyceraldehyde
 b) Dihydroxy acetone
 c) Tartonic acid
 d) Glyceraldehyde and dihydroxy acetone
372. An organic compound C_3H_6O neither gives precipitate with semicarbazide nor reacts with sodium. It could be
 a) CH_3CH_2CHO b) CH_3COCH_3 c) $CH_2 = CHCH_2OH$ d) $CH_2 = CHOCH_3$
373. Which one among the following is Williamson's synthesis?
 a) $\begin{array}{l} CH_3 \\ \diagdown \\ C=O \\ \diagup \\ CH_3 \end{array} \xrightarrow[Conc. HCl]{Zn-Hg} CH_3-CH_2-CH_3$
 b) $CH_3-CHO \xrightarrow[-H_2O]{Dil. NaOH} CH_3-CH=CH-CHO$
 c) $C_2H_5I + C_2H_5ONa \rightarrow C_2H_5 \cdot O \cdot C_2H_5 + NaI$
 d) $HCHO \xrightarrow{NaOH} HCOONa + CH_3OH$
374. Which compound is capable of strong hydrogen bonding?
 a) C_4H_9OH b) C_3H_7OH c) C_2H_5OH d) $C_5H_{11}OH$
375. $CH \equiv CH \xrightarrow{O_3/NaOH} X \xrightarrow{Zn/CH_3COOH} Y$ is:
 a) CH_2OH-CH_2OH b) CH_3CH_2OH c) CH_3COOH d) CH_3OH
376. Which of the following statements is not correct?
 a) All alcohols are miscible with water b) Only lower alcohols are miscible with water
 c) All alcohols are not poisonous d) Methanol is not poisonous
377. The major product of the following reaction is:

 a) A hemiacetal
 b) An acetal
 c) An ether
 d) An ester
378. Widespread deaths due to liquor poisoning occurs due to presence of:
 a) Lead compounds in liquor
 b) Methyl alcohol in liquor
 c) Ethyl alcohol in liquor
 d) Carbonic acid in liquor
379. An alcohol produced during the manufacture of soap is:
 a) Butanol b) Glycerol c) Ethanol d) Ethylene glycol
380. Which of the following reactions gives an dialkyl oxonium salt?
 a) Ethyl alcohol + sodium metal
 b) Diethyl ether + hydrochloric acid
 c) Tertiary amine + alkyl halide
 d) Nitromethane + sodium metal
381. The reaction of *neo*-pentyl alcohol with concentrated HCl gives
 a) *neo*-pentyl chloride b) 2-chloro-2-methylbutane

c) 2-methyl-2-butene

d) A mixture of *neo*-pentyl chloride and 2-methyl-2-butene


382. RCH_2CH_2OH can be converted to RCH_2CH_2COOH by the following sequence of steps





a) PBr_3, KCN, H_3O^+

b) $PBr_3, KCN, H_2/P^+$

c) KCN, H_3O^+

d) HCN, PBr_3, H_3O^+

383.  $\xrightarrow[CH_2Cl_2]{mCPBA}$ A, A is



384. When phenyl magnesium bromide reacts with *t*-butanol, the product would be

a) Benzene

b) Phenol

c) *t*-butyl benzene

d) *t*-butyl phenyl ether

385. Which of the following is not cleaved by HIO_4 ?

A. Glycerol

B. Glycol

C. Propan-1,3-diol

D. Methoxy-2-propanol

a) A, B, C, D

b) A, B

c) B, C

d) C, D

386. Ethyl propanoate on reduction with $LiAlH_4$ yields:

a) Methanol

b) Ethanol and propanol

c) Propane

d) Mixture of ethanol and methanol

387. When acetyl chloride is reduced with $LiAlH_4$, the product formed is:

a) Methyl alcohol

b) Ethyl alcohol

c) Acetaldehyde

d) Acetone

388. The correct order of acid strength of the following compounds is

V. Phenol

VI. *p*-cresol

VII. *m*-nitrophenol

VIII. *p*-nitrophenol

a) VIII > II > I > IV

b) IV > III > I > II

c) II > IV > I > III

d) I > II > IV > III

389. Alkyd resins, made of glycerol are used:

a) As substitute for white chalk

b) Instead of alkanes

c) For paints and coatings

d) For making alcohol

390. Which reagent is more effective to convert but-2-enal to but-2-enol?

a) $KMnO_4$

b) $NaBH_4$

c) H_2/Pt

d) $K_2Cr_2O_7/H_2SO_4$

391. An organic compound A containing C, H and O has a pleasant odour with boiling point of $78^\circ C$. On boiling A with concentrated H_2SO_4 , a colourless gas is produced which decolourises bromine water and alkaline $KMnO_4$. The organic liquid A is






a) C_2H_5Cl

b) $C_2H_5COOCH_3$

c) C_2H_5OH

d) C_2H_6

392. Identify (X) in the sequence:



393. Phenol on reaction with CHCl_3 and NaOH give benzaldehyde. Intermediate of this reaction is


- a) Carbocation b) Carbanion c) Radical d) Carbene

394. Increasing order of acid strength among *tert.* butanol, isopropanol and ethanol is:

- a) Ethanol, isopropanol, *tert.* butanol
 b) *tert.* butanol, isopropanol, ethanol
 c) Isopropanol, *tert.* butanol, ethanol
 d) *tert.* butanol, ethanol, isopropanol

395. A neutral compound gives colour with ceric ammonium nitrate. It suggests that the compound has:

- a) Alcohol gp. b) Aldehyde gp. c) Ether gp. d) Ketone gp.



In the above reaction A is

- a) $\text{CH}_3-\overset{\text{OH}}{\underset{\text{OH}}{\text{C}}}=\text{CH}_2$ b) $\text{CH}_2=\overset{\text{OH}}{\underset{\text{OH}}{\text{C}}}-\text{OCH}_3$ c) $\text{CH}_2 = \text{CHOH}$ d) None of these





397. Which compound will have highest boiling point?

- a) CH_4 b) CH_3OH c) $\text{C}_2\text{H}_5\text{OH}$ d) HCHO

398. What is formed when glycerol reacts with excess of HI?

- a) $\begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{CHI} \\ | \\ \text{CH}_2\text{OH} \end{array}$ b) $\begin{array}{c} \text{CH}_2 \\ || \\ \text{CH} \\ | \\ \text{CH}_2\text{I} \end{array}$ c) $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CHI} \\ | \\ \text{CH}_3 \end{array}$ d) $\begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{C}=\text{O} \\ | \\ \text{CH}_3 \end{array}$

399. Which of the following is not soluble in NaHCO_3 solution?

- a)  b)  c)  d) 


400. Pyroligneous acid doesn't contain




- a) Acetic acid b) $\text{C}_2\text{H}_5\text{OH}$ c) CH_3OH d) CH_3COCH_3

401. Power alcohol is a mixture of petrol and alcohol in the ratio:

- a) 4 : 1 b) 1 : 4 c) 2 : 1 d) 1 : 2

402. The final product obtained in the reaction,



- a)  b) 
 c)  d) None of the above

403. Which one of the following gases is liberated when ethyl alcohol is heated with methyl magnesium iodide?

- a) Methane b) Ethane c) Carbon dioxide d) Propane

404. Phenol \xrightarrow{X} forms a tribromo derivative "X" is

- a) Bromine in benzene b) Bromine in water
 c) Potassium bromide solution d) Bromine in carbon tetrachloride at 0°C


405. Phenol is more acidic than alcohol because

- a) Phenol is more soluble in polar solvents b) Alcohol does not lose hydrogen atom

- c) Phenoxide ion is stabilised by resonance d) Phenoxide ion doesn't exhibit resonance
406. Which of the following is the best method for making *iso*-propylmethyl ether?
- a) $\text{CH}_3\text{I} + (\text{CH}_3)_2\text{CHOH} \rightarrow$ b) $\text{CH}_3\text{I} + (\text{CH}_3)_2\text{CHO}^- \rightarrow$
 c) $(\text{CH}_3)_2\text{CHI} + \text{CH}_3\text{O}^- \rightarrow$ d) $(\text{CH}_3)_2\text{CHCl} + \text{CH}_3\text{OH} \rightarrow$
407. If the boiling point of ethanol (molecular weight=46) is 78°C , what is the boiling point of diethyl ether? (molecular weight=74)
- a) 100°C b) 78°C c) 86°C d) 34°C
408. An organic compound *A* reacts with PCl_5 to give *B*. The compound *B* with sodium metal gives *n*-butane. Thus, *A* and *B* are:
- a) $\text{C}_2\text{H}_5\text{OH}$ and $\text{C}_2\text{H}_5\text{Cl}$
 b) $\text{C}_2\text{H}_5\text{Cl}$ and $\text{C}_2\text{H}_5\text{ONa}$
 c) $\text{C}_3\text{H}_7\text{OH}$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCl}$
 d) $\text{C}_4\text{H}_9\text{OH}$ and $\text{C}_4\text{H}_9\text{OCl}$
409. Acetic acid is obtained from ethyl alcohol by the process of:
- a) Distillation b) Reduction c) Fermentation d) Dehydration
410. Intermolecular dehydration of alcohols gives:
- a) Alkenes b) Ketones c) Alkynes d) Ethers
411. Glycerol on warming with excess of HI :
- a) 2-iodopropane b) 1-iodopropane c) 1,2,3-tri-iodopropane d) None of these
412. Cumene process is the most important commercial method for the manufacture of phenol. Cumene is
- a) 1-methyl ethyl benzene b) Ethyl benzene
 c) Vinyl benzene d) Propyl benzene
413. Which of the following alcohols cannot be oxidized by potassium dichromate in the presence of sulphuric acid?


- a) $\text{CH}_3\text{CH}_2\text{OH}$ b) $\text{C}_6\text{H}_5\text{OH}$ c) $\text{C}_6\text{H}_5\text{CHOHCH}_3$ d) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}} - \text{OH}$

414. Which of the following is stable compound?
- a) $\text{CCl}_3\text{CH}(\text{OH})_2$ b) $\text{CH}_2=\text{CHOH}$ c) $\text{CH}_3-\text{CH}(\text{OH})_2$ d) $\text{HC}(\text{OH})_3$



The major product obtained in this reaction is


- a) $(\text{CH}_3)_2\text{CHCH} = \text{CH}_2$
 b) $(\text{CH}_3)_2\text{C} = \text{CH} - \text{CH}_3$
 c) 1 : 1 mixture of (a) and (b)
 d) None of the above



Product obtained is




- a) $\begin{array}{c} \text{R} \\ | \\ \text{R}' - \text{C} - \text{CH}_2\text{OH} \\ | \\ \text{R}'' \end{array}$ b) $\begin{array}{c} \text{R}' \\ | \\ \text{RCH}_2 - \text{C} - \text{OH} \\ | \\ \text{R}'' \end{array}$ c) $\begin{array}{c} \text{R} \\ | \\ \text{R}'\text{CH}_2 - \text{C} - \text{OH} \\ | \\ \text{R}'' \end{array}$ d) $\begin{array}{c} \text{R} \\ | \\ \text{R}''\text{CH}_2 - \text{C} - \text{OH} \\ | \\ \text{R}' \end{array}$

417. The reaction involved in the oil of winter green test is salicylic acid $\xrightarrow[\text{Conc. H}_2\text{SO}_4]{\Delta}$ product. The product is treated with Na_2CO_3 solution. The missing reagent in the above reaction is
- a) Phenol b) NaOH c) Ethanol d) Methanol
418. An example of a compound with functional group $-\text{O}-$ is:
- a) Acetic acid b) Methyl alcohol c) Diethyl ether d) Acetone
419. Phenol gives characteristic colouration with
- a) Iodine solution b) Bromine water
c) Aqueous FeCl_3 solution d) Ammonium hydroxide
420. The correct order of the ease with which primary, secondary and tertiary alcohols can be dehydrated using concentrated H_2SO_4 is :
- a) Tertiary > secondary > primary
b) Primary > secondary > tertiary
c) Secondary > tertiary > primary
d) Secondary > primary > tertiary
421. Which are explosives?
- a) Wood pulp (dynamite)
b) Cellulose nitrate (blasting gelatin)
c) Gun cotton or cellulose nitrate and Vaseline (cordite)
d) All of the above
422. Some time explosion occurs while distilling ethers. It is due to the presence of
- a) Oxide b) Ketones c) Aldehyde d) Peroxides
423. Acidity of phenol is due to
- a) Hydrogen bonding b) Phenolic group
c) Benzene ring d) Resonance stabilisation of its anion
424. Glycerol on reacting with sodium gives:
- a) Disodium glycerollate
b) Monosodium glycerollate
c) Trisodium glycerollate
d) None of the above
425. The compound which reacts fastest with Lucas reagent at room temperature is
- a) 1-butanol b) 2-butanol c) 2-methylpropanol d) 2-methylpropan-2-ol
426. Mild oxidation of glycerol with $\text{H}_2\text{O}_2/\text{FeSO}_4$ gives
- a) Glyceraldehyde
b) Dihydroxy acetone
c) Both (a) and (b)
d) None of the above
427. To prepare 2-propanol from CH_3MgI , the other chemical required is:
- a) HCHO b) CH_3CHO c) $\text{C}_2\text{H}_5\text{OH}$ d) CO_2
428. The first oxidation product of primary alcohol is:
- a) A ketone b) An ester c) An aldehydes d) A hydrocarbon
429. Phenol is soluble in water because
- a) Of weak hydrogen bonding between phenol and water molecules
b) Of intermolecular hydrogen bonding between phenol molecules
c) If has a higher boiling point than that of water
d) None of the above
430. Consider the following reaction,




product X is

- a) Picric acid b) Styphnic acid c) Salicylic acid d) Benzoic acid
431. Glycerol on treatment with oxalic acid at 110°C forms:
 a) Formic acid b) CO₂ and CO c) Allyl alcohol d) glycol
432. At 530 K, glycerol reacts with oxalic acid to produce
 a) Allyl alcohol b) Formic acid c) Glyceraldehydes d) Formaldehyde
433. Absolute alcohol is prepared from rectified spirit by:
 a) Fractional distillation
 b) Steam distillation
 c) Azeotropic distillation
 d) Vacuum distillation
434. Williamson's synthesis is used to prepare
 a) Diethyl ether b) Acetone c) PVC d) Bakelite
435. Anisole can be prepared by the action of methyl iodide on sodium phenate. The reaction is called
 a) Wurtz's reaction b) Williamson's reaction
 c) Fittig's reaction d) Etard's reaction
436. When *o*- or *p*-phenol sulphonic acid is treated with bromine water, the product formed is
 a) 2, 4-dibromophenol b) 2, 4, 6-tribromophenol
 c) 3-bromophenol boric acid d) 3, 5-dibromophenol
437. Esterification of alcohols involves:
 a) H of alcohol and OH of acid
 b) OH of alcohol and H of acid
 c) OH of alcohol and OH of acid
 d) H of alcohol and H of acid
438. An organic liquid A containing C, H and O has a pleasant odour with a b.p. of 78°C. On boiling A with conc. H₂SO₄ a colourless gas is produced which decolourises bromine water and alkaline KMnO₄. One mole of this gas also takes one mole of H₂. The organic liquid A is:
 a) C₂H₅Cl b) C₂H₅CHO c) C₂H₆ d) C₂H₅OH
439. In the presence of an acid catalyst, two alcohol molecules will undergo dehydration to give:
 a) Ester
 b) Anhydride
 c) Ether
 d) Unsaturated hydrocarbon
440. Complete combustion of ether gives:
 a) C₂H₅OH b) CO₂ and H₂O c) C₂H₄ d) C₂H₂
441. Product is



d) Both (a) and (b)



In above reactions the role of Cl_2 in step-1 and step-2 respectively is

a) Oxidation, chlorination

b) Reduction, chlorination

c) Oxidation, addition

d) Reduction, substitution

443. An enzyme which brings about the conversion of starch into maltose is known as:


a) Maltase

b) Zymase


c) Invertase

d) Diastase


444. Strength of acidity is in order




(I)



(II)



(III)



(IV)

a) II > I > III > IV

b) III > IV > I > II

c) I > IV > III > II

d) IV > III > I > II

445. Ethyl alcohol is denatured by:


a) Methanol and formic acid

b) KCN

c) CH_3OH and C_6H_6

d) CH_3OH and pyridine

446. For the sequence of reaction,



The compound A in the sequence is

a) 2-butanone

b) Acetaldehyde

c) Acetone

d) Propanal

447. A compound with molecular formula $\text{C}_4\text{H}_{10}\text{O}_3$ is converted by the action of acetyl chloride to a compound with molecular weight 190. The original compound has:





a) One OH group

b) Two OH groups

c) Three OH groups

d) No OH group

448. The main product of the following reaction is



449. Which of the following compound is oxidised to prepare methyl ethyl ketone?

a) 2-propanol

b) 1-butanol

c) 2-butanol

d) Ter-butyl alcohol

450. The value of $\text{C}-\text{O}-\text{C}$ angle in ether molecule is:

a) 180°

b) 150°

c) 90°

d) 110°

451. What amount of bromine will be required to convert 2 g of phenol into 2, 4, 6-tribromo phenol?

- a) 4.00 b) 6.00 c) 10.22 d) 20.44

452. Chloroform which is a good solvent for aromatic impurities is:

- a) Dichloro dimethyl ether
b) Dichlorodiethyl ether
c) Mono chloro ether
d) Diethyl ether

453. The characteristic group of secondary alcohol is:

- a) $-\text{CH}_2\text{OH}$
b) >CHOH
c) >COH
d) $-\text{COOH}$


454. The compound on dehydrogenation gives a ketone. The original compound is

- a) Primary alcohol b) Secondary alcohol c) Tertiary alcohol d) Carboxylic acid

455. 1-phenyl ethanol can be prepared from benzaldehyde by the action of:

- a) CH_3Br b) CH_3Br and AlBr_3 c) CH_3I , Mg and HOH d) $\text{C}_2\text{H}_5\text{I}$ and Mg

456. Which one of the following will most readily be dehydrated in acidic conditions?



457. On reduction with LiAlH_4 , a ketone yields:

- a) Primary alcohol b) Secondary alcohol c) Tertiary alcohol d) All of these


458. The decreasing order of boiling points of 1° , 2° , 3° alcohol is:

- a) $1^\circ > 2^\circ > 3^\circ$ b) $3^\circ > 2^\circ > 1^\circ$ c) $2^\circ > 1^\circ > 3^\circ$ d) None of these

459. The formula for vinyl alcohol is:

- a) $\text{CH}_2=\text{CHCH}_2\text{OH}$ b) $\text{C}_6\text{H}_5\text{CHOHCH}_3$ c) $\text{CH}_2=\text{COHCH}_3$ d) $\text{CH}_2=\text{CHOH}$

460. Consider the reaction,



A and B respectively are

- a) 1, 2-epoxycyclohexane, *trans*-2-bromocyclohexanol b) 1, 2-epoxycyclohexane, *cis*-2-bromocyclohexanol
c) *trans*-2-bromocyclohexanol 1,2-epoxyethane d) *cis*-2-bromocyclohexanol 1,2-epoxyethane

461. Alcoholic fermentation of sugar gives 3% glycerol. The yield can be increased to 25% if fermentation is made in presence of:

- a) Na_2SO_4 b) Na_3PO_4 c) Na_2S d) None of these

462. The reaction,



is called :

- a) Fischer-Speier esterification

- b) Clemmensen condensation
 c) Claisen condensation
 d) None of the above
463. When isopropyl alcohol vapours are passed over heated copper it gives:
 a) Acetone b) Ethyl alcohol c) Methyl alcohol d) Acetaldehyde
464. Glycol on oxidation with....gives oxalic acid.
 a) Acidic KMnO_4 b) Acidic $\text{K}_2\text{Cr}_2\text{O}_7$ c) Nitric acid d) HIO_4
465. When compound X is oxidised by acidified potassium dichromate, compound Y is formed. Compound Y on reduction with LiAlH_4 gives X . X and Y respectively are
 a) $\text{C}_2\text{H}_5\text{OH}$, CH_3COOH
 b) CH_3COCH_3 , CH_3COOH
 c) $\text{C}_2\text{H}_5\text{OH}$, CH_3COCH_3
 d) CH_3CHO , CH_3COCH_3
466. The reaction of ethanol with H_2SO_4 does not give:
 a) C_2H_4 b) $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$ c) C_2H_2 d) $\text{C}_2\text{H}_5\text{HSO}_4$
467. Lucas reagent produces cloudiness immediately with:
 a) n -butanol b) Isopropanol c) n -propanol d) Tertiary butanol
468. Primary alcohols can be obtained from the reaction of the RMgX with:
 a) HCHO b) H_2O c) CO_2 d) CH_3CHO
469. The major product obtained on interaction of phenol with sodium hydroxide and carbon dioxide is
 a) Benzoic acid b) Salicylaldehyde c) Salicylic acid d) Phthalic acid
470. Chlorobenzene $\xrightarrow[X]{\text{Reaction}}$ Phenol $\xrightarrow[Y]{\text{Reaction}}$ Salicylaldehyde
 X and Y reactions are respectively.....
 a) Fries rearrangement and Kolbe-Schmidt b) Cumene and Reimer-Tiemann
 c) Dow and Reimer-Tiemann d) Dow and Friedel-Craft
471. Phenol $\xrightarrow{\text{NaNO}_2/\text{H}_2\text{SO}_4}$ B $\xrightarrow{\text{H}_2\text{O}}$ C $\xrightarrow{\text{NaOH}}$ D
 Name of the above reaction is
 a) Liebermann's reaction b) Phthalein fusion test
 c) Reimer-Tiemann reaction d) Schotten-Baumann reaction
472. Vinyl carbinol is:
 a) $\text{HOH}_2\text{C}-\text{CH}=\text{CH}_2$ b) $\text{CH}_3\text{C}(\text{OH})=\text{CH}_2$ c) $\text{CH}_3-\text{CH}=\text{CH}-\text{OH}$ d) $\begin{array}{c} \text{CH}_3-\text{C}=\text{CH}_2 \\ | \\ \text{CH}_3\text{OH} \end{array}$
473. Choose the incorrect statement
 a) Ordinary ethyl alcohol is known as rectified spirit
 b) The alcohol sold in the market for polishing etc, is known as methylated spirit
 c) Absolute alcohol is 100% ethanol
 d) Power alcohol is 100% ethanol
474. The reaction of ethanol with concentrated H_2SO_4 at room temperature gives
 a) $\text{CH}_3\text{CH}_2\text{OH}_2^+\text{HSO}_4^-$ b) $\text{CH}_3\text{CH}_2\text{OSO}_2\text{OH}$
 c) $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ d) $\text{H}_2\text{C}=\text{CH}_2$
475. $\begin{array}{c} \text{H}_2\text{C}-\text{CH}_2 \\ \diagdown \quad / \\ \text{O} \end{array} \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) CH}_3\text{MgCl}} X \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) CH}_3\text{MgCl}} X$
 The product obtained in this reaction is
 a) $\text{CH}_3\text{CH}_2\text{OH}$ b) $(\text{CH}_3)_2\text{CHOH}$
 c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ d) $\text{HO}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$
476. When ethylene glycol is heated with a mixture of concentrated HNO_3 and concentrated H_2SO_4 , it produces
 a) COOH b) $\text{CO}_2 + \text{H}_2$ c) CH_2ONO_2 d) CH_2ONO_2




477. Cyclohexanol on reaction with PBr_3 in presence of pyridine gives

- a) Bromocyclohexane b) Bromocyclohexane c) 1-bromocyclohexanol d) None of these





478. On treatment with a concentrated solution of zinc chloride in concentrated HCl at room temperature, an alcohol immediately gives, an oily product. The alcohol can be

- a) $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$
b) $\text{CH}_3\text{CHOHCH}_3$



- d) Any of these






479. The most unlikely representation of resonance structures of *p*-nitrophenoxide ion is



480. Ethylene glycol gives oxalic acid on oxidation with

- a) Acidified $\text{K}_2\text{Cr}_2\text{O}_7$ b) Acidified KMnO_4 c) Alkaline KMnO_4 d) Periodic acid

481. In the reaction,



482. Diethyl ether may be regarded as anhydride of:

- a) $\text{C}_2\text{H}_5\text{COOH}$ b) $\text{C}_2\text{H}_5\text{OH}$ c) $\text{C}_2\text{H}_5\text{CHO}$ d) $\text{C}_2\text{H}_5\text{COOC}_2\text{H}_5$

483. Glycol reacts with PCl_3 and gives ethylene dichloride. What will be the product, if it reacts with P + I_2 ?

- a) Ethylene iodide b) Ethylene iodohydrin c) Ethylene d) None of these

484. Methyl alcohol reacts with phosphorus trichloride to form:

- a) Methane b) Methyl chloride c) Acetyl chloride d) Dimethyl ether

485. Arrange the following in order of decreasing acidic strength. *p*-nitrophenol (I), *p*-cresol (II), *m*-cresol (III), phenol (IV)

- a) $\text{I} > \text{II} > \text{III} > \text{IV}$ b) $\text{IV} > \text{III} > \text{II} > \text{I}$ c) $\text{I} > \text{III} > \text{II} > \text{IV}$ d) $\text{III} > \text{II} > \text{I} > \text{IV}$

486. A diazonium chloride reacts with ϕOH to give an azodye. The reaction is called

- a) Diazotisation b) Condensation c) Coupling d) Reduction

487. Which alcohol is most acidic?

- a) Methanol b) Ethanol c) Isopropyl alcohol d) *t*-butyl alcohol

488. Which reagent can distinguish $\text{C}_2\text{H}_5\text{OH}$ and ϕOH ?

- a) SOCl_2 b) CH_3COCl c) $(\text{CH}_3\text{CO})_2\text{O}$ d) CH_3COOH


489. *Iso*-butyl alcohol $\xrightarrow{\text{P/I}_2} \xrightarrow{\text{AgNO}_2} \xrightarrow{\text{HNO}_2} \xrightarrow{\text{NaOH}} \text{A}$

True statement about A is

- a) Blue coloured solution
c) Red precipitate
490. Acetone on reduction gives:
a) CH_3COOH b) CH_3CHO c) $\text{C}_2\text{H}_5\text{OH}$ d) $(\text{CH}_3)_2\text{CHOH}$
491. Sodium ethoxide and ethyl chloride on heating will give:
a) Ether b) Ethyl alcohol c) Acetaldehyde d) Acetic acid
492. Pinacol is
a) 3-methylbutan-2-ol b) 2, 3-dimethyl-2, 3-butanediol
c) 2, 3-dimethyl-2-propanone d) None of the above
493. The product in the reaction is:
$$\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{P} + \text{I}_2} \text{A} \xrightarrow[\text{Ether}]{\text{Mg}} \text{B} \xrightarrow{\text{HCHO}} \text{C} \xrightarrow{\text{H}_2\text{O}} \text{D}$$



a) Propanal b) Butanal c) *n*-butanol d) *n*-propanol
494. In esterification of an acid, the other reagent is:
a) Aldehyde b) Alcohol c) Amine d) Water
495. $\text{C}_2\text{H}_5\text{OH}$ and $\text{C}_2\text{H}_5\text{OH}$ can be distinguished by
a) $\text{Br}_2 + \text{H}_2\text{O}$ b) FeCl_3 c) $\text{I}_2 + \text{NaOH}$ d) Both (b) and (c)
496. Identify (Z) in the series:
$$\text{CH}_2=\text{CH}_2 \xrightarrow{\text{HBr}} (\text{X}) \xrightarrow{\text{Hydrolysis}} (\text{Y}) \xrightarrow[\text{I}_2 (\text{excess})]{\text{NaOH}} (\text{Z})$$

a) $\text{C}_2\text{H}_5\text{I}$ b) $\text{C}_2\text{H}_5\text{OH}$ c) CHI_3 d) CH_3CHO
497. Phenol can be converted to *o*-hydroxybenzaldehyde by
a) Kolbe's reaction b) Reimer-Tiemann reaction
c) Wurtz reaction d) Cannizaro reaction
498. An organic compound 'X' with molecular formula, $\text{C}_7\text{H}_8\text{O}$ is insoluble in aqueous NaHCO_3 but dissolves in NaOH . When treated with bromine water 'X' rapidly gives 'Y' $\text{C}_7\text{H}_5\text{OBr}_3$.
The compounds 'X' and 'Y' respectively, are
a) Benzyl alcohol and 2, 4, 6-tribromo-3-methoxy benzene
b) Benzyl alcohol and 2, 4, 6-tribromo-3-methyl phenol
c) *o*-cresol and 3, 4, 5-tribromo-2-methyl phenol
d) Methoxybenzene and 2, 4, 6-tribromo-3-methoxy benzene
499. Which of the following compound would not evolve CO_2 when treated with NaHCO_3 solution?
a) Salicylic acid b) Phenol c) Benzoic acid d) 4-nitrobenzoic acid
500. For which pair iodoform test cannot be used as distinction test?
a) Propanol-1 and propanol-2
b) Butanol-2 and 2-methyl propan-2-ol
c) Butanol-1 and butanol-2
d) Pentanol-1 and pentanol-3
501. Tonics usually contain small amount of:
a) Formalin b) Vinegar c) Alcohol d) Ether
502. Primary, secondary and tertiary alcohols can be distinguished by performing
a) Beilstein's test b) Victor Meyer's test c) Fehling's solution test d) Hofmann's test
503. Ethanol reacts with thionyl chloride to give ethyl chloride and:
a) S, SO_2 b) SO_2 , HCl c) Cl_2 , SO_3 d) SO_3 , HCl
504. The product C in the following sequence of reaction,
$$\text{C}_2\text{H}_5\text{Br} \xrightarrow{\text{NaOH} (aq)} \text{A} \xrightarrow{\text{Na}} \text{B} \xrightarrow{\text{CH}_3\text{I}} \text{C}$$
 is:
a) Butane b) Ethane c) Methyl ethyl ether d) propane
505. Which of the following is an anaesthetic?
a) Ether b) Thiobarburates c) Trichloromethane d) All of these
506. In the reaction,




(vapour)



The molecular formula of X is



507. In which of the following bond angles on sp^3 -hybridized are not contracted due to lone pair of electron?



508. By which the following reactions can *trans*-cyclopentane-1, 2-diol be obtained?



d) None of these

509. A compound X, when boiled with Na_2CO_3 solution gives glycol as the product. What is X ?

a) Ethylene

b) Ethylene oxide

c) Ethyl bromide

d) Ethyl hydrogen sulphate

510. Glycerol is present as a triester in:


a) Petroleum

b) Kerosene oil





c) Vegetable oil and fats

d) Naphtha




511.




by RMgX which is the incorrect pair?



512. Which alcohol cannot be oxidized by MnO_2 ?



513. The reaction,






Is called



a) Laderer Mannasse reaction

b) Claisen condensation

- c) Benzoin condensation
d) Etard reaction
514. An alcohol is not oxidised in alkaline or neutral solution but in acidic solution it is turned first to acetone and then to acetic acid. It is a:
a) Primary alcohol b) Secondary alcohol c) Tertiary alcohol d) None of these
515. In the reaction $\text{Ph} - \begin{array}{c} \text{CH}_3\text{CH}_3 \\ | \quad | \\ \text{C} - \text{C} - \text{Ph} \\ | \quad | \\ \text{OH} \quad \text{OH} \end{array} \xrightarrow{\text{Conc. H}_2\text{SO}_4} \text{A}$ The product A is
- a) $\text{Ph} - \begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ | \quad | \\ \text{C} = \text{C} - \text{Ph} \end{array}$ b) $\text{Ph} - \begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ | \quad | \\ \text{C} - \text{C} - \text{Ph} \\ \diagdown \quad \diagup \\ \text{O} \end{array}$ c) $\text{Ph} - \begin{array}{c} \text{CH}_3 \quad \text{O} \\ | \quad || \\ \text{C} - \text{C} - \text{CH}_3 \\ | \\ \text{Ph} \end{array}$ d) $\text{Ph} - \begin{array}{c} \text{CH}_3\text{O} \\ | \quad || \\ \text{C} - \text{C} - \text{Ph} \\ | \\ \text{CH}_3 \end{array}$
516. Which reagent will convert propionic acid to propanol-1?
a) KMnO_4 b) LiAlH_4 c) Cr_2O_3 d) MnO_2
517. Which of the following is a gas?
a) Methane thiol b) Ethane thiol c) Isobutyl thiol d) Propyl thiol
518. Alcohols may behave as:
a) Bronsted acid b) Lewis base c) Neutral d) All of these
519. The reaction;
 $\text{RCOOH} \xrightarrow{\text{C}_2\text{H}_5\text{OH} + \text{Na}} \text{RCH}_2\text{OH}$ is called:
a) Corey House reaction
b) Bonveault-Blanc reaction
c) Clemmensen reduction
d) None of the above
520. Absolute alcohol is prepared by
a) Vacuum distillation b) Azeotropic distillation
c) Steam distillation d) None of the above
521. On heating glycerol with conc. H_2SO_4 , a compound is obtained which has bad odour. The compound is:
a) Acrolein b) Formic acid c) Allyl alcohol d) Methyl isocyanide
522. Pyroligneous acid contains:
a) CH_3COOH (10%), CH_3OH b) $\text{C}_2\text{H}_5\text{OH}$ (10%), CH_3OH c) CH_3COCH_3 (10%), $\text{C}_2\text{H}_5\text{OH}$ d) None of the above
523. Ethyl alcohol reacts with HCl but not with HCN because:
a) $\text{C}_2\text{H}_5\text{OH}$ is weak base and HCN is weak base
b) $\text{C}_2\text{H}_5\text{OH}$ is strong acid and HCN is weak acid
c) HCl is strong acid and $\text{C}_2\text{H}_5\text{OH}$ is weak base
d) None of the above
524. When wine is put in air it becomes sour due to:
a) Oxidation of $\text{C}_2\text{H}_5\text{OH}$ into CH_3COOH
b) Bacteria
c) Virus
d) Formic acid formation
525. Dunstan's test is used for identification of
a) Acetone b) Ethanol c) Glycerol d) Glycol
526. An alcohol on oxidation is found to give CH_3COOH and $\text{CH}_3\text{CH}_2\text{COOH}$. The alcohol is:
a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ b) $(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CH}_3$ c) $\text{CH}_3(\text{CH}_2)_2\text{CHOH}$ d) $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3$
527. The enzymes which are used to convert starch into ethyl alcohol are
a) Maltase, diastase b) Diastase, maltase, zymase



d) Invertase, diastase, maltase



529. Ethyl alcohol can be prepared from Grignard reagent by the reaction of

- a) HCHO b) $R_2\text{CO}$ c) RCN d) RCOCl


530. The correct order of the solubility of different alcohols in water is

- a) Ethanol > *n*-propanol > *n*-butyl alcohol
 b) *n*-propyl alcohol > ethyl alcohol > *n*-butyl alcohol
 c) ethyl alcohol > *n*-butyl alcohol > *n*-propyl alcohol
 d) *n*-butyl alcohol > *n*-propyl alcohol > ethyl alcohol





531. Germinated Barley (an enzyme) is a source of enzyme:

- a) Zymase b) Diastase c) Maltase d) Invertase

532. In the reaction,




The products are



533. Methylphenyl ether can be obtained by reacting

- a) Phenolate ions and methyl iodide b) Methoxide ions and bromobenzene
 c) Methanol and phenol d) Bromobenzene and methyl bromide



In the above sequence *X* can be

- a) H_2/Ni
 b) NaBH_4
 c) $\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+$
 d) Both (a) and (b)

535. To distinguish between salicylic acid and phenol one can use

- a) NaHCO_3 solution b) 5% NaOH solution c) Neutral FeCl_3 d) Bromine water

536. Diethyl ether finds its use in medicine as:


- a) Pain killer b) Hypnotic c) Antiseptic d) Anaesthetic

537. Ethyl chloride reacts with sodium ethoxide to form a compound *A*. Which of the following reactions also yields *A*?

- a) C_2H_5Cl , KOH (alc.), Δ
 c) C_2H_5Cl , Mg(dry ether)


- b) $2C_2H_5OH$, conc. H_2SO_4 , $140^\circ C$
 d) C_2H_2 , dil H_2SO_4 , $HgSO_4$

538. In the following sequence of reactions,



The compound 'D' is

- a) Butanal
 b) *n*-butyl alcohol
 c) *n*-propyl alcohol
 d) Propanal



This reaction is called

- a) Reimer-Tiemann reaction
 b) Liebermann's nitroso reaction
 c) Dakin reaction
 d) Lederer -Manasse reaction


540. Carbocation is not the intermediate in








- a) Hydroboration-oxidation of an alkene
 b) Oxymercuration-demercuration of an alkene
 c) Reaction of HCl with CH_3CH_2OH
 d) All of the above

541. The number of isomeric alcohols of formula $C_4H_{10}O$ is:

- a) 2
 b) 4
 c) 7
 d) 8

542. The final product of the following reaction is/are



- a) 
 b)  + 
 c)  + 
 d)  + 

543. Anisole is the product obtained from phenol by the reaction known as

- a) Coupling
 b) Etherification
 c) Oxidation
 d) Esterification

544. Propan-1-ol can be prepared from propane by

- a) H_2O/H_2SO_4
 b) $Hg(OAc)_2/H_2O$ followed by $NaBH_4$
 c) B_2H_6 followed by H_2O_2
 d) CH_3CO_2H/H_2SO_4

545. Lubricant used in watch is:

- a) Coconut oil
 b) Pine oil
 c) Animal oil
 d) Glycerol

546. Methyl alcohol on oxidation with acidified $K_2Cr_2O_7$ gives:


- a) CH_3COCH_3
 b) CH_3CHO
 c) $HCOOH$
 d) CH_3COOH

547. Lucas reagent is a mixture of:






- a) Conc. HCl + anhydrous $ZnCl_2$
 b) Conc. HCl + hydrous $ZnCl_2$
 c) Conc. HNO_3 + hydrous $ZnCl_2$
 d) Conc. HNO_3 + anhydrous $ZnCl_2$

548. If methanol vapour is passed over heated copper at $300^\circ C$, it forms formaldehyde by:

- a) Hydrogenation b) Dehydrogenation c) Dehydration d) Oxidation
549. Terylene is formed by the reaction of one of the following alcohols:
 a) 2-chloroethanol b) 1,2,3-propanetriol c) Ethanediol d) Phenol
550. Alcoholic fermentation by starch or sugar is brought about by:
 a) CO_2 b) Sodium bicarbonate c) Yeast d) phosphates
551. General formula for alcohols is:
 a) >COH b) >CHOH c) $-\text{CH}_2\text{OH}$ d) All of these




A and B respectively are

- a) Both 
- b) Both 
- c)  , 
- d)  , 


553. When phenol reacts with phthalic anhydride in presence of H_2SO_4 and heated and hot reaction mixture is poured in NaOH solution, then product formed is
 a) Alizarin b) Methyl orange c) Fluorescein d) Phenolphthalein

554. Correct order of dehydration of IX.



- a) $A > B > C > D$ b) $B > C > A > D$ c) $D > A > C > A$ d) $D > A > B > C$



555. The following reaction is known as




- a) Perkin reaction b) Gattermann reaction
 c) Kolbe reaction d) Gattermann-aldehyde reaction

556. In the Liebermann test for phenols, the blue or green colour produced is due to the formation of





557. Four hydroxy compounds have functional groups as shown



The purple colour with FeCl_3 will be given by

- a) A only b) A and B c) C only d) A, B, C and D

558. Ether in contact with air for a long time form peroxides. The presence of peroxide in ether can be tested by adding Fe^{2+} ion and then adding

- a) KCN b) SnCl_2 c) HgCl_2 d) KCNS

559. Fermentation is:

- a) Exothermic b) Endothermic c) Reversible d) None of these

560. Which could not be obtained from wood?

- a) CH_3OH b) $\text{C}_2\text{H}_5\text{OH}$ c) Wood tar d) Wood charcoal

561. Methanol and ethanol can be distinguished by the following:

- a) By reaction with metallic sodium
b) By reaction with caustic soda
c) By heating with iodine and washing soda
d) By heating with zinc and inorganic mineral acid

562. Acetic anhydride reacts with diethyl ether in the presence of anhydrous AlCl_3 to give

- a) $\text{CH}_3\text{CH}_2\text{COOH}$ b) $\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_3$ c) $\text{CH}_3\text{COOCH}_3$ d) $\text{CH}_3\text{COOC}_2\text{H}_5$

563. Which of the following is insoluble in alcohol?

- a) Resins and varnishes b) Soaps and varnishes c) Rubbers and plastics d) Dyes and drugs

564. 1-propanol and 2-propanol can be distinguished by

- a) Oxidation with alkaline KMnO_4 followed by reaction with Fehling solution
b) Oxidation with acidic dichromate followed by reaction with Fehling solution
c) Oxidation by heating with copper followed by reaction with Fehling solution
d) Oxidation with concentrated H_2SO_4 followed by reaction with Fehling solution

565. Which of the following does not react with sodium metal?

- a) $(\text{CH}_3)_2\text{O}$ b) $\text{CH}_3\text{CH}_2\text{OH}$ c) CH_3COOH d) $\text{C}_6\text{H}_5\text{OH}$





566. Purity of ether before using it as anaesthetic agent is tested by:

- a) $\text{KI} + \text{starch}$ b) CuSO_4 c) H_2SO_4 d) None of these


567. Alcoholic beverages contain

- a) Isopropyl alcohol b) *n*-propyl alcohol c) Ethyl alcohol d) Methyl alcohol



568. Picric acid is





569. The final product (IV) in the sequence of reactions



is





570. The products of combustion of an aliphatic thiol (RSH) at 298 K are

- a) $\text{CO}_2(\text{g}), \text{H}_2\text{O}(\text{g})$ and $\text{SO}_2(\text{g})$ b) $\text{CO}_2(\text{g}), \text{H}_2\text{O}(\text{l})$ and $\text{SO}_2(\text{g})$
 c) $\text{CO}_2(\text{l}), \text{H}_2\text{O}(\text{l})$ and $\text{SO}_2(\text{g})$ d) $\text{CO}_2(\text{g}), \text{H}_2\text{O}(\text{l})$ and $\text{SO}_2(\text{l})$

571. During alcoholic fermentation inorganic salts like ammonium sulphate or ammonium phosphate are added:

- a) To decrease the freezing point of solution
 b) Which act as food for ferment cells
 c) Which prevent the growth of undesirable bacteria
 d) Which produce desirable enzymes

572. To obtain unsaturated alcohols from unsaturated aldehydes the following reagent is used for reduction:

- a) Na amalgam/ H_2O b) Dil. H_2SO_4 c) Zn/HCl d) LiAlH_4

573. Hydroboration oxidation of 4-methyl octene would give

- a) 4-methyl octanol b) 2-methyl decane
 c) 4-methyl heptanol d) 4-methyl-2-octanone

574. $Z \xrightarrow{\text{PCl}_5} X \xrightarrow{\text{Alc. KOH}} Y \xrightarrow[2. \text{H}_2\text{O}; \text{boil}]{1. \text{Conc. H}_2\text{SO}_4} Z$ is :

- a) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{OH}$ b) $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\ | \\ \text{OH} \end{array}$ c) $(\text{C}_2\text{H}_5)_3 \text{C} - \text{OH}$ d) $\text{CH}_3 - \text{CH} = \text{CH}_2$

575. The general molecular formula, which represents the homologous series of alkanols is:

- a) $\text{C}_n\text{H}_{2n+1}\text{O}$ b) $\text{C}_n\text{H}_{2n+2}\text{O}$ c) $\text{C}_n\text{H}_{2n}\text{O}_2$ d) $\text{C}_n\text{H}_{2n}\text{O}$


576. On reacting with neutral ferric chloride, phenol gives

- a) Red colour b) Blue colour c) Violet colour d) Green colour

577. There are four alcohols P, Q, R and S which have 3, 2, 1 and zero alpha hydrogen atom(s). Which one of the following will not respond to Victor-Meyer's test?


- a) P b) Q c) R d) S





578. Which doesn't form in the acid catalysed rearrangement of cumene hydroperoxide?



579. Ethanol is more soluble in water but ether is less soluble because:

- a) Ethanol forms strong hydrogen bonds in water whereas ether forms weaker hydrogen bonding
 b) Ether is more volatile than ethanol
 c) The molecular weight of ether is more than that of ethanol
 d) None of the above



- a)  b)  c)  d) 

CHEMISTRY (QUESTION BANK)**11.ALCOHOLS, PHENOLS AND ETHERS****: ANSWER KEY :**


1)	a	2)	c	3)	b	4)	d	149)	b	150)	b	151)	c	152)	d
5)	b	6)	c	7)	a	8)	b	153)	a	154)	d	155)	a	156)	c
9)	a	10)	b	11)	d	12)	d	157)	c	158)	b	159)	a	160)	d
13)	c	14)	d	15)	a	16)	c	161)	c	162)	b	163)	c	164)	a
17)	c	18)	d	19)	b	20)	c	165)	c	166)	a	167)	d	168)	c
21)	b	22)	a	23)	b	24)	a	169)	c	170)	a	171)	d	172)	d
25)	a	26)	b	27)	c	28)	c	173)	d	174)	a	175)	d	176)	d
29)	b	30)	b	31)	a	32)	d	177)	c	178)	a	179)	c	180)	d
33)	a	34)	b	35)	b	36)	c	181)	b	182)	d	183)	d	184)	b
37)	b	38)	b	39)	b	40)	d	185)	a	186)	c	187)	d	188)	b
41)	d	42)	a	43)	c	44)	b	189)	c	190)	b	191)	d	192)	c
45)	a	46)	b	47)	d	48)	b	193)	b	194)	d	195)	d	196)	c
49)	b	50)	c	51)	b	52)	b	197)	d	198)	d	199)	d	200)	a
53)	d	54)	c	55)	b	56)	a	201)	d	202)	c	203)	c	204)	c
57)	d	58)	b	59)	d	60)	a	205)	b	206)	c	207)	b	208)	a
61)	c	62)	d	63)	d	64)	b	209)	a	210)	d	211)	c	212)	d
65)	a	66)	a	67)	d	68)	d	213)	c	214)	a	215)	c	216)	c
69)	a	70)	a	71)	c	72)	a	217)	b	218)	a	219)	b	220)	a
73)	d	74)	d	75)	b	76)	c	221)	a	222)	a	223)	c	224)	a
77)	c	78)	d	79)	a	80)	c	225)	d	226)	a	227)	c	228)	b
81)	c	82)	b	83)	d	84)	a	229)	d	230)	c	231)	b	232)	b
85)	a	86)	c	87)	d	88)	d	233)	a	234)	c	235)	c	236)	a
89)	c	90)	c	91)	b	92)	b	237)	d	238)	b	239)	c	240)	d
93)	b	94)	c	95)	d	96)	c	241)	d	242)	d	243)	a	244)	a
97)	d	98)	a	99)	d	100)	c	245)	b	246)	a	247)	a	248)	d
101)	a	102)	b	103)	d	104)	c	249)	d	250)	c	251)	c	252)	b
105)	a	106)	c	107)	d	108)	c	253)	b	254)	c	255)	b	256)	b
109)	c	110)	b	111)	a	112)	a	257)	d	258)	a	259)	a	260)	b
113)	c	114)	d	115)	a	116)	d	261)	b	262)	c	263)	b	264)	a
117)	d	118)	c	119)	c	120)	c	265)	a	266)	b	267)	a	268)	c
121)	a	122)	a	123)	c	124)	b	269)	b	270)	c	271)	c	272)	d
125)	a	126)	c	127)	b	128)	d	273)	c	274)	b	275)	d	276)	c
129)	c	130)	a	131)	c	132)	a	277)	c	278)	c	279)	c	280)	a
133)	a	134)	a	135)	d	136)	d	281)	a	282)	d	283)	c	284)	a
137)	b	138)	c	139)	a	140)	d	285)	a	286)	c	287)	c	288)	d
141)	b	142)	d	143)	d	144)	b	289)	b	290)	c	291)	d	292)	c
145)	b	146)	a	147)	c	148)	b	293)	d	294)	d	295)	d	296)	b

297) b	298) b	299) a	300) c	445) d	446) c	447) b	448) a
301) c	302) d	303) a	304) d	449) c	450) d	451) c	452) b
305) d	306) c	307) c	308) d	453) b	454) b	455) c	456) a
309) b	310) d	311) a	312) c	457) b	458) a	459) d	460) a
313) c	314) c	315) a	316) b	461) c	462) a	463) a	464) c
317) a	318) c	319) a	320) b	465) a	466) c	467) d	468) a
321) b	322) c	323) b	324) b	469) c	470) c	471) a	472) a
325) d	326) b	327) c	328) b	473) d	474) b	475) c	476) c
329) c	330) a	331) d	332) d	477) a	478) c	479) c	480) c
333) d	334) c	335) c	336) d	481) d	482) b	483) c	484) b
337) c	338) b	339) a	340) a	485) a	486) c	487) a	488) a
341) b	342) c	343) c	344) d	489) a	490) d	491) a	492) b
345) a	346) a	347) b	348) a	493) a	494) b	495) d	496) c
349) d	350) a	351) b	352) b	497) b	498) c	499) b	500) d
353) a	354) d	355) b	356) c	501) c	502) b	503) b	504) c
357) d	358) c	359) c	360) d	505) d	506) c	507) c	508) c
361) c	362) c	363) c	364) a	509) b	510) c	511) d	512) a
365) b	366) d	367) c	368) d	513) a	514) c	515) c	516) b
369) a	370) b	371) d	372) d	517) a	518) d	519) b	520) b
373) c	374) c	375) a	376) a	521) a	522) a	523) c	524) a
377) b	378) b	379) b	380) b	525) c	526) d	527) b	528) c
381) b	382) a	383) a	384) a	529) a	530) a	531) b	532) d
385) d	386) b	387) b	388) b	533) a	534) b	535) a	536) d
389) c	390) b	391) c	392) b	537) b	538) c	539) c	540) d
393) d	394) b	395) a	396) b	541) b	542) b	543) b	544) c
397) c	398) c	399) c	400) b	545) d	546) c	547) a	548) b
401) a	402) a	403) a	404) b	549) c	550) c	551) d	552) d
405) c	406) b	407) d	408) a	553) d	554) b	555) d	556) d
409) c	410) d	411) a	412) a	557) c	558) d	559) a	560) b
413) d	414) a	415) b	416) b	561) c	562) d	563) c	564) c
417) d	418) c	419) c	420) a	565) a	566) a	567) c	568) c
421) d	422) d	423) d	424) a	569) c	570) b	571) b	572) d
425) d	426) c	427) b	428) c	573) a	574) b	575) b	576) c
429) a	430) b	431) a	432) a	577) d	578) d	579) a	580) a
433) c	434) a	435) b	436) b				
437) a	438) d	439) c	440) b				
441) c	442) a	443) d	444) b				






CHEMISTRY (QUESTION BANK)

11.ALCOHOLS, PHENOLS AND ETHERS












: HINTS AND SOLUTIONS :




(Y)
(yellow ppt
tri-iodomethane)




Formaldehyde gives primary alcohol with Grignard reagent while any other aldehyde except formaldehyde give secondary alcohol.



Sodiummethoxide bromo ethoxyethane
ethane





It is tertiary alcohol.

23 (b)


Peroxides are decomposed on heating with H_2SO_4

24 (a)

Glycerol is not reduced because of extensive H-bonding.


25 (a)

When an alkyl aryl ether is heated with HI, halogen goes with alkyl group. Therefore, heating anisole (methyl phenyl ether) with HI phenol and methyl iodide are obtained.



26 (b)

C – Br bond is weaker as compared to C – Cl bond




29 (b)

Higher concentration of substrate less easily undergoes fermentation; (a), (c), (d) favours fermentation.

30 (b)

The correct order of stability of carbocation is as follows




31 (a)

Alcohols with same molecular weight are expected to have almost same boiling point however two more factors other than molecular weight are important, they are namely H-boiling and surface area of molecule. Both these factors are least in 3° alcohols and maximum in 1° alcohols. Hence, 3° alcohols have least boiling

point while 1° alcohols have maximum boiling point.



32 (d)



will not with ϕNNCl to give dye

33 (a)

Phenol cannot be distinguished from ethanol by sodium because both evolve hydrogen with sodium.




34 (b)

Ethers do not contain acidic H-atom.

35 (b)

Ethylene glycol reacts with excess of PCl_5 to give ethylene chloride.




36 (c)

Due to low f.p. and mobile nature.

37 (b)


Structure of phenolphthalein in basic medium is as follows.



38 (b)


$\text{C}_2\text{H}_5\text{O}^-$ will attract the proton from phenol converting the later into phenoxide ion. This would then make nucleophilic attack on the methylene carbon of alkyl iodide, but $\text{C}_2\text{H}_5\text{O}^-$ is in excess $\text{C}_2\text{H}_5\text{O}^-$ is better nucleophile than $\text{C}_6\text{H}_5\text{O}^-$ (phenoxide) ion since while in the former the negative charge is localised over oxygen and in the later it is delocalised over the whole molecular frame work. So, it is $\text{C}_2\text{H}_5\text{O}^-$ ion that

would make nucleophilic attack at ethyl iodide to give diethyl ether (Williamson's synthesis).




40 (d) -OH gp. directly attached to benzene nucleus represents for phenolic gp.

41 (d) The pinacol-pinacolone rearrangement involves dehydration of diols through the formation of carbocation intermediate which rearranges to more stable compound.




42 (a) Oxidation of glycerol by KMnO_4 is violent.



sp^3 and sp^3 -hybridisations of carbon and oxygen in electronic structure of ether.

45 (a) Protonation of -OH is first step. Conversion of poor leaving group (-OH) into good leaving group (-OH₂⁺).




46 (b) It contains (R)₃COH.

48 (b) When an electron attracting group (like -NO₂, -Cl) is attached to the phenol ring, it stabilises the negative charge on the oxygen of phenoxide ion. Due to this reason acidic character of phenol increases. But when an electron donating group (like -CH₃) is attached to the

phenol ring, it destabilises the ring and hence, acidic character of phenol decreases. Thus, the correct order of acidic character is *p*-nitrophenol > *p*-chlorophenol > phenol > *o*-cresol.

49 (b) Glycerol react with oxalic acid at 110°C temperature, it gives methanoic acid (formic acid).



50 (c)


$$\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3\text{---C---CH}_2\text{OH} \\ | \\ \text{CH}_3 \end{array} \longrightarrow \begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3\text{---C}=\text{CH---CH}_3 \\ | \\ \text{CH}_3 \end{array}$$

due to rearrangement of carbocation following alkyl shift.

51 (b)

First sulfonation is the means to block *para* position and to reduce the reactivity of phenolic ring against strong oxidising agent HNO₃. (The use of conc. HNO₃ over phenol cause the oxidation of ring mainly). The strong acidic medium in second step cause desulfonation (ipso mechanism) also.

52 (b) Glycols are dihydric alcohols (having two hydroxyl groups). Ethylene glycol is the first member of this series.



CH₂OH
ethylene glycol

53 (d)

Absolute alcohol is 100% alcohol.

54 (c)

The order of reactivity depends upon the stability of the carbocation formed

ie, FCH₂⁺C⁺HCH₃, FCH₂CH₂⁺C⁺HCH₃, CH₃⁺C⁺HCH₃

and Ph⁺C⁺H₂. The stability order of carbocations is

Ph⁺C⁺H₂ > CH₃⁺C⁺HCH₃ > FCH₂CH₂⁺C⁺HCH₃ >

FCH₂⁺C⁺HCH₃. Thus, the order of reactivity follows

the order IV > III > II > I

55 (b)

Glycerol trinitrate adsorbed on Kieselguhr is called dynamite; an explosive.

57 (d)

4 alcohols (butan-1-ol; butan-2-ol; 2-methyl butan-1-ol; 2-methyl butan-2-ol) and 3 ethers (diethyl ether, methyl-propyl ether and methyl isopropyl ether).





59 (d)

ROH + HBr → R - Br + H₂O

The rate of reaction is fastest for 3° alcohol.

The rate of reaction decreases as follows

3° > 2° > 1°



60 (a)

Alcohols (ROH) are hydroxy derivatives of alkane or alkyl derivative of water.



61 (c)

Presence of two isopropyl groups on oxygen atom of ether shows more powerful inductive effect.


63 (d)

Alcohol is initially protonated by the acid to form protonated alcohol or oxonium ion. It is then

attacked by a second molecule of alcohol which acts as nucleophile



64 (b)



65 (a)


Electron withdrawing groups (like -NO₂) increase the acidity of phenols by stabilising corresponding phenoxide ion. The effect of -NO₂ group will be minimum at *m*-position due to lack of increased delocalisation of electrons in it. Hence, *m*-nitrophenol is the weakest acid among these.

66 (a)


Fusel oil is a mixture of pentanol and butanol with other organic substances.

67 (d)

Benzene sulphonic acid and *p*-nitro phenol react with NaHCO₃ and evolve CO₂ gas.



Because benzene sulphonic acid *p*-nitrophenol are stronger acids, so they are capable to evolve CO₂ with NaHCO₃.




68 (d)

Secondary alcohols give blue colour in Victor Meyer test


69 (a)

Conc. HCl + anhydrous $ZnCl_2$ is called as Lucas reagent. It is used to distinguish primary, secondary and tertiary alcohol.

70 (a)



71 (c)



2-propanol acetone acetic acid

73 (d)



Phenol reacts with PCl_5 to form chlorobenzene. Halogenation of phenol does not take place with HX

74 (d)

Alcohol has polar H which makes intermolecular H-bonding possible. Ether is non-polar hence no H-bonding. Lack of H-bonding in ether makes it more volatile than alcohol.

76 (c)

In the given sequence of reaction, the alcohol is tertiary.



77 (c)


It is better to call nitroglycerine as glycerol trinitrate an inorganic ester of HNO_3 and glycerol.

78 (d)

Br is replaced by OH gp.

80 (c)

Glycerol is dehydrated on heating with $KHSO_4$.



81 (c)

1° alkyl halides on treatment with an alkoxide ion tend to undergo substitution to form ethers. So sodium tert butoxide and ethyl bromide reagent is used


82 (b)

A mixture of $H_2O_2 + FeSO_4$ is called Fenton's reagent used as oxidant.


83 (d)

A simple ether is one which possesses same alkyl groups on O atom, . e., ROR.

84 (a)



85 (a)




86 (c)


Presence of two or more OHgp. on a carbon atom makes it unstable and compound loses H_2O molecule.

87 (d)

A mixture of conc. HCl + anhy $ZnCl_2$ is called Lucas reagent. In Lucas test tertiary alcohols immediately give turbidity while secondary alcohols give turbidity after 5 min. Primary alcohols give no reaction with Lucas reagent at room temperature.




Primary alcohol and hence, no white cloudiness or turbidity at room temperature.



Secondary alcohol

White cloudiness or turbidity appears within about 5 min.



Tertiary alcohol

Or turbidity appears immediately.

88 (d)

To have tertiary alkyl-alkyl ether one needs sod. Tertiary alkoxide and alkyl halide.

89 (c)

Due to H-bonding.

90 (c)

Glyoxal is a trivial name for ethane-1-2-dial.

91 (b)


Ethyl alcohol is mixed with methyl alcohol to denaturate it in order to prevent its use for drinking purposes.

93 (b)


The density of glycerol is higher than propanol due to extensive intermolecular hydrogen

bonding. Glycerol contains three -OH groups while propanol contains only one -OH group.

95 (d)



Due to *-I* and *-R* influence, NO_2 in *ortho*-position should have raised the acidity to the maximum extent. But it is due to intramolecular H-bonding, *ortho*-nitrophenol is less acidic than *para*-nitrophenol.



96 (c)

$\text{CH}_3\text{OC}_2\text{H}_5 + \text{HI} \rightarrow \text{CH}_3\text{I} + \text{C}_2\text{H}_5\text{OH}$;
O-atom goes with higher alkyl gp.

97 (d)


Glycerol is $\text{CH}_2\text{OHCHOHCH}_2\text{OH}$

98 (a)

Due to intermolecular hydrogen bonding, alcohols are less volatile than ether

99 (d)

In the presence of conc. H_2SO_4 , two molecules of phenol condense with phthalic anhydride to form phenolphthalein




100 (c)

The mixture shows positive deviations from Raoult's law; *i. e.*, $\Delta H_{\text{mix}} > 0$, $\Delta V_{\text{mix}} > 0$.

101 (a)


Benzyl phenyl ether is an unsymmetrical ether so halide ion of HI attached to the simple alkyl group and reaction takes place by following mechanism.

1. Protonation of ether



Benzyl phenyl ether

2. Nucleophilic attack



102 (b)


$\text{CH}_2\text{OH} \xrightarrow[170^\circ\text{C}]{\text{H}_2\text{SO}_4} \text{C}_2\text{H}_4 + \text{H}_2\text{O}$; Removal of H_2O from substrate molecule is called dehydration.

It is classified as elimination reaction.

104 (c)

The acid H_2SO_4 is added to adjust pH in between 4 to 4.5 which is favourable for the growth of yeast and unfavourable for the growth of undesired bacteria.

105 (a)



This is acid catalysed cleavage of cyclic ether where nucleophile attacks the more substituted carbon.

106 (c)

$\text{R}-\text{S}-\text{R}$ or $\text{R}-\text{SR}'$ are thioethers.

107 (d)

All are dehydrating agents.

108 (c)

$\text{C}_2\text{H}_5\text{OH} + [\text{O}] \xrightarrow[\text{H}_2\text{SO}_4/\Delta]{\text{KMnO}_4/\text{H}_4} \text{CH}_3\text{COOH} \xrightarrow[\text{H}_2\text{SO}_4/\Delta]{\text{C}_2\text{H}_5\text{OH}(Y)} \text{CH}_3\text{COOC}_2\text{H}_5$
ethanol (X) ethyl ethanoate (esterification)

ethanoic acid

Hence, $X = \text{CH}_3\text{COOH}$



$Y = \text{C}_2\text{H}_5\text{OH}$

109 (c)

Lower members are soluble in water due to H-bonding and solubility decreases with increasing hydrophobic character.

138 (c)

Ether on reaction with excess of HI produce two molecules of alkyl halide.



Ethyl iodide


When equimolar quantities of ether and HI are present, then one molecule of alkyl halide and one molecule of alcohol are formed.

139 (a)

It is a substitute of petrol.

141 (b)

Zymase enzyme act on glucose and give ethyl alcohol and carbon dioxide.



142 (d)

Only CH_2OH group is oxidized to $-\text{COOH}$; Double bond is not affected.

143 (d)

Both Zn-Hg/HCl and $\text{NH}_2\text{NH}_2, \text{OH}^-$ reduce CO to CH_2 , but acid sensitive reagents are not reduced by Zn-Hg/HCl .

144 (b)


Glycerol has 3 $-\text{OH}$ groups and thus shows extensive H-bonding.

145 (b)

The best method to prepare cyclohexene from cyclohexanol is by conc. H_3PO_4 because in given options dehydrating agent is conc. H_3PO_4 .

146 (a)

Diethyl sulphate in the presence of NaOH acts as alkylating agent, it causes alkylation of phenol to give ethyl phenyl ether which is also called phenetole.



147 (c)

Tertiary alcohols are dehydrated on passing over heated Cu ; Primary and secondary are dehydrogenated.

148 (b)

The process is called hydroboration.

149 (b)

Secondary alcohols give turbidity within 5 min with Lucas reagent

151 (c)



Diethyl ether itself being a Lewis base is not attacked by nucleophiles, *ie*, OH^- ion. All others contain an electrophilic carbon and are readily attacked by nucleophile

152 (d)

Ethers acts as Lewis base only towards strong acids.

154 (d)

When ethyl alcohol is heated with conc. H_2SO_4 at $160^\circ - 170^\circ\text{C}$, the product obtained is ethylene (C_2H_4).




ethylene

But at lower temperature ether is formed.

155 (a)

Phenol is heated with phthalic anhydride in presence of conc H_2SO_4 to give phenolphthalein which gives pink colour with alkali




156 (c)

Large is $\text{H}-\text{X}$ bond length, more is acidic nature of halogen acid.


157 (c)

Rectified spirit is $\text{C}_2\text{H}_5\text{OH} +$ water mixture obtained after distillation of fermented liquid. On further careful fractional distillation (rectification) gives II fraction as 93 to 95% ethyl alcohol (rectified spirit).


158 (b)



159 (a)




This is Liebermann's nitroso reaction of phenol. When phenol is warmed with sodium nitrite and 1 cc. conc. H_2SO_4 , blue colour is obtained which on adding water, becomes red. This again turns to blue on adding NaOH. Deep blue colour is due to the formation of sodium salt of indophenol.



160 (d)

H_2SO_4 acts as catalyst as well as dehydrating agent for the reaction,



161 (c)


Dynamite is known as nobel's oil.

162 (b)

The order of increasing acidic strength is *p*-methyl phenol < phenol < *m*-nitrophenol < *p*-nitrophenol


164 (a)

$\text{C}_6\text{H}_5\text{O}^-$ is a weaker nucleophile than $\text{C}_2\text{H}_5\text{O}^-$. Therefore, the better nucleophile, *ie*, $\text{C}_2\text{H}_5\text{O}^-$ will attack $\text{C}_6\text{H}_5\text{I}$ to form diethyl ether.





165 (c)

Oxymercuration-demercuration occurs by a more stable carbocation.



166 (a)

m-cresol due to phenoxide ion in H_2O solvent, gives tribromoderivative at all *ortho* and *para* positions.




167 (d)


	Reagent	Phenol	Benzoic acid	Conclusion
A	Aqueous NaOH	Salt formation	Salt formation	No specific colour change
B	Neutral FeCl_3	Violet colour	Buff-coloured precipitate	Thus, FeCl_3 can be used to make distinction

168 (c)

$-\text{OH}$ group is an activating group, hence increase electron density on *o*- and *p*-position in benzene ring. Thus, phenol very easily undergoes nitration (electrophilic substitution and give trinitrophenol, *ie*, picric acid).




169 (c)



(a), (b), (d) may be formed but (c) is never formed Hence, correct choice → (c).

170 (a)


Greater the conjugation, greater the stability due to resonance and easier the dehydration. Thus, the correct order of dehydration is



172 (d)

Reduction by H_2 is favoured by catalyst.

173 (d)



(R is not CH_3)

174 (a)

Tertiary alcohols give alkene.

177 (c)

Tertiary alcohols are easily dehydrated.


178 (a)

Peroxide oxidizes Fe^{2+} to Fe^{3+} which gives red colour with KCNS.

182 (d)

Peppermint is soluble in alcohol.

183 (d)



184 (b)

Molasses, the brown syrupy liquid left after crystallization contains about 50% sugar.


185 (a)

$R-SH$ are thiols or mercaptans.

186 (c)

Chloral hydrate [$CCl_3CH(OH)_2$] is stable due to H-bonding


188 (b)



189 (c)


Grignard reagent ($RMgX$) reacts with only those compounds which contains acidic hydrogen or which contains carbonyl group. Dimethyl ether (CH_3OCH_3) due to absence of both acidic hydrogen and carbonyl group does not react with Grignard reagent.

190 (b)



Halogen goes with simpler alkyl gp.

191 (d)



193 (b)

(A) CH_3CN ; (B) CH_3COOH ; (C) C_2H_5OH .

194 (d)

Ester + $NaOH \rightarrow$ Sodium salt of acid + Alcohol.

195 (d)


All are dehydrating agents.

197 (d)

Alcohol $\xrightarrow{[O]}$ Aldehyde or ketones with same carbon atoms.

198 (d)


Phenol, on refluxing with chloroform and sodium hydroxide followed by hydrolysis yields *o*-hydroxy benzaldehyde




199 (d)

Molecular formula C_3H_8O ($C_nH_{2n+2}O$) suggests that the organic compound is either alcohol or ether.

Since, the compound on reaction with HI gives two different compounds, It must be an unsymmetrical ether, and its formula must be $CH_3OC_2H_5$ (methoxyethane).



Methoxyethane X Y



Aqueous iodoform

200 (a)

Glycerol is generally used as an antifreeze reagent for making explosives.

201 (d)


Follow IUPAC rules.

203 (c)

Alcohol forms a azeotropic mixture with water and absolute alcohol is obtained by this mixture (rectified spirit) by adding benzene and then carrying out fractional distillation.

204 (c)

Etherates are complexes of ethers with Lewis acid



205 (b)


Glycerol has no use in match boxes.

207 (b)


Alkoxide has metal-oxygen bond.

208 (a)

Alcohols are more acidic than alkynes but less acidic than water thus, the correct order of acidity is



209 (a)



210 (d)


Ethers are relatively less active due to the absence of functional group.

211 (c)

During germination of grape juice, a brown crust is formed at the top which is called argol. This contains potassium hydrogen tartrate and is used for preparation of tartaric acid and Rochelle salt.

212 (d)

The process of benzylation of compounds containing active hydrogen such as phenol, aniline, alcohol etc, with benzoyl chloride in the presence of aqueous NaOH is called Schotten-Baumann reaction.



213 (c)


Gashol or power alcohol (ethanol + petrol) a fuel for generating power.

215 (c)

Alcohol + Acid \rightleftharpoons Ester (fruity smelling)




Alcohol (C_2H_5OH) when react with acid they produce ester and esters have fruity smell.




ethyl alcohol acetic acid fruity smell of ester

216 (c)


2° alkyl halides tend to undergo elimination. Thus bromocyclopentane on treatment with sodium ethoxide gives cyclopentene rather than cyclophenyl ethyl ether




217 (b)



(Aldehyde and acid of same carbon atoms)




(Acid of less carbon atom)



(Both of less carbon atoms)


218 (a)

Pinacol $(CH_3)_2C(OH)C(OH)(CH_3)_2$ with dil H_2SO_4 or HCl undergoes dehydration and rearranges to form ketones (pinacolone)




219 (b)

This is base catalysed cleavage of cyclic ethers where nucleophile attacks least substituted carbon.



221 (a)

By Dow process large quantities of phenol are formed by heating chlorobenzene with a 10% solution of caustic soda or sodium carbonate at $300^\circ C$ under very high pressure (200 atm)




222 (a)

Tincture of iodine is a solution of I_2 in alcohol.

223 (c)

Phenol on heating with CCl_4 and aqueous KOH gives salicylic acid. This reaction is Reimer-Tiemann reaction.




224 (a)

Cl_2 in absence of moisture has no action over CH_3OH . In presence of moisture it oxidizes CH_3OH to HCHO .


226 (a)



(Aldehyde and acid of same carbon atoms)




(Acid of less carbon atom)



(Both of less carbon atoms)

227 (c)



Note: Ethers are functional isomers of alcohols

228 (b)

The reaction of alcohol with conc. HCl and anhydrous ZnCl_2 following $\text{S}_{\text{N}}1$ pathway, so greater the stability of carbocation formed, faster is the reaction.

2-methyl propan-2-ol gives 3° carbocation. Hence, it reacts rapidly with conc. HCl and anhydrous ZnCl_2 (Lucas reagent).

229 (d)

Solubility of alcohols decreases with increasing mol. wt. because of increasing hydrophobic nature of alkyl gp.

230 (c)


It is name reaction.

231 (b)

Alcohol $\xrightarrow{[O]}$ aldehyde or ketones with same carbon atoms.


233 (a)

Catechol is most acidic out of all dihydric phenols.



The reaction is Williamson's synthesis type reaction.

234 (c)




235 (c)

Alcohol is very good solvent for many species.

236 (a)

The reaction follows Saytzeff rule which says that during dehydration reaction hydrogen is taken preferably from carbon atom having lesser hydrogen atoms.




237 (d)

Peroxides decompose violently on heating.


238 (b)

3° alcohols are resistant to oxidation under drastic condition. They first form ketone and then acid by losing one carbon at each step.




\therefore Acid having 2C is formed when 3° alcohol is oxidised under drastic conditions.

240 (d)




241 (d)

Br_2 is formed by a redox reaction :





$-\text{OH}$ group is the activating group and there is S_{E} at o - and p -positions giving yellowish white precipitate of 2, 4, 6-tribromophenol :



242 (d)

Tertiary halides do not undergo Williamson's synthesis. To get *t*-alkyl-alkyl ether, *t*-alkoxide and alkyl halide should be used.


244 (a)



The esterification by $RCOOH$ and $R'OH$ does not take place in alkaline medium.

245 (b)

Because the difference in mass between $-CH_2OH$ group and $-COOH$ group is 14, thus the compound which undergoes oxidation is a primary alcohol. ($-CH_2OH$ is the functional group of primary alcohols).




Primary alcohol acid

246 (a)

The formula represents for alcohol. Also secondary alcohol gives acid with less no. of carbon atoms.


247 (a)




HIO_4 oxidises $-CH_2OH$ to $HCHO$ and breaks the $C-C$ bond of terminal CH_2OH gps.

249 (d)

By Williamson's synthesis, alkyl halide on reaction with sodium alkoxide gives ether.




250 (c)




251 (c)

Williamson's synthesis It involves the heating of alkyl halide with sodium or potassium alkoxides. This reaction is used for the preparation of ethers e.g.,



252 (b)


CH_3OH and C_2H_5OH can be differentiated by using Na_2CO_3 and I_2 . C_2H_5OH gives yellow precipitate of CHI_3 whereas CH_3OH does not react with it.



(iodoform
Yellow ppt)

254 (c)


Dehydration of alcohol is in order




Thus, (C), a 3° alcohol is dehydrated very easily.

255 (b)

Primary alcohols get dehydrogenated with reduced copper at 573 K, to give corresponding aldehydes.






256 (b)



257 (d)

Proton donors are acids. Among given choices C_2H_5OH can give proton (H^+) most easily. $\therefore C_2H_5OH$ is most acidic among C_2H_6 , CH_4 , $CH \equiv CH$ and C_2H_5OH


258 (a)



2-bromopropane

259 (a)

The reactivity of primary halides is in the order, $CH_3 > CH_3CH_2 > CH_3CH_2CH_2$ and the tendency of alkyl halides to undergo elimination is $3^\circ > 2^\circ > 1^\circ$. Hence, for better yield, the alkyl halide should be primary and alkoxide should be secondary or tertiary.




260 (b)

283 (c)


Ethers are Lewis base and forms complex compounds with Lewis acids.

284 (a)

All those compounds which have $\left(\begin{array}{c} \text{OH} \quad \text{OH} \\ | \quad | \\ -\text{C} - \text{C}- \\ | \quad | \end{array} \right)$ groups are oxidised by periodic acid (HIO_4). Thus is not oxidised.





285 (a)




Phenol shows stabilization due to resonance.


286 (c)



2° alcohol through 2° carbocation ($\text{CH}_3\text{CH}^+\text{CH}_3$)



3° alcohol through 3° carbocation [$(\text{CH}_3)_3\text{C}^+$]




Thus, best alternate is (c)

287 (c)

Diethyl ether when heated with CO at 150°C and 500 atm pressure in presence of BF_3 forms ethyl propionate.

289 (b)

When phenolic ether is heated with HI, it gives alkyl halide and phenol




290 (c)

The red colour is due to anion of nitrolic acid.

291 (d)

Picric acid is 2, 4, 6-trinitrophenol. It is due to presence of three $-\text{I}$ showing $-\text{NO}_2$ groups, is more acidic than acetic acid and benzoic acid.




2,4,6-trinitrophenol
(picric acid)

292 (c)

CH_3OH does not contain $\text{CH}_3-\text{CHOH}-$ unit.

293 (d)



294 (d)



295 (d)


Ethers are supposed to have no functional group.

296 (b)


More is the branching in molecule, lesser is surface area and weaker are intermolecular forces.

298 (b)

The dehydration of 1-butanol gives 2-butene as the main product because 2-carbocation is stabler than 1°.



300 (c)



301 (c)

Longer the bond length, lesser will be dissociation energy and hence, more reactivity.


Among halogen acids bond length increases from HCl to HI.

\therefore Order of reactivity of halogen acids towards alcohol is



302 (d)

Alkyl halides react with sodium alkoxide to give ether. This is called Williamson's synthesis of ether.



303 (a)

Alcohols although possess low mol. wt. than thiol but they show H-bonding.

306 (c)

Alcohols are neutral and do not influence pH.

307 (c)


Enzymes are protinuous molecules derived from living organisms.

308 (d)

Both ether and chloroform are anaesthetic agents.


309 (b)

Picric acid is *sym*-trinitrophenol.




310 (d)

Among hydrogen halides, as the size of halide ion increases, its reactivity towards ethyl alcohol also increases. Thus, the order of reactivity of hydrogen halide is



312 (c)

Chloral forms chloral hydrate with water due to H-bonding.



314 (c)

Primary, secondary and tertiary alcohols are distinguished by Lucas test. A mixture of anhydrous ZnCl_2 + conc HCl is called Lucas reagent.

316 (b)

Sodium alkoxide is $R - \bar{\text{O}} \text{Na}^+$.

317 (a)

Due to the formation of stable tertiary carbon atom as an intermediate.

319 (a)


Reactivity order for H-atom of alcohol is,

Primary > Secondary > Tertiary



320 (b)

Phenol condenses with aliphatic and aromatic aldehydes in the *o*- and *p*-positions, the most important example being the condensation with formaldehyde. This is known as Lederer-Manasse reaction



322 (c)


C—O—C angle is 100° and thus, ethers $R-O-R$ have dipole moment.

323 (b)

It is a fact.

324 (b)

Presence of electron withdrawing group such as NO_2 , CHO etc, on benzene nucleus, makes phenol more acidic by stabilising phenoxide ion while presence of electron releasing groups such as $-\text{CH}_3$ $-\text{C}_2\text{H}_5$ destabilises the phenoxide ion, thus makes the phenol less acidic. Hence, the order of acidity of given compound is



327 (c)

Mixture of anhydrous ZnCl_2 and conc. HCl is known as Lucas reagent. Lucas test is used for the distinction between primary, secondary and tertiary alcohols.


The tertiary alcohol reacts immediately with Lucas reagent producing turbidity.

The secondary alcohol gives turbidity within 5-10 min and primary alcohol doesn't give turbidity at all at room temperature. In the given alternates, 2-hydroxy-2-methyl propane is 3° alcohol, so it is more reactive.

328 (b)

—OH gp. is on allyl gp. ($\text{CH}_2=\text{CH}-\text{CH}_2-$)


329 (c)




Due to strong H-bonding and weaker hydrophobic character.

351 (b)

Jones reagent oxidises 1° alcohols to aldehydes and 2° alcohols to ketones without affecting C = C double bond.




352 (b)



354 (d)

+R group present in phenol decreases the acidity while -R group presents at *ortho* or at *para* position increases the acidity of phenols. Thus, the correct order of acidity is



355 (b)

Due to the presence of two lone pair of electrons on oxygen atom.

356 (c)


__do__

357 (d)

The enzyme catalysed conversion of starch into sugar.


358 (c)

The process is called hydroboration and is used to convert terminal alkenes to 1-ol.



359 (c)

Sodium phenoxide reacts with CO₂ at 400 K and 4 – 7 atm pressure to give sodium salicylate. This is called Kolbe's reaction




361 (c)



Cannizzaro's reaction

362 (c)

It is aldol condensation.



363 (c)


Ethers have two alkyl groups on oxygen atom.

364 (a)

Pepsin hydrolyses —CONH— (peptide bonds) to —COOH and —NH₂.

365 (b)


1° alcohols are converted into aldehyde by reaction with Na₂Cr₂O₇ and H₂SO₄.



368 (d)


H-bonding in molecule gives rise to higher b.p.

369 (a)



This reaction is ethinylation. The catalyst used are copper acetylide or sod. alkoxide.

370 (b)




1° alcohol on oxidation gives aldehyde having same number of carbon and aldehyde on oxidation gives acid having same number of carbon atoms.

It means, D will be alcohol having two carbon atoms that is C₂H₅OH (ethyl alcohol) and the alcohol on oxidation will give CH₃CHO (acetaldehyde)



372 (d)

Alcohols (—OH) react with sodium and carbonyl

compounds  give precipitate with semicarbazide.

Since, the compound with molecular formula, C₃H₆O does not give precipitate with semicarbazide and does react with sodium, it is neither a carbonyl compound nor an alcohol. Hence, it must be an ether, *i.e.*, CH₂ = CHOCH₃


373 (c)

Williamson's synthesis is used for the preparation of ethers, specially mixed ethers.


374 (c)

An increase in hydrophobic character decreases H-bonding.

375 (a)



377 (b)




Note : Because of its special structure, there are two ether oxygen attached to same carbon, making it acetal.

378 (b)


CH_3OH is toxic and injurious to health and therefore also used for denaturation of alcohol.

384 (a)


Phenyl magnesium bromide reacts with *t*-butanol to produce benzene because phenyl group (electronegative group) is associated with active hydrogen of alcohol, *i.e.*, -H of -OH group of alcohol.



386 (b)



387 (b)



388 (b)

Phenols are acidic in nature due to resonance stabilisation of phenoxide ion. Presence of electrons releasing groups such as $-\text{CH}_3$ destabilises ion and decreases the acidic nature of phenols. On the other hand presence of electron withdrawing group in the ring stabilise phenoxide ion and increases the acidic nature of phenols. Further more *meta*-isomer is less acidic. Then *para* because it is stabilised by inductive effect only. Thus, correct order is $\text{IV} > \text{III} > \text{I} > \text{II}$


389 (c)

Also known as glyptal resin; A class of synthetic resin obtained by the reaction of polyhydric alcohol with poly basic organic acids or

anhydrides, e.g., Glycerol and phthalic anhydride, generally used for surface coating.

390 (b)


NaBH_4 , LiAlH_4 has no action on $\text{C}=\text{C}$.



391 (c)


The organic liquid *A* is $\text{C}_2\text{H}_5\text{OH}$

i. Ethyl alcohol is a colourless liquid with a characteristic pleasant smell, having boiling point 78.1°C .



(which decolourises Br_2 water and alk. KMnO_4)


392 (b)




Which gives iodoform test.

393 (d)

Reimer-Tiemann reaction this involves the treatment of phenol with chloroform in aqueous sodium hydroxide solution followed by acid hydrolysis. Salicylaldehyde is formed.



In the above reaction, chloroform first reacts with sodium hydroxide to produce dichloro carbene which is the intermediate in this reaction.





394 (b) Reactions involving H-atom of alcohol show the order $1^\circ > 2^\circ > 3^\circ$.

395 (a) A characteristic test for alcoholic gp.

397 (c) The boiling point of alcohols is higher than the boiling points of corresponding alkanes and aldehydes due to H-bonding. As the molecule mass increases, boiling point increases. Thus, C_2H_5OH has the higher boiling point among the given.

398 (c) When glycerol reacts with HI, *iso*-propyl iodide is obtained




is not soluble in NaHCO_3


400 (b) Pyroligneous acid is used for the preparation of acetic acid. It contains about 10% acetic acid, and was originally treated by neutralising with lime and then distilling off the volatile compounds like methanol and acetone

401 (a) Power alcohol is used to generate power.

403 (a) Ethyl alcohol reacts with methyl magnesium iodide as follows



405 (c) Due to resonance the phenoxide ion is more stable whereas resonance is not possible in alkoxide ion.



$R - O^\ominus$ no resonance is possible, Since, phenoxide ion is better stabilized by resonance, the phenol has more tendency to form phenoxide ion by releasing H^+ ion. So, phenols are acidic in nature.

407 (d) Molecular weight of diethyl ether is more than ethanol. Therefore, it should have higher boiling point than ethanol. But it is not so. It is due to intermolecular hydrogen bonding. Ethyl alcohol has intermolecular hydrogen bonding but diethyl ether has no hydrogen bonding. The compounds shows intermolecular hydrogen bonding has higher m.p. and b.p. than compounds having no hydrogen bonding. Therefore, the boiling point of diethyl ether will be less than ethanol (78°C).


409 (c)

$$\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{Acetyl bacilli}} \text{CH}_3\text{COOH}$$

410 (d)

$$\text{ROH} + \text{HOR} \xrightarrow{-\text{H}_2\text{O}} \text{ROR}$$

412 (a) Cumene is isopropyl benzene (1-methyl ethyl benzene). It on oxidation gives phenol.




414 (a)

Due to H-bonding.


415 (b)

Many a time unexpected products result during dehydration of alcohols.



417 (d)

Methanol reacts with salicylic acid in the presence of a few drops of conc H_2SO_4 to give methyl salicylate having the smell of oil of winter green.



418 (c)

Ethers are $R-O-R'$ or $R-O-R$.

419 (c)

Phenol gives characteristic colouration (violet) with aqueous FeCl_3 solution.

420 (a)

Reactivity of $-\text{OH}$ gp. of alcohols (due to +ve IE of alkyl group).

421 (d)

Remember these.

422 (d)


In presence of air and light, ether form peroxides which cause explosion during distillation

423 (d)


All the reaction proceed by stable ions. After the lose of H^+ ion, phenol forms phenoxide ion. The phenoxide ion is resonance stabilized, thus makes the phenol more acidic.

426 (c)


With mild oxidising agent like bromine water or H_2O_2 in the presence of FeSO_4 (Fenton's reagent), glycerol is oxidised to a mixture of glyceraldehyde and dihydroxy acetone



428 (c)




431 (a)



432 (a)


Oxalic acid on reaction with glycerol at 530K temperature furnish allyl alcohol.



433 (c)


Alcohol forms a azeotropic mixture with water and absolute alcohol is obtained by this mixture (rectified spirit) by adding benzene and then carrying out fractional distillation.

434 (a)



435 (b)


The reaction of alkyl halide with sodium alkoxide to give ether (alkoxy alkane) is known as Williamson's synthesis. In this reaction an ether (anisole) is prepared by the action of alkyl halide (methyl iodide) on sodium alkoxide (sodium phenate), so it is an example of Williamson's synthesis.




The reaction is called Fischer-Speier esterification.

465 (a)

When ethyl alcohol is oxidised by acidified potassium dichromate, CH_3COOH (Y) is obtained as




Carboxylic acid undergoes reduction with LiAlH_4 to give primary alcohol as




467 (d)

Reactivity order of OH towards Lucas reagent is, Tertiary > Secondary > Primary alcohol.

470 (c)



471 (a)



This reaction is an example of coupling reaction

472 (a)


H of CH_3OH (carbinol) is replaced by vinyl gp.

473 (d)

Alcohols which are used for generating power is called power alcohol. A mixture of 20% ethanol and 80% gasoline is used in internal combustion engines to derive power

474 (b)


When one H_2SO_4 reacts with ethyl alcohol at room temperature, ethyl hydrogen sulphate is formed



Ethyl hydrogen sulphate

477 (a)


Cyclohexanol on reaction with PBr_3 in presence of pyridine gives bromocyclohexane.



478 (c)

In Lucas test, when alcohol is mixed with conc HCl and anhydrous ZnCl_2 at room temperature, if oily product is formed immediately, the alcohol can be tertiary


479 (c)



Since, N being an element of second period can't contain more than 10 electrons in its valence shell

480 (c)

In presence of HNO_3 or alkaline KMnO_4



484 (b)

PCl_5 replaces $-\text{OH}$ group by $-\text{Cl}$.

486 (c)

Phenol forms azo dye, with benzene diazonium chloride. This reaction is called coupling reaction

487 (a)


Reactivity of H-atom of alcohol, (due to + IE of alkyl gp.)

Primary > Secondary > Tertiary.


Also CH_3OH is more acidic than $\text{C}_2\text{H}_5\text{OH}$ due to more +IE of $-\text{C}_2\text{H}_5$ gp.

489 (a)


Iso-butyl alcohol is secondary alcohol



490 (d)




491 (a)




492 (b)

2, 3-dimethyl butane-2, 3-diol is known as pinacol




494 (b)




495 (d)

$\text{C}_2\text{H}_5\text{OH}$ and $\text{C}_6\text{H}_5\text{OH}$ can be distinguished by neutral FeCl_3 solution or $\text{I}_2 + \text{NaOH}$ solution. $\text{C}_2\text{H}_5\text{OH}$ gives iodoform test with $\text{I}_2 + \text{NaOH}$ solution while phenol does not give yellow ppt. of iodoform.



$\text{C}_6\text{H}_5\text{OH}$ reacts with neutral FeCl_3 solution to give purple colour while $\text{C}_2\text{H}_5\text{OH}$ doesn't give any colour with neutral FeCl_3 solution.


497 (b)



(Reimer-Tiemann's reaction)

498 (c)


Compound 'X' ($\text{C}_7\text{H}_8\text{O}$) is insoluble in aqueous NaHCO_3 but soluble in NaOH , so it is a phenol. Since, the number of carbon atoms remains the same after bromination, the compound must be *meta* *cresol* and reactions takes place as follows



499 (b)

Phenol doesn't decompose sodium carbonate or sodium bicarbonate, *i.e.*, CO_2 is not evolved because phenol is a weaker acid than carbonic acid.

500 (d)



Unit attached to C or H in it to shown iodoform reaction.


501 (c)

Alcohol is used as solvent for many drugs.


502 (b)

In Victor's Meyer's test, 1° -alcohol gives red colour, 2° -alcohol gives blue colour while 3° -alcohol gives no colour.

503 (b)



504 (c)




505 (d)

All are anaesthetic agents.

506 (c)


Alcohols are oxidised by not copper to give aldehydes.



507 (c)

Bond angle is 110° due to steric hindrance of bulky alkyl groups.

513 (a)



This reaction is called Laderer Mannasse reaction.

514 (c)
do

517 (a)
CH₃SH is gas with foul smell and thus, mixed with LPG to detect its leakage.

518 (d)
Alcohols are neutral as they do not influence the pH. Due to O—H bond, they possess Bronsted acid nature showing cleavage of O—H bond. Also due to the presence of lone pair of electron on oxygen atom, they act as Lewis base. The reactivity order is based on +IE of alkyl groups.

Lewis base order : 3° > 2° > 1°

Bronsted acid order : 1° > 2° > 3°

519 (b)
Reduction of acid and acid derivatives producing alcohol by C₂H₅OH + Na is called Bonveault-Blanc reaction.

520 (b)
Absolute alcohol is 100% pure ethanol. The fractional distillation of aqueous solution of ethanol gives a constant boiling azeotropic mixture which contains 95% ethanol. To get 100% ethanol, a small amount of benzene is added with azeotropic mixture and then distilled. It is called azeotropic distillation.

522 (a)
Pyroligneous acid obtained during destructive distillation of wood contains mainly acetic acid (9-10 %), methyl alcohol (2-2.5%) and acetone about 0.5%; the other distillation products are wood gas, wood charcoal, wood tar.



523 (c)
Weak base reacts with strong acid.

525 (c)
Dunstan's test is used for identification of glycerol

529 (a)
(RMgX) + HCHO → 1° alcohol
Grignard reagent

(RMgX) + RCHO → 2° alcohol

(RMgX) + RCOR → 3° alcohol




530 (a)
Solubility of alcohols in water decreases as the size of alkyl group increases because tendency to form hydrogen bonding decreases. So, the order of solubility is as
Ethanol > *n*-propanol > *n*-butyl alcohol

531 (b)
Germinated barley called malt contains diastase enzyme.

533 (a)
Methyl phenyl ether is obtained by the reaction of phenolate ions and methyl iodine.
 $\text{C}_6\text{H}_5\text{O}^- + \text{CH}_3\text{I} \rightarrow \text{C}_6\text{H}_5\text{OCH}_3 + \text{I}^-$
Methyl phenyl ether


534 (b)
NaBH₄ and LiAlH₄ attacks only carbonyl group and reduce it into alcohol group. They do not attack on double bond.




535 (a)
Salicylic acid + NaHCO₃ → effervescence of CO₂
Phenol + NaHCO₃ → No reaction
∴ NaHCO₃ is used to distinguish between phenol and salicylic acid.

536 (d)
Both ether and chloroform are anaesthetic agents.

537 (b)
Ethyl chloride reacts with sodium ethoxide to form diethyl ether as




Diethyl ether is also obtained by reaction of ethyl alcohol with conc. H₂SO₄ at 140°C.




539 (c)

Conversion of -CHO group present in phenol ring into -OH in the presence of H_2O_2 is called Darkin reaction.



Conversion of amino acids into methyl α -acetamide ketones, when heated with acetic anhydride in pyridine solution is often referred to as the Darkin west reaction.



540 (d)


No one of the given reactions involve of formation of carbocation intermediate

541 (b)

Butanol-1, butanol-2, 2-methylpropanol-1, 2-methylpropanol-2.


542 (b)

Phenol on reaction with chloroform and KOH gives salicylaldehyde, which with 50% KOH solution undergoes Cannizaro's reaction.




543 (b)

In presence of NaOH or KOH, phenol reacts with alkyl halide and gives phenolic ether ($\text{C}_6\text{H}_5\text{OR}$).




Vapours of $\text{C}_6\text{H}_5\text{OH}$ and CH_3OH , with red hot ThO_2 (thoria) give anisole (phenolic ether).



anisole

544 (c)

Alkenes undergo addition reaction with diborane. The addition compounds on hydrolysis with $\text{H}_2\text{O}_2/\text{OH}^-$ yield alcohols



545 (d)

Glycerol is used as lubricant in watches.


546 (c)



547 (a)

Lucas reagent is anhyd. $\text{ZnCl}_2 + \text{HCl}$ (conc.) used to distinguish *p*, *s* and *t* alcohols.

548 (b)



549 (c)

Terylene is formed by the action of glycol ($\text{CH}_2\text{OHCH}_2\text{OH}$) on dimethyl terephthalate. It is also called dacron.

550 (c)

Yeast contains maltase, invertase, zymase enzymes.

551 (d)

General formula for alcohols is $\text{C}_n\text{H}_{2n+1}\text{OH}$. Primary alcohols have $-\text{CH}_2\text{OH}$ gp. Secondary alcohols have $>\text{CHOH}$ gp. and tertiary alcohols have

553 (d)


When phenol reacts with phthalic anhydride in presence of conc. H_2SO_4 and heated, then mixture is poured in NaOH solution the product formed is phenolphthalein.

555 (d)


In the presence of anhydrous ZnCl_2 , phenol form salicylaldehyde. It is Gattermann-aldehyde reaction.

556 (d)

Phenol gives Libermann's nitroso reaction.




This blue colour is formed due to the formation of




557 (c)

In rearrangement of cumene hydroperoxide



is not formed


579 (a)



Both shows H-bonding, however the increase in hydrophobic character (due to two alkyl groups in ether), the H-bonding weakens.

580 (a)

NaBH₄ reduces aldehyde to 1° alcohol.



CHEMISTRY (QUESTION BANK)**11.ALCOHOLS, PHENOLS AND ETHERS****Assertion - Reasoning Type**

This section contain(s) 0 questions numbered 1 to 0. Each question contains STATEMENT 1(Assertion) and STATEMENT 2(Reason). Each question has the 4 choices (a), (b), (c) and (d) out of which **ONLY ONE** is correct.

- a) Statement 1 is True, Statement 2 is True; Statement 2 **is** correct explanation for Statement 1
 b) Statement 1 is True, Statement 2 is True; Statement 2 **is not** correct explanation for Statement 1
 c) Statement 1 is True, Statement 2 is False
 d) Statement 1 is False, Statement 2 is True

1

Statement 1: Benzoic acid is stronger acid than phenol

Statement 2: Resonating structures of phenoxide ion has negative charge on less electronegative carbon atom

2

Statement 1:

Dehydration of $\text{CH}_3\text{CH}_2\overset{\text{OH}}{\text{CH}}\text{CH}_3$ gives $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ as the minor product and $\text{CH}_3\text{CH}=\text{CHCH}_3$ as the major product

Statement 2:

$\text{CH}_2=\text{CH}-\overset{\text{OH}}{\text{CH}}-\text{CH}_3$ can be dehydrated more readily than $\text{CH}_3\text{CH}_2\overset{\text{OH}}{\text{CH}}\text{CH}_3$

3

Statement 1: When glycerol is heated with hydrogen iodide, only allyl iodide is formed

Statement 2: With hydrogen iodide, glycerol forms three different compounds like propene, isopropyl iodide and allyl iodide

4

Statement 1: Ether have low dipole moments and sp^3 hybridised oxygen atom

Statement 2: Ethers behaves as Lewis base

5

Statement 1: Alcohols have higher boiling points than alkanes

Statement 2: O – H bond of alcohol is highly polarized

CHEMISTRY (QUESTION BANK)

11.ALCOHOLS, PHENOLS AND ETHERS

: ANSWER KEY :

- 1) a 2) b 3) d 4) b |
5) a


CHEMISTRY (QUESTION BANK)**11.ALCOHOLS, PHENOLS AND ETHERS****: HINTS AND SOLUTIONS :**

1 (a)


In phenoxide ion all the resonating structures except one have negative charge on less electronegative carbon atom while in benzoate ion, both the resonating forms have negative charge on more electronegative oxygen atom

3 (d)

When glycerol is heated with a small amount of HI, allyl iodide is formed



When glycerol is heated with a large amount of HI, the allyl iodide first formed is reduced to propene, which in presence of excess of HI forms isopropyl iodide



4 (b)

Due to the presence of lone pairs of electrons on the oxygen atom, ethers behave as Lewis bases. They dissolve in acids like conc H_2SO_4 and conc HCl to form oxonium salts



5 (a)

The high polarisation of O – H bond in alcohols gives rise to intramolecular hydrogen bonding, *ie*, molecules are brought nearer to each other and held together by attractive forces. No such hydrogen bonding exists in alkanes

Session : 2025-26

AS PER NEW NTA SYLLABUS

Total Questions : 58 I

CHEMISTRY (QUESTION BANK)

11.ALCOHOLS, PHENOLS AND ETHERS

Matrix-Match Type

This section contain(s) 0 question(s). Each question contains Statements given in 2 columns which have to be matched. Statements (A, B, C, D) in **columns I** have to be matched with Statements (p, q, r, s) in **columns II**.

1. Match the list given below

Column-I

Column- II

- | | |
|---|----------------------------|
| (A) Phenolic esters on heating in presence of AlCl_3 | (p) Claisen rearrangement |
| (B) Ary-allyl ethers are heated | (q) Houben-Hoesch reaction |
| (C) Polyhydroxy phenol undergo acylation with $\text{RNC} + \text{HCl}$ | (r) Fries rearrangement |

CODES :

- | | A | B | C | D |
|----|---|---|---|---|
| a) | a | b | c | |
| b) | c | b | a | |
| c) | b | c | a | |
| d) | c | a | b | |

CHEMISTRY (QUESTION BANK)

11.ALCOHOLS, PHENOLS AND ETHERS

: ANSWER KEY :

1) d

CHEMISTRY (QUESTION BANK)

11.ALCOHOLS, PHENOLS AND ETHERS

: HINTS AND SOLUTIONS :

1 (d)
In Fries rearrangement phenolic esters are heating in presence of AlCl_3

Claisen rearrangement-Ary-allyl ethers are heated.

Houben-Hoesch reaction- Polyhydroxy phenol undergo acylation with $\text{RNC} + \text{HCl}$