

## MATHS ( QUESTION BANK )

### 10. VECTOR ALGEBRA

#### Single Correct Answer Type

1. A unit vector in  $xy$ -plane that makes an angle  $45^\circ$  with the vector  $(\hat{i} + \hat{j})$  and an angle of  $60^\circ$  with the vector  $(3\hat{i} - 4\hat{j})$ , is
  - a)  $\hat{i}$
  - b)  $\frac{1}{\sqrt{2}}(\hat{i} - \hat{j})$
  - c)  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j})$
  - d) None of these
2. Let  $\vec{a}$  and  $\vec{b}$  be unit vectors inclined at an angle  $2\alpha (0 \leq \alpha \leq \pi)$  each other, then  $|\vec{a} + \vec{b}| < 1$ , if
  - a)  $\alpha = \frac{\pi}{2}$
  - b)  $\alpha < \frac{\pi}{3}$
  - c)  $\alpha > \frac{2\pi}{3}$
  - d)  $\frac{\pi}{3} < \alpha < \frac{2\pi}{3}$
3. The cartesian form of the plane  $\vec{r} = (s - 2t)\hat{i} + (3 - t)\hat{j} + (2s + t)\hat{k}$  is
  - a)  $2x - 5y - z - 15 = 0$
  - b)  $2x - 5y + z - 15 = 0$
  - c)  $2x - 5y - z + 15 = 0$
  - d)  $2x + 5y - z + 15 = 0$
4. If  $\vec{a} = 4\hat{i} + 6\hat{j}$  and  $\vec{b} = 3\hat{j} + 4\hat{k}$ , the vector form of the component of  $\vec{a}$  along  $\vec{b}$  is
  - a)  $\frac{18}{5}(3\hat{i} + 4\hat{k})$
  - b)  $\frac{18}{25}(3\hat{j} + 4\hat{k})$
  - c)  $\frac{36}{25}(3\hat{j} + 4\hat{k})$
  - d)  $\frac{19}{18}(2\hat{i} + 3\hat{j})$
5. A force  $\vec{F} = 2\hat{i} + \hat{j} - \hat{k}$  acts at a point  $A$ , whose position vectors is  $2\hat{i} - \hat{j}$ . The moment of  $\vec{F}$  about the origin is
  - a)  $\hat{i} + 2\hat{j} - 4\hat{k}$
  - b)  $\hat{i} - 2\hat{j} - 4\hat{k}$
  - c)  $\hat{i} + 2\hat{j} + 4\hat{k}$
  - d)  $\hat{i} - 2\hat{j} + 4\hat{k}$
6. If  $\vec{a}, \vec{b}, \vec{c}$  are linearly independent vectors, then  $\frac{(\vec{a} + 2\vec{b}) \times (2\vec{b} + \vec{c}) \cdot (5\vec{c} + \vec{a})}{\vec{a} \cdot (\vec{b} \times \vec{c})}$  is equal to
  - a) 10
  - b) 14
  - c) 18
  - d) 12
7. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are perpendicular to  $\vec{b} + \vec{c}, \vec{c} + \vec{a}$  and  $\vec{a} + \vec{b}$  respectively and if  $|\vec{a} + \vec{b}| = 6, |\vec{b} + \vec{c}| = 8$  and  $|\vec{c} + \vec{a}| = 10$ , then  $|\vec{a} + \vec{b} + \vec{c}|$  is equal to
  - a)  $5\sqrt{5}$
  - b) 50
  - c)  $10\sqrt{2}$
  - d) 10
8. If  $\vec{a}, \vec{b}, \vec{c}$  are three mutually perpendicular vectors of equal magnitude, then the angle  $\theta$  which  $\vec{a} + \vec{b} + \vec{c}$  makes with any one of three given vectors is given by
  - a)  $\cos^{-1} \frac{1}{\sqrt{3}}$
  - b)  $\cos^{-1} \frac{1}{3}$
  - c)  $\cos^{-1} \frac{2}{\sqrt{3}}$
  - d) None of these
9. Forces  $3 O\vec{A}, 5 O\vec{B}$  act along  $OA$  and  $OB$ . If their resultant passes through  $C$  on  $AB$ , then
  - a)  $C$  is a mid-point of  $AB$
  - b)  $C$  divides  $AB$  in the ratio  $2 : 1$
  - c)  $3 AC = 5 CB$
  - d)  $2 AC = 3 CB$
10. The centre of the circle given by  $\vec{r} \cdot (\hat{i} + 2\hat{j} + 2\hat{k}) = 15$  and  $\vec{r} \cdot (\hat{j} + 2\hat{k}) = 4$  is
  - a) (1,2,4)
  - b) (3,1,4)
  - c) (1,3,4)
  - d) None of these
11. Consider a tetrahedron with faces  $F_1, F_2, F_3, F_4$ . Let  $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$  be the vectors whose magnitudes are respectively equal to areas of  $F_1, F_2, F_3, F_4$  and whose directions are perpendicular to these faces in outward direction. Then,  $|\vec{v}_1 + \vec{v}_2 + \vec{v}_3 + \vec{v}_4|$  equals









74. The vectors  $\vec{a}(x) = \cos x \hat{i} + (\sin x)\hat{j}$  and  $\vec{b}(x) = x \hat{i} + \sin x \hat{j}$  are collinear for
- a) Unique value of  $x, 0 < x < \frac{\pi}{6}$                       b) Unique value of  $x, \frac{\pi}{6} < x < \frac{\pi}{3}$   
c) No value of  $x$     d) Infinitely many values of  $x, 0 < x < \frac{\pi}{2}$
75. A unit vector in  $xy$ -plane makes an angle of  $45^\circ$  with the vector  $\hat{i} + \hat{j}$  and an angle of  $60^\circ$  with the vector  $3\hat{i} - 4\hat{j}$  is
- a)  $\hat{i}$     b)  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$     c)  $\frac{\hat{i} - \hat{j}}{\sqrt{2}}$     d) None of these
76. The vector  $\vec{a}$  lies in the plane of vectors  $\vec{b}$  and  $\vec{c}$ , which of the following is correct
- a)  $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$                       b)  $\vec{a} \cdot \vec{b} \times \vec{c} = 1$                       c)  $\vec{a} \cdot \vec{b} \times \vec{c} = -1$                       d)  $\vec{a} \cdot \vec{b} \times \vec{c} = 3$
77. If the volume of parallelopiped with coterminous  $4\hat{i} + 5\hat{j} + \hat{k}$  and  $3\hat{i} - 9\hat{j} + p\hat{k}$  is 34 cu units, then  $p$  is equal to
- a) 4    b) -13    c) 13    d) 6
78. The value of  $\frac{(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2}{2\vec{a}^2\vec{b}^2}$  is
- a)  $\vec{a} \cdot \vec{b}$     b) 1    c) 0    d)  $\frac{1}{2}$
79. The magnitude of cross product of two vectors is  $\sqrt{3}$  times the dot product. The angle between the vectors is
- a)  $\frac{\pi}{6}$     b)  $\frac{\pi}{3}$     c)  $\frac{\pi}{2}$     d)  $\frac{\pi}{4}$
80. If  $G$  is the intersection of diagonals of a parallelogram  $ABCD$  and  $O$  is any point, then  $O\vec{A} + O\vec{B} + O\vec{C} + O\vec{D} =$
- a)  $2\vec{OG}$     b)  $4\vec{OG}$     c)  $5\vec{OG}$     d)  $3\vec{OG}$
81. If  $\vec{a} = (-1, 1, 1)$  and  $\vec{b} = (2, 0, 1)$ , then the vector  $\vec{X}$  satisfying the conditions  
(i) that it is coplanar with  $\vec{a}$  and  $\vec{b}$   
(ii) that it is perpendicular to  $\vec{b}$ , (iii) that  $\vec{a} \cdot \vec{X} = 7$  is,
- a)  $-3\hat{i} + 4\hat{j} + 6\hat{k}$                       b)  $-\frac{3}{2}\hat{i} + \frac{5}{2}\hat{j} + 3\hat{k}$                       c)  $3\hat{i} + 16\hat{j} - 6\hat{k}$                       d) None of these
82. If  $ABCDEF$  is a regular hexagon, then  $\vec{AC} + \vec{AD} + \vec{EA} + \vec{FA} =$
- a)  $2\vec{AB}$     b)  $3\vec{AB}$     c)  $\vec{AB}$     d)  $\vec{0}$
83.  $[(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c}) (\vec{b} \times \vec{c}) \times (\vec{c} \times \vec{a}) (\vec{c} \times \vec{a}) \times (\vec{a} \times \vec{b})]$  is equal to
- a)  $[\vec{a}\vec{b}\vec{c}]^2$     b)  $[\vec{a}\vec{b}\vec{c}]^3$     c)  $[\vec{a}\vec{b}\vec{c}]^4$     d) None of these
84. Suppose  $\vec{a} = \lambda \hat{i} - 7\hat{j} + 3\hat{k}$ ,  $\vec{b} = \lambda \hat{i} + \hat{j} + 2\lambda \hat{k}$ . If the angle between  $\vec{a}$  and  $\vec{b}$  is greater than  $90^\circ$ , then  $\lambda$  satisfies the inequality
- a)  $-7 < \lambda < 1$                       b)  $\lambda > 1$     c)  $1 < \lambda < 7$     d)  $-5 < \lambda < 1$
85. Let  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar vectors such that  $\vec{r}_1 = \vec{a} - \vec{b} + \vec{c}, \vec{r}_2 = \vec{b} + \vec{c} - \vec{a}, \vec{r}_3 = \vec{c} + \vec{a} + \vec{b}, \vec{r} = 2\vec{a} - 3\vec{b} + 4\vec{c}$   
If  $\vec{r} = \lambda_1\vec{r}_1 + \lambda_2\vec{r}_2 + \lambda_3\vec{r}_3$ , then
- a)  $\lambda_1 = 7$     b)  $\lambda_1 + \lambda_3 = 3$     c)  $\lambda_1 + \lambda_2 + \lambda_3 = 3$     d)  $\lambda_3 + \lambda_2 = 2$
86. If  $\vec{a}, \vec{b}, \vec{c}, \vec{d}$  are the position vectors of points  $A, B, C$  and  $D$  respectively such that  $(\vec{a} - \vec{d}) \cdot (\vec{b} - \vec{c}) = (\vec{b} - \vec{a}) \cdot (\vec{c} - \vec{d}) = 0$ , then  $D$  is the
- a) Centroid of  $\Delta ABC$   
b) Circumcentre of  $\Delta ABC$   
c) Orthocenter of  $\Delta ABC$   
d) None of these

87.  $A, B, C, D, E, F$  in that order, are the vertices of a regular hexagon with center origin. If the position vectors  $A$  and  $B$  are respectively,  $4\hat{i} + 3\hat{j} - \hat{k}$  and  $-3\hat{i} + \hat{j} + \hat{k}$ , then  $\overline{DE}$  is equal to  
 a)  $7\hat{i} + 2\hat{j} - 2\hat{k}$       b)  $-7\hat{i} - 2\hat{j} + 2\hat{k}$       c)  $3\hat{i} - \hat{j} - \hat{k}$       d)  $-4\hat{i} - 3\hat{j} + 2\hat{k}$
88. If  $|\vec{a} \times \vec{b}| = |\vec{a} \cdot \vec{b}|$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is  
 a)  $\Pi$       b)  $\frac{2\pi}{3}$       c)  $\frac{\pi}{4}$       d)  $\frac{\pi}{2}$
89. The ratio in which  $\hat{i} + 2\hat{j} + 3\hat{k}$  divides the join of  $-2\hat{i} + 3\hat{j} + 5\hat{k}$  and  $7\hat{i} - \hat{k}$  is  
 a) 2:1      b) 2:3      c) 3:4      d) 1:4
90. The values of  $x$  for which the angle between the vectors  $\vec{a} = x\hat{i} - 3\hat{j} - \hat{k}$  and  $\vec{b} = 2x\hat{i} - x\hat{j} - \hat{k}$  is acute and angle between  $\vec{b}$  and  $y$ -axis lies between  $\pi/2$  and  $\pi$  are  
 a)  $-1$       b) All  $x > 0$       c)  $1$       d) All  $x < 0$
91. The moment about the point  $M(-2, 4, -6)$  of the force represented in magnitude and position  $\overline{AB}$  where the points  $A$  and  $B$  have the coordinates  $(1, 2, -3)$  and  $(3, -4, 2)$  respectively, is  
 a)  $8\hat{i} - 9\hat{j} - 14\hat{k}$       b)  $2\hat{i} - 6\hat{j} + 5\hat{k}$       c)  $-3\hat{i} + 2\hat{j} - 3\hat{k}$       d)  $-5\hat{i} - 8\hat{j} - 8\hat{k}$
92. The angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{5\pi}{6}$  and the projection of  $\vec{a}$  in the direction of  $\vec{b}$  is  $\frac{-6}{\sqrt{3}}$ , then  $|\vec{a}|$  is equal to  
 a)  $6$       b)  $\frac{\sqrt{3}}{2}$       c)  $12$       d)  $4$
93. The equation of the line passing through the points  $a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  is  
 a)  $(a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$       b)  $(a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) - t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$   
 c)  $a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + (b_1\hat{i} + b_2\hat{j} + b_3\hat{k})t$       d) None of the above
94. The vector  $\vec{b} = 3\hat{i} + 4\hat{k}$  is to be written as the sum of a vector  $\vec{a} = \hat{i} + \hat{j}$  and a vector  $\vec{\beta}$  perpendicular to  $\vec{a}$ . Then  $\vec{a} =$   
 a)  $\frac{3}{2}(\hat{i} + \hat{j})$       b)  $\frac{2}{3}(\hat{i} + \hat{j})$       c)  $\frac{1}{2}(\hat{i} + \hat{j})$       d)  $\frac{1}{3}(\hat{i} + \hat{j})$
95. A parallelogram is constructed on  $3\vec{a} + \vec{b}$  and  $\vec{a} - 4\vec{b}$ , where  $|\vec{a}| = 6$  and  $|\vec{b}| = 8$  and  $\vec{a}$  and  $\vec{b}$  are anti-parallel, then the length of the longer diagonal is  
 a)  $40$       b)  $64$       c)  $42$       d)  $48$
96. If the vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  from the sides  $BC, CA$  and  $AB$  respectively of a triangle  $ABC$  then  
 a)  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$       b)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} = 0$   
 c)  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$       d)  $\vec{a} \times \vec{a} + \vec{a} \times \vec{c} + \vec{c} \times \vec{a} = 0$
97. The vectors  $\vec{a}$  and  $\vec{b}$  of equal magnitude 5 originating from a point and directs respectively towards north-east and north-west. Then, the magnitude of  $\vec{a} - \vec{b}$  is  
 a)  $3\sqrt{2}$       b)  $2\sqrt{3}$       c)  $2\sqrt{5}$       d)  $5\sqrt{2}$
98. If the vectors  $\vec{a} = \hat{i} + a\hat{j} + a^2\hat{k}, \vec{b} = \hat{i} + b\hat{j} + b^2\hat{k}$  and  $\vec{c} = \hat{i} + c\hat{j} + c^2\hat{k}$  are three non-coplanar vectors and  $\begin{vmatrix} a & a^2 & 1+a^3 \\ b & b^2 & 1+b^3 \\ c & c^2 & 1+c^3 \end{vmatrix} = 0$ , then the value of  $abc$  is  
 a)  $0$       b)  $1$       c)  $2$       d)  $-12$
99. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 3\hat{j} + 5\hat{k}$  and  $\vec{c} = 7\hat{i} + 9\hat{j} + 11\hat{k}$  then the area of parallelogram having diagonals  $\vec{a} + \vec{b}$  and  $\vec{b} + \vec{c}$  is  
 a)  $4\sqrt{6}$  sq units      b)  $\frac{1}{2}\sqrt{21}$  sq units      c)  $\frac{\sqrt{6}}{2}$  sq units      d)  $\sqrt{6}$  sq units
100. Let  $\vec{a} = \hat{i} + \hat{j} - \hat{k}, \vec{b} = \hat{i} - \hat{j} + \hat{k}$  and  $\vec{c}$  be a unit vector perpendicular to  $\vec{a}$  and coplanar with  $\vec{a}$  and  $\vec{b}$ , then it is given by

- a)  $\frac{1}{\sqrt{6}}(2\hat{i} - \hat{j} + \hat{k})$       b)  $\frac{1}{\sqrt{2}}(\hat{j} + \hat{k})$       c)  $\frac{1}{\sqrt{6}}(\hat{i} - 2\hat{j} + \hat{k})$       d)  $\frac{1}{2}(\hat{j} - \hat{k})$
101. If  $\vec{a} \cdot \hat{i} = 4$ , then  $(\vec{a} \times \hat{j}) \cdot (2\hat{j} - 3\hat{k}) =$   
a) 12      b) 2      c) 0      d) -12
102. If  $\vec{a} + 2\vec{b} + 4\vec{c} = \vec{0}$  and  $(\vec{a} \times \vec{b}) + (\vec{b} \times \vec{c}) + (\vec{c} \times \vec{a}) = \lambda(\vec{b} \times \vec{c})$ , then  $\lambda$  is equal to  
a) 4      b) 7      c) 8      d) 9
103. Forces acting on a particle have magnitude 5, 3 and 1 unit and act in the direction of the vectors  $6\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $3\hat{i} - 2\hat{j} + 6\hat{k}$  and  $2\hat{i} - 3\hat{j} - 6\hat{k}$  respectively. They remain constant while the particle is displaced from the points  $A(2, -1, -3)$  to  $B(5, -1, 1)$ . The work done is  
a) 11 unit      b) 33 unit      c) 10 unit      d) 30 unit
104. For any vector  $\vec{r}$ , the value of  $\hat{i} \times (\vec{r} \times \hat{i}) + \hat{j} \times (\vec{r} \times \hat{j}) + \hat{k} \times (\vec{r} \times \hat{k})$ , is  
a)  $\vec{0}$       b)  $2\vec{r}$       c)  $-2\vec{r}$       d) None of these
105. The vector equation of the plane passing through the origin and the line of intersection of the planes  $\vec{r} \cdot \vec{a} = \lambda$  and  $\vec{r} \cdot \vec{b} = \mu$ , is  
a)  $\vec{r} \cdot (\lambda \vec{a} - \mu \vec{b}) = 0$       b)  $\vec{r} \cdot (\lambda \vec{b} - \mu \vec{a}) = 0$       c)  $\vec{r} \cdot (\lambda \vec{a} + \mu \vec{b}) = 0$       d)  $\vec{r} \cdot (\lambda \vec{b} + \mu \vec{a}) = 0$
106. If  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar and  $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = k[\vec{a}, \vec{b}, \vec{c}]$ , then  $k$  is equal to  
a) 0      b) 1      c) 2      d) 3
107. If  $\vec{a} = (\hat{i} + \hat{j} + \hat{k})$ ,  $\vec{a} \cdot \vec{b} = 1$  and  $\vec{a} \times \vec{b} = \hat{j} - \hat{k}$ , then  $\vec{b}$  is  
a)  $\hat{i} - \hat{j} + \hat{k}$       b)  $2\hat{j} - \hat{k}$       c)  $\hat{i}$       d)  $2\hat{i}$
108. A tetrahedron has vertices at  $O(0,0,0)$ ,  $A(1,2,1)$ ,  $B(2,1,3)$  and  $C(-1,1,2)$ . Then, the angle between the faces  $OAB$  and  $ABC$  will be  
a)  $\cos^{-1}\left(\frac{19}{35}\right)$       b)  $\cos^{-1}\left(\frac{17}{31}\right)$       c)  $30^\circ$       d)  $90^\circ$
109. If  $2\vec{a} + 3\vec{b} - 5\vec{c} = \vec{0}$ , then ratio in which  $\vec{c}$  divides  $\overline{AB}$  is  
a) 3:2 internally      b) 3:2 externally      c) 2:3 internally      d) 2:3 externally
110. The perimeter of the triangle whose vertices have the position vectors  $\hat{i} + \hat{j} + \hat{k}$ ,  $5\hat{i} + 3\hat{j} - 3\hat{k}$  and  $2\hat{i} + 5\hat{j} + 9\hat{k}$  is given by  
a)  $15 + \sqrt{157}$       b)  $15 - \sqrt{157}$       c)  $\sqrt{15} + \sqrt{157}$       d)  $\sqrt{15} - \sqrt{157}$
111. If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = 5$ ,  $|\vec{b}| = 4$  and  $|\vec{c}| = 3$ , then the value of  $|\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}|$  is  
a) 25      b) 50      c) -25      d) 20
112. If  $\vec{a}$  is any vector, then  $(\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2 =$   
a)  $\vec{a}^2$       b)  $2\vec{a}^2$       c)  $3\vec{a}^2$       d)  $4\vec{a}^2$
113. If  $\vec{a}, \vec{b}, \vec{c}$  are three non-zero vectors (no two of which are collinear), such that the pairs of vectors  $(\vec{a} + \vec{b}, \vec{c})$  and  $(\vec{b} + \vec{c}, \vec{a})$  are collinear, then  $\vec{a} + \vec{b} + \vec{c} =$   
a)  $\vec{a}$       b)  $\vec{b}$       c)  $\vec{c}$       d)  $\vec{0}$
114. Let  $\vec{a}, \vec{b}, \vec{c}$  be three non-coplanar vectors and  $\vec{r}$  be any vector in space such that  $\vec{r} \cdot \vec{a} = 1$ ,  $\vec{r} \cdot \vec{b} = 2$  and  $\vec{r} \cdot \vec{c} = 3$ . If  $[\vec{a}, \vec{b}, \vec{c}] = 1$ , then  $\vec{r}$  is equal to  
a)  $\vec{a} + 2\vec{b} + 3\vec{c}$       b)  $\vec{b} \times \vec{c} + 2\vec{c} \times \vec{a} + 3\vec{a} \times \vec{b}$   
c)  $(\vec{b} \cdot \vec{c})\vec{a} + 2(\vec{c} \cdot \vec{a})\vec{b} + 3(\vec{a} \cdot \vec{b})\vec{c}$       d) None of these
115. If  $\vec{x} + \vec{y} + \vec{z} = \vec{0}$ ,  $|\vec{x}| = |\vec{y}| + |\vec{z}| = 2$ , and  $\theta$  is angle between  $\vec{y}$  and  $\vec{z}$ , then the value of  $\operatorname{cosec}^2\theta + \cot^2\theta$  is equal to  
a)  $4/3$       b)  $5/3$       c)  $1/3$       d) 1
116. If  $\vec{a} \cdot \vec{b} = -|\vec{a}||\vec{b}|$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is  
a)  $45^\circ$       b)  $180^\circ$       c)  $90^\circ$       d)  $60^\circ$





- a)  $\frac{a\vec{a} + b\vec{b} + c\vec{c}}{a + b + c}$       b)  $\frac{a\vec{a} + b\vec{b} + c\vec{c}}{\sqrt{a^2 + b^2 + c^2}}$       c)  $\frac{1}{3}[\vec{a} + \vec{b} + \vec{c}]$       d)  $\frac{\vec{a} + \vec{b} + \vec{c}}{a + b + c}$
146. If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then which of the following values of  $\vec{a} \cdot \vec{b}$  is not possible?  
a)  $\sqrt{3}$       b)  $\sqrt{3}/2$       c)  $1/\sqrt{2}$       d)  $-1/2$
147. The two vectors  $\{\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = 4\hat{i} - \lambda\hat{j} + 6\hat{k}\}$  are parallel, if  $\lambda$  is equal to  
a) 2      b) -3      c) 3      d) -2
148. Force acting on a particle have magnitude 5,3 and 1 unit act in the direction of the vectors  $6\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $3\hat{i} - 2\hat{j} + 6\hat{k}$  and  $2\hat{i} - 3\hat{j} - 6\hat{k}$  respectively. They remain constant while the particle is displaced from the point  $A(2, -1, -3)$  to  $B(5, -1, 1)$ . The work done is  
a) 11 units      b) 33 units      c) 10 units      d) 30 units
149. If  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$  and  $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$ , then  
a) Either  $\vec{a} = \vec{0}$  or  $\vec{b} = \vec{c}$       b)  $\vec{a} \parallel (\vec{b} - \vec{c})$       c)  $\vec{a} \perp (\vec{b} - \vec{c})$       d) None of these
150. The two vectors  $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = 4\hat{i} - \lambda\hat{j} + 6\hat{k}$  are parallel if  $\lambda =$   
a) 2      b) -3      c) 3      d) -2
151. If  $\vec{a}, \vec{b}, \vec{c}$  are unit coplanar vectors, then  $[2\vec{a} - \vec{b}, 2\vec{b} - \vec{c}, 2\vec{c} - \vec{a}]$  is equal to  
a) 1      b) 0      c)  $-\sqrt{3}$       d)  $\sqrt{3}$
152. The angle between the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  when  $\vec{a} = (1, 1, 4)$  and  $\vec{b} = (1, -1, 4)$  is  
a)  $45^\circ$       b)  $90^\circ$       c)  $15^\circ$       d)  $30^\circ$
153. Let  $P(3, 2, 6)$  be a point in space and  $Q$  be a point on the line  $\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(-3\hat{i} + \hat{j} + 5\hat{k})$ . Then, the value of  $\mu$  for which the vector  $\overrightarrow{PQ}$  is parallel to the plane  $x - 4y + 3z = 1$  is  
a)  $\frac{1}{4}$       b)  $-\frac{1}{4}$       c)  $\frac{1}{8}$       d)  $-\frac{1}{8}$
154. The area of triangle having vertices as  $\hat{i} - 2\hat{j} + 3\hat{k}, -2\hat{i} + 3\hat{j} - \hat{k}, 4\hat{i} - 7\hat{j} + 7\hat{k}$  is  
a) 36 sq units      b) 0 sq units      c) 39 sq units      d) 11 sq units
155. If  $\vec{r} \times \vec{a} = \vec{b} \times \vec{a}; \vec{r} \times \vec{b} = \vec{a} \times \vec{b}; \vec{a} \neq \vec{0}; \vec{b} \neq \vec{0}; \vec{a} \neq \lambda \vec{b}, \vec{a}$  is not perpendicular to  $\vec{b}$ , then  $\vec{r} =$   
a)  $\vec{a} - \vec{b}$       b)  $\vec{a} + \vec{b}$       c)  $\vec{a} \times \vec{b} + \vec{a}$       d)  $\vec{a} \times \vec{b} + \vec{b}$
156. If  $\vec{a} + \vec{b} + \vec{c}$  are three unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , where  $\vec{0}$  is null vector, then  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is  
a) -3      b) -2      c)  $-\frac{3}{2}$       d) 0
157. The edges of a parallelepiped are unit length and are parallel to non-coplanar unit vectors  $\vec{a}, \vec{b}, \vec{c}$  such that  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = \frac{1}{2}$ . Then, the volume of the parallelepiped is  
a)  $\frac{1}{\sqrt{2}}$  cu unit      b)  $\frac{1}{2\sqrt{2}}$  cu unit      c)  $\frac{\sqrt{3}}{2}$  cu unit      d)  $\frac{1}{\sqrt{3}}$  cu unit
158. If  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{b} = \hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k})$ , then length of  $\vec{b}$  is equal to  
a)  $\sqrt{12}$       b)  $2\sqrt{12}$       c)  $3\sqrt{14}$       d)  $2\sqrt{14}$
159. A vector  $\vec{a}$  has components  $2p$  and  $1$  with respect to a rectangular cartesian system. This system is rotated through a certain angle about the origin in the counter clockwise sense, if this respect to new system  $\vec{a}$  has components  $p + 1$  and  $1$ , then  
a)  $p = 0$       b)  $p = 1$  or  $p = \frac{-1}{2}$       c)  $p = -1$       d)  $p = 1$  or  $p = -1$
160. If the vectors  $\vec{r}_1 = a\hat{i} + \hat{j} + \hat{k}, \vec{r}_2 = \hat{i} + b\hat{j} + \hat{k}, \vec{r}_3 = \hat{i} + \hat{j} + c\hat{k} (a \neq 1, b \neq 1, c \neq 1)$  are coplanar, then the value of  $\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c}$ , is  
a) -1      b) 0      c) 1      d) None of these
161. A non-zero vectors  $\vec{a}$  is such that its projection along the vectors  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$  and  $\frac{-\hat{i} + \hat{j}}{\sqrt{2}}$  and  $\hat{k}$  are equal, then unit vector along  $\vec{a}$  is

$$a) \frac{\sqrt{2}\hat{j} - \hat{k}}{\sqrt{3}}$$

$$b) \frac{\hat{j} - \sqrt{2}\hat{k}}{\sqrt{3}}$$

$$c) \frac{\sqrt{2}}{\sqrt{3}}\hat{j} + \frac{\hat{k}}{\sqrt{3}}$$

$$d) \frac{\hat{j} - \hat{k}}{\sqrt{2}}$$

162. Let  $P, Q, R$  and  $S$  be the points on the plane with position vectors  $-2\hat{i} - \hat{j}, 4\hat{i}, 3\hat{i} + 3\hat{j}$  and  $-3\hat{i} + 2\hat{j}$  respectively. The quadrilateral  $PQRS$  must be

- a) Parallelogram, which is neither a rhombus nor a rectangle  
 b) Square  
 c) Rectangle, but not a square  
 d) Rhombus, but not a square

163. If  $\vec{a}, \vec{b}, \vec{c}$  are linearly independent vectors and  $\Delta = \begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{a} \cdot \vec{c} & \vec{b} \cdot \vec{c} & \vec{c} \cdot \vec{c} \end{vmatrix}$ , then

- a)  $\Delta = 0$   
 b)  $\Delta = 1$   
 c)  $\Delta =$  any non-zero value  
 d) None of these

164. If  $\vec{a} = -2\hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 5\hat{j}$  and  $\vec{c} = 4\hat{i} + 4\hat{j} - 2\hat{k}$ , then the projection of  $3\vec{a} - 2\vec{b}$  on the axis of the vector  $\vec{c}$  is

- a) 11                                      b) -11                                      c) 33                                      d) -33

165. A tetrahedron has vertices at  $O(0, 0), A(1, 2, 1), B(2, 1, 3)$  and  $C(-1, 1, 2)$ . Then, the angle between the faces  $OAB$  and  $ABC$  will be

- a)  $\cos^{-1}\left(\frac{19}{35}\right)$                                       b)  $\cos^{-1}\left(\frac{7}{31}\right)$                                       c)  $30^\circ$                                       d)  $90^\circ$

166. If  $\vec{a} + 2\vec{b} + 3\vec{c} = \vec{0}$  and  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$  is equal to  $\lambda(\vec{b} \times \vec{c})$ , then  $\lambda =$

- a) 3                                      b) 4                                      c) 5                                      d) None of these

167.  $\vec{a} \times [\vec{a} \times (\vec{a} \times \vec{b})]$  is equal to

- a)  $(\vec{a} \times \vec{a}) \cdot (\vec{b} \times \vec{a})$                                       b)  $\vec{a} \cdot (\vec{b} \times \vec{a}) - \vec{b} \cdot (\vec{a} \times \vec{b})$   
 c)  $[\vec{a} \cdot (\vec{a} \times \vec{b})]\vec{a}$                                       d)  $(\vec{a} \cdot \vec{a})(\vec{b} \times \vec{a})$

168. If  $\vec{a}$  is a unit vector such that  $\vec{a} \times (\hat{i} + 2\hat{j} + \hat{k}) = \hat{i} - \hat{k}$ , then  $\vec{a} =$

- a)  $-\frac{1}{3}(2\hat{i} + \hat{j} + 2\hat{k})$                                       b)  $\hat{j}$                                       c)  $\frac{1}{3}(\hat{i} + 2\hat{j} + 2\hat{k})$                                       d)  $\hat{i}$

169. The median  $AD$  of the triangle  $ABC$  is bisected at  $E, BE$  meets  $AC$  in  $F$ , then  $AF:AC =$

- a)  $3/4$                                       b)  $1/3$                                       c)  $1/2$                                       d)  $1/4$

170. Vectors  $\vec{a}$  and  $\vec{b}$  are inclined at an angle  $\theta = 120^\circ$ . If  $|\vec{a}| = 1, |\vec{b}| = 2$ , then  $[(\vec{a} + 3\vec{b}) \times (3\vec{a} + \vec{b})]^2$  is equal to

- a) 190                                      b) 275                                      c) 300                                      d) 192

171. If  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar vectors, then  $(\vec{a} + \vec{b} + \vec{c}) \cdot [(\vec{a} + \vec{b}) \times (\vec{a} \times \vec{c})]$  is

- a) 0                                      b)  $2[\vec{a} \cdot \vec{b} \cdot \vec{c}]$                                       c)  $-\vec{a} \cdot \vec{b} \cdot \vec{c}$                                       d)  $[\vec{a} \cdot \vec{b} \cdot \vec{c}]$

172. If  $\vec{a} = \hat{i} + \hat{j} - \hat{k}, \vec{b} = \hat{i} - \hat{j} + \hat{k}$  and  $\vec{c}$  is a unit vector perpendicular to the vector  $\vec{a}$  and coplanar with  $\vec{a}$  and  $\vec{b}$ , then a unit vector  $\vec{d}$  perpendicular to both  $\vec{a}$  and  $\vec{c}$  is

- a)  $\frac{1}{\sqrt{6}}(2\hat{i} - \hat{j} + \hat{k})$                                       b)  $\frac{1}{\sqrt{2}}(\hat{j} + \hat{k})$                                       c)  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j})$                                       d)  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{k})$

173. If  $G$  is the centroid of the  $\Delta ABC$ , then  $\vec{GA} + \vec{BG} + \vec{GC}$  is equal to

- a)  $2\vec{GB}$                                       b)  $2\vec{GA}$                                       c)  $\vec{0}$                                       d)  $2\vec{BG}$

174. A non-zero vector  $\vec{a}$  is parallel to the line of intersection of the plane determined by vectors  $\hat{i}, \hat{i} - \hat{j}$  and the plane determined by the vectors  $\hat{i} + \hat{j}, \hat{i} - \hat{k}$ . The angle between  $\vec{a}$  and  $\hat{i} + 2\hat{j} - 2\hat{k}$  is

- a)  $\frac{\pi}{3}$                                       b)  $\frac{\pi}{6}$                                       c)  $\frac{\pi}{4}$                                       d) None of these

175. If  $|\vec{a}| = 5, |\vec{b}| = 6$  and  $\vec{a} \cdot \vec{b} = -25$ , then  $|\vec{a} \times \vec{b}|$  is equal to

- a) 25                                      b)  $6\sqrt{11}$                                       c)  $11\sqrt{5}$                                       d)  $5\sqrt{11}$





206. The angle between the straight lines  $\vec{r} = (2 - 3t)\hat{i} + (1 + 2t)\hat{j} + (2 + 6t)\hat{k}$  and  $\vec{r} = (1 + 4s)\hat{i} + (2 - s)\hat{j} + (8s - 1)\hat{k}$  is
- a)  $\cos^{-1}\left(\frac{\sqrt{41}}{34}\right)$       b)  $\cos^{-1}\left(\frac{21}{34}\right)$       c)  $\cos^{-1}\left(\frac{43}{63}\right)$       d)  $\cos^{-1}\left(\frac{34}{63}\right)$
207. A vector which makes equal angles with the vectors  $\frac{1}{3}(\hat{i} - 2\hat{j} + 2\hat{k})$ ,  $\frac{1}{5}(-4\hat{i} - 3\hat{k})$  and  $\hat{j}$  is
- a)  $5\hat{i} + \hat{j} + 5\hat{k}$       b)  $-5\hat{i} + \hat{j} + 5\hat{k}$       c)  $5\hat{i} - \hat{j} + 5\hat{k}$       d)  $5\hat{i} + \hat{j} - 5\hat{k}$
208. In a  $\Delta ABC$ , if  $\vec{AB} = \hat{i} - 7\hat{j} + \hat{k}$  and  $\vec{BC} = 3\hat{i} + \hat{j} + 2\hat{k}$ , then  $|\vec{CA}| =$
- a)  $\sqrt{61}$       b)  $\sqrt{52}$       c)  $\sqrt{51}$       d)  $\sqrt{41}$
209. If  $\hat{i}, \hat{j}, \hat{k}$  are unit orthonormal vectors and  $\vec{a}$  is a vector, if  $\vec{a} \times \vec{r} = \hat{j}$ , then  $\vec{a} \cdot \vec{r}$  is
- a) 0      b) 1      c) -1      d) Arbitrary scalar
210. If the scalar product of the vector  $\hat{i} + \hat{j} + 2\hat{k}$  with the unit vector along  $m\hat{i} + 2\hat{j} + 3\hat{k}$  is equal to 2, then one of the value of  $m$  is
- a) 3      b) 4      c) 5      d) 6
211. Let  $\vec{a}$  and  $\vec{b}$  are non-collinear vectors. If there exists scalars  $\alpha, \beta$  such that  $\alpha\vec{a} + \beta\vec{b} = \vec{0}$ , then
- a)  $\alpha = \beta \neq 0$       b)  $\alpha + \beta = 0$       c)  $\alpha = \beta = 0$       d)  $\alpha \neq \beta$
212. The vector  $\vec{a} = \hat{i} + \hat{j} + m\hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j} + (m + 1)\hat{k}$  and  $\vec{c} = \hat{i} - \hat{j} + m\hat{k}$  are coplanar, if  $m$  is equal to
- a) 1  
b) 4  
c) 3  
d) No value of  $m$  for which vectors are coplanar
213. The unit vector in  $XOY$  plane and making angles  $45^\circ$  and  $60^\circ$  respectively with  $\vec{a} = 2\hat{i} + 2\hat{j} - \hat{k}$  and  $b = 0\hat{i} + \hat{j} - \hat{k}$ , is
- a)  $-\frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{k}$   
b)  $\frac{1}{\sqrt{2}}\hat{i} - \frac{1}{\sqrt{2}}\hat{k}$   
c)  $\frac{1}{3\sqrt{2}}\hat{i} + \frac{4}{3\sqrt{2}}\hat{j} + \frac{1}{3\sqrt{2}}\hat{k}$   
d) None of these
214. The value of  $\lambda$ , for which the four points  $2\hat{i} + 3\hat{j} - \hat{k}$ ,  $\hat{i} - 2\hat{j} + 3\hat{k}$ ,  $3\hat{i} + 4\hat{j} - 2\hat{k}$ ,  $\hat{i} - 6\hat{j} + \lambda\hat{k}$  are coplanar, is
- a) 2      b) 4      c) 6      d) 8
215. If  $|\vec{a}| = |\vec{b}|$ , then
- a)  $(\vec{a} + \vec{b})$  is parallel to  $\vec{a} - \vec{b}$   
b)  $\vec{a} + \vec{b}$  is  $\perp$  to  $\vec{a} - \vec{b}$   
c)  $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 2|\vec{a}|^2$   
d) None of these
216. The area of a parallelogram whose adjacent sides are given by the vectors  $\hat{i} + 2\hat{j} + 3\hat{k}$  and  $-3\hat{i} - 2\hat{j} + \hat{k}$  (in sq unit), is
- a)  $\sqrt{180}$  sq unit      b)  $\sqrt{140}$  sq unit      c)  $\sqrt{80}$  sq unit      d)  $\sqrt{40}$  sq unit
217. If  $P$  is any point with in a triangle  $ABC$ , then  $\vec{PA} + \vec{CP}$  is equal to
- a)  $\vec{AC} + \vec{CB}$       b)  $\vec{BC} + \vec{BA}$       c)  $\vec{CB} + \vec{AB}$       d)  $\vec{CB} + \vec{BA}$
218. Let the unit vectors  $\vec{a}$  and  $\vec{b}$  be perpendicular to each other and the unit vector  $\vec{c}$  be inclined at an angle  $\theta$  to both  $\vec{a}$  and  $\vec{b}$ . If  $\vec{c} = x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})$ , then
- a)  $x = \cos \theta, y = \sin \theta, z = \cos 2\theta$   
b)  $x = \sin \theta, y = \cos \theta, z = -\cos 2\theta$   
c)  $x = y = \cos \theta, z^2 = \cos 2\theta$

- d)  $x = y = \cos \theta, z^2 = -\cos 2\theta$
219. If  $\vec{a}, \vec{b}, \vec{c}$  are vectors such that  $\vec{a} \cdot \vec{b} = 0$  and  $\vec{a} + \vec{b} = \vec{c}$ , then  
 a)  $|\vec{a}|^2 + |\vec{b}|^2 = |\vec{c}|^2$       b)  $|\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2$       c)  $|\vec{b}|^2 = |\vec{a}|^2 = |\vec{c}|^2$       d) None of these
220. If  $OACB$  is a parallelogram with  $\vec{OC} = \vec{a}$  and  $\vec{AB} = \vec{b}$ , then  $\vec{OA} =$   
 a)  $\vec{a} + \vec{b}$       b)  $\vec{a} - \vec{b}$       c)  $\frac{1}{2}(\vec{b} - \vec{a})$       d)  $\frac{1}{2}(\vec{a} - \vec{b})$
221. Five points given by  $A, B, C, D, E$  are in plane. Three forces  $\vec{AC}, \vec{AD}$  and  $\vec{AE}$  act at  $A$  and three forces  $\vec{CB}, \vec{DB}, \vec{EB}$  act at  $B$ . Then, their resultant is  
 a)  $2\vec{AC}$       b)  $3\vec{AB}$       c)  $3\vec{DB}$       d)  $2\vec{BC}$
222. The vector  $\vec{a} = \alpha\hat{i} + 2\hat{j} + \beta\hat{k}$  lies in the plane of the vectors  $\vec{b} = \hat{i} + \hat{j}$  and  $\vec{c} = \hat{j} + \hat{k}$  and bisects the angle between  $\vec{b}$  and  $\vec{c}$ . Then, which one of the following gives possible value of  $\alpha$  and  $\beta$ ?  
 a)  $\alpha=1, \beta=1$       b)  $\alpha=2, \beta=2$       c)  $\alpha=1, \beta=2$       d)  $\alpha=2, \beta=1$
223. A unit vector perpendicular to the plane of  $\vec{a} = 2\hat{i} - 6\hat{j} - 3\hat{k}, \vec{b} = 4\hat{i} + 3\hat{j} - \hat{k}$  is  
 a)  $\frac{4\hat{i} + 3\hat{j} - \hat{k}}{\sqrt{26}}$       b)  $\frac{2\hat{i} - 6\hat{j} - 3\hat{k}}{7}$       c)  $\frac{3\hat{i} - 2\hat{j} + 6\hat{k}}{7}$       d)  $\frac{2\hat{i} - 3\hat{j} - 6\hat{k}}{7}$
224. Vectors  $\vec{a}$  and  $\vec{b}$  are inclined at angle  $\theta = 120^\circ$ . If  $|\vec{a}| = 1, |\vec{b}| = 2$ , then  $[(\vec{a} + 3\vec{b}) \times (3\vec{a} - \vec{b})]^2$  is equal to  
 a) 300      b) 325      c) 275      d) 225
225. If  $\vec{a} \cdot \hat{i} = 4$  then  $(\vec{a} \times \hat{j}) \cdot (2\hat{j} - 3\hat{k})$  is equal to  
 a) 12      b) 2      c) 0      d) -12
226. The volume (in cubic unit) of the tetrahedron with edges  $\hat{i} + \hat{j} + \hat{k}, \hat{i} - \hat{j} + \hat{k}$  and  $\hat{i} + 2\hat{j} - \hat{k}$  is  
 a) 4      b)  $\frac{2}{3}$       c)  $\frac{1}{6}$       d)  $\frac{1}{3}$
227. If  $|\vec{a} \times \vec{b}| = 4, |\vec{a} \cdot \vec{b}| = 2$ , then  $|\vec{a}|^2 + |\vec{b}|^2 =$   
 a) 6      b) 2      c) 20      d) 8
228. If  $\vec{a}, \vec{b}, \vec{c}$  be three unit vectors such that  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{1}{2}\vec{b}, \vec{b}$  and  $\vec{c}$  being non-parallel. If  $\theta_1$  is the angle between  $\vec{a}$  and  $\theta_2$  is the angle between  $\vec{a}$  and  $\vec{c}$ , then  
 a)  $\theta_1 = \frac{\pi}{6}, \theta_2 = \frac{\pi}{3}$       b)  $\theta_1 = \frac{\pi}{3}, \theta_2 = \frac{\pi}{6}$       c)  $\theta_1 = \frac{\pi}{2}, \theta_2 = \frac{\pi}{3}$       d)  $\theta_1 = \frac{\pi}{3}, \theta_2 = \frac{\pi}{2}$
229. If  $P, Q, R$  are the mid-points of the sides  $AB, BC$  and  $CA$  of  $\Delta ABC$  are  $O$  is a point within the triangle, then  $\vec{OA} + \vec{OB} + \vec{OC} =$   
 a)  $2(\vec{OP} + \vec{OQ} + \vec{OR})$       b)  $\vec{OP} + \vec{OQ} + \vec{OR}$       c)  $4(\vec{OP} + \vec{OQ} + \vec{OR})$       d)  $6(\vec{OP} + \vec{OQ} + \vec{OR})$
230.  $(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2$  is equal to  
 a)  $\vec{a}^2 \vec{b}^2$       b)  $\vec{a}^2 + \vec{b}^2$       c) 1      d)  $2\vec{a} \cdot \vec{b}$
231. If  $\vec{a}$  is a vector of magnitude 50, collinear with the vector  $\vec{b} = 6\hat{i} - 8\hat{j} - \frac{15}{2}\hat{k}$  and makes an acute angle with the positive direction of z-axis, then  $\vec{a}$  is equal to  
 a)  $-24\hat{i} + 32\hat{j} + 30\hat{k}$       b)  $24\hat{i} - 32\hat{j} - 30\hat{k}$       c)  $12\hat{i} - 16\hat{j} - 15\hat{k}$       d)  $-12\hat{i} + 16\hat{j} - 15\hat{k}$
232. If  $ABCDEF$  is a regular hexagon with  $\vec{AB} = \vec{a}$  and  $\vec{BC} = \vec{b}$ , then  $\vec{CE}$  equals  
 a)  $\vec{b} - \vec{a}$       b)  $-\vec{b}$       c)  $\vec{b} - 2\vec{a}$       d)  $\vec{b} + \vec{a}$
233. If the vectors  $2\hat{i} - 3\hat{j} + 4\hat{k}$  and  $\hat{i} + 2\hat{j} - \hat{k}$  and  $m\hat{i} - \hat{j} + 2\hat{k}$  are coplanar, then the value of  $m$  is  
 a)  $\frac{5}{8}$       b)  $\frac{8}{5}$       c)  $-\frac{7}{4}$       d)  $\frac{2}{3}$
234. If  $\vec{a}, \vec{b}, \vec{c}$  are the three vectors mutually perpendicular to each other to form a right handed system and  $|\vec{a}| = 1, |\vec{b}| = 3$  and  $|\vec{c}| = 5$ , then  $[\vec{a} - 2\vec{b}, \vec{b} - 3\vec{c}, \vec{c} - 4\vec{a}]$  is equal to  
 a) 0      b) -24      c) 3600      d) -215
235. The value of  $\hat{i} \times (\hat{j} \times \hat{k}) + \hat{j} \times (\hat{k} \times \hat{i}) + \hat{k} \times (\hat{i} \times \hat{j})$  is







- c)  $\vec{r} = (2k, 4k, 5k)k \in R$  d) None of these
278. The vector  $\vec{a}$  coplanar with the vectors  $\hat{i}$  and  $\hat{j}$ , perpendicular to the vector  $\vec{b} = 4\hat{i} - 3\hat{j} + 5\hat{k}$  such that  $|\vec{a}| = |\vec{b}|$  is  
a)  $\sqrt{2}(3\hat{i} + 4\hat{j})$  or,  $-\sqrt{2}(3\hat{i} + 4\hat{j})$   
b)  $\sqrt{2}(4\hat{i} + 3\hat{j})$  or,  $-\sqrt{2}(4\hat{i} + 3\hat{j})$   
c)  $\sqrt{3}(4\hat{i} + 5\hat{j})$  or,  $-\sqrt{3}(4\hat{i} + 5\hat{j})$   
d)  $\sqrt{3}(5\hat{i} + 4\hat{j})$  or,  $-\sqrt{3}(5\hat{i} + 4\hat{j})$
279. Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be vectors with magnitude 3, 4 and 5 respectively and  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , then the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is  
a) 47 b) 25 c) 50 d) -25
280. If  $\vec{a}, \vec{b}, \vec{c}$  are the position vectors of the vertices of an equilateral triangle, whose orthocenter is at the origin, then  
a)  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  b)  $\vec{a}^2 = \vec{b}^2 + \vec{c}^2$  c)  $\vec{a} + \vec{b} = \vec{c}$  d) None of these
281. If  $4\hat{i} + 7\hat{j} + 8\hat{k}, 2\hat{i} + 3\hat{j} + 4\hat{k}$  and  $2\hat{i} + 5\hat{j} + 7\hat{k}$  are the position vectors of the vertices  $A, B$  and  $C$  respectively of triangle  $ABC$ . The position vector of the point where the bisector of angle  $A$  meets  $BC$  is  
a)  $\frac{1}{2}(6\hat{i} + 13\hat{j} + 18\hat{k})$  b)  $\frac{2}{3}(6\hat{i} + 12\hat{j} - 8\hat{k})$  c)  $\frac{1}{3}(-6\hat{i} - 8\hat{j} - 9\hat{k})$  d)  $\frac{2}{3}(-6\hat{i} - 12\hat{j} + 8\hat{k})$
282. If the vectors  $\vec{a} = 2\hat{i} + 3\hat{j} + 6\hat{k}$  are collinear and  $|\vec{b}| = 21$ , then  $\vec{b} =$   
a)  $\pm 3(2\hat{i} + 3\hat{j} + 6\hat{k})$  b)  $\pm(2\hat{i} + 3\hat{j} - 6\hat{k})$  c)  $\pm 21(2\hat{i} + 3\hat{j} + 6\hat{k})$  d)  $\pm 21(\hat{i} + \hat{j} + \hat{k})$
283. The value of  $[\vec{a} - \vec{b}, \vec{b} - \vec{c}, \vec{c} - \vec{a}]$ , where  $|\vec{a}| = 1, |\vec{b}| = 5, |\vec{c}| = 3$ , is  
a) 0 b) 1 c) 6 d) None of these
284. The distance between the line  $\vec{r} = 2\hat{i} - 2\hat{j} + 3\hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$  and the plane  $\vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 5$  is  
a)  $\frac{10}{9}$  b)  $\frac{3}{10}$  c)  $\frac{10}{3\sqrt{3}}$  d)  $\frac{10}{9}$
285. In a parallelogram  $ABCD$ ,  $|\vec{AB}| = a, |\vec{AD}| = b$  and  $|\vec{AC}| = c$ . Then,  $\vec{DB} \cdot \vec{AB}$  has the value  
a)  $\frac{3a^2 + b^2 - c^2}{2}$  b)  $\frac{a^2 + 3b^2 - c^2}{2}$  c)  $\frac{a^2 - b^2 + 3c^2}{2}$  d)  $\frac{a^2 + 3b^2 + c^2}{2}$
286. If  $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$  and  $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$ , then the angle between the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  is  
a)  $60^\circ$  b)  $90^\circ$  c)  $45^\circ$  d)  $55^\circ$
287. If  $\vec{a} = \hat{i} + \hat{j} - \hat{k}, \vec{b} = -\hat{i} + 2\hat{j} + 2\hat{k}$  and  $\vec{c} = -\hat{i} + 2\hat{j} - \hat{k}$ , then a unit vector normal to the vectors  $\vec{a} + \vec{b}$  and  $\vec{b} - \vec{c}$  is  
a)  $\hat{i}$  b)  $\hat{j}$  c)  $\hat{k}$  d) None of these
288. If  $\vec{a}, \vec{b}, \vec{c}$  and three vectors such that  $\vec{a} = \vec{b} + \vec{c}$  and the angle between  $\vec{b}$  and  $\vec{c}$  is  $\frac{\pi}{2}$  then  
a)  $a^2 = b^2 + c^2$  b)  $b^2 = c^2 + a^2$  c)  $c^2 = a^2 + b^2$  d)  $2a^2 - b^2 = c^2$
289. If the position vector of  $A$  with respect to  $O$  is  $3\hat{i} - 2\hat{j} + 4\hat{k}$  and  $\vec{AB} = 3\hat{i} - \hat{j} + \hat{k}$ . Then the position vector of  $B$  with respect to  $O$  is  
a)  $-\hat{j} + 3\hat{k}$  b)  $6\hat{i} - 3\hat{j} + 5\hat{k}$  c)  $\hat{j} - 3\hat{k}$  d)  $\hat{i} - 3\hat{j} + 5\hat{k}$
290. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 3\hat{j} + 5\hat{k}$  and  $\vec{c} = 7\hat{i} + 9\hat{j} + 11\hat{k}$ , then the area of the parallelogram having diagonals  $\vec{a} + \vec{b}$  and  $\vec{b} + \vec{c}$  is  
a)  $4\sqrt{6}$  b)  $\frac{1}{2}\sqrt{21}$  c)  $\frac{\sqrt{6}}{2}$  d)  $\sqrt{6}$
291. The angle between the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$ , where  $\vec{a} = (1, 1, 4)$  and  $\vec{b} = (1, -1, 4)$  is  
a)  $90^\circ$  b)  $45^\circ$  c)  $30^\circ$  d)  $15^\circ$
292. Area of rhombus is ....., where diagonals are  $\vec{a} = 2\hat{i} - 3\hat{j} + 5\hat{k}$  and  $\vec{b} = -\hat{i} + \hat{j} + \hat{k}$   
a)  $\sqrt{21.5}$  b)  $\sqrt{31.5}$  c)  $\sqrt{28.5}$  d)  $\sqrt{38.5}$

293. If the vectors  $\hat{i} - 2x\hat{j} - 3y\hat{k}$  and  $\hat{i} + 3x\hat{j} + 2y\hat{k}$  are orthogonal to each other, then the locus of the point  $(x, y)$  is  
a) A circle                      b) An ellipse                      c) A parabola                      d) A straight line
294. If the position vectors of the vertices of a triangle are  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$ , then the triangle is  
a) Equilateral                      b) Isosceles                      c) Right angled isosceles                      d) Right angled
295. The two variable vectors  $3x\hat{i} + y\hat{j} - 3\hat{k}$  and  $x\hat{i} - 4y\hat{j} + 4\hat{k}$  are orthogonal to each other, then the locus of  $(x, y)$  is  
a) Hyperbola                      b) Circle                      c) Straight line                      d) Ellipse
296. If  $|\vec{a}| = |\vec{b}| = |\vec{a} + \vec{b}| = 1$ , then  $|\vec{a} - \vec{b}|$  is equal to  
a) 1                      b)  $\sqrt{2}$                       c)  $\sqrt{3}$                       d) None of these
297. The angle between the vectors  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $2\hat{i} - \hat{j} - \hat{k}$  is  
a)  $\pi/2$                       b)  $\pi/4$                       c)  $\pi/3$                       d) None of these
298. A unit vector coplanar with  $\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + 2\hat{j} + \hat{k}$  and perpendicular to  $\hat{i} + \hat{j} + \hat{k}$  is  
a)  $\left(\frac{\hat{j} - \hat{k}}{\sqrt{2}}\right)$                       b)  $\left(\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}\right)$                       c)  $\left(\frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}}\right)$                       d)  $\left(\frac{\hat{i} + 2\hat{j} + \hat{k}}{\sqrt{6}}\right)$
299. The length of the longer diagonal of the parallelogram constructed on  $5\vec{a} + 2\vec{b}$  and  $\vec{a} - 3\vec{b}$  if it is given that  $|\vec{a}| = 2\sqrt{2}$ ,  $|\vec{b}| = 3$  and angle between  $\vec{a}$  and  $\vec{b}$  is  $\pi/4$ , is  
a) 15                      b)  $\sqrt{113}$                       c)  $\sqrt{593}$                       d)  $\sqrt{369}$
300. The position vector of the point where the line  $\vec{r} = \hat{i} - \hat{j} + \hat{k} + t(\hat{i} + \hat{j} - \hat{k})$  meets the plane  $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 5$  is  
a)  $5\hat{i} + \hat{j} - \hat{k}$                       b)  $5\hat{i} + 3\hat{j} - 3\hat{k}$                       c)  $2\hat{i} + \hat{j} + 2\hat{k}$                       d)  $5\hat{i} + \hat{j} + \hat{k}$
301. If  $\vec{a} + \vec{b} + \vec{c} = 0$ ,  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$ ,  $|\vec{c}| = 7$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is  
a)  $\pi/6$                       b)  $2\pi/3$                       c)  $5\pi/3$                       d)  $\pi/3$
302. If  $\vec{a}$  is perpendicular to  $\vec{b}$  and  $\vec{c}|\vec{a}| = 2$ ,  $|\vec{b}| = 3$ ,  $|\vec{c}| = 4$  and the angle between  $\vec{b}$  and  $\vec{c}$  is  $\frac{2\pi}{3}$ , then  $[\vec{a} \vec{b} \vec{c}]$  is equal to  
a)  $4\sqrt{3}$                       b)  $6\sqrt{3}$                       c)  $12\sqrt{3}$                       d)  $18\sqrt{3}$
303. The position vectors of the points  $A, B, C$  are  $(2\hat{i} + \hat{j} - \hat{k})$ ,  $(3\hat{i} - 2\hat{j} + \hat{k})$  and  $(\hat{i} + 4\hat{j} - 3\hat{k})$  respectively. These points  
a) Form an isosceles triangle                      b) Form a right angled triangle  
c) Are collinear                      d) Form a scalene triangle
304. If  $\vec{a} = 4\hat{i} + 6\hat{j}$  and  $\vec{b} = 3\hat{j} + 4\hat{k}$ , then the vector form of component of  $\vec{a}$  along  $\vec{b}$  is  
a)  $\frac{18}{10\sqrt{3}}(3\hat{j} + 4\hat{k})$                       b)  $\frac{18}{25}(3\hat{j} + 4\hat{k})$                       c)  $\frac{18}{\sqrt{3}}(3\hat{j} + 4\hat{k})$                       d)  $3\hat{j} + 4\hat{k}$
305. Two vectors  $\vec{a}$  and  $\vec{b}$  are non-collinear. If vectors  $\vec{c} = (x - 2)\vec{a} + \vec{b}$  and  $\vec{d} = (2x + 1)\vec{a} - \vec{b}$  are collinear, then  $x =$   
a)  $1/3$                       b)  $1/2$                       c) 1                      d) 0
306. Through the point  $P(\alpha, \beta, \gamma)$  a plane is drawn at right angles to  $OP$  to meet the coordinate axes are  $A, B, C$  respectively. If  $OP = p$  then equation of plane  $\overline{ABC}$  is  
a)  $\alpha x + \beta y + \gamma z = p$                       b)  $\frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = p$   
c)  $2\alpha x + 2\beta y + 2\gamma z = p^2$                       d)  $\alpha x + \beta y + \gamma z = p^2$
307. If  $ABCDEF$  is a regular hexagon with  $\overline{AB} = \vec{a}$  and  $\overline{BC} = \vec{b}$ , then  $\overline{CE}$  equals  
a)  $\vec{b} - \vec{a}$                       b)  $-\vec{b}$                       c)  $\vec{b} - 2\vec{a}$                       d) None of these
308. A unit vector perpendicular to both  $\hat{i} + \hat{j}$  and  $\hat{j} + \hat{k}$ , is  
a)  $\hat{i} - \hat{j} + \hat{k}$                       b)  $\hat{i} + \hat{j} + \hat{k}$                       c)  $\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$                       d)  $\frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$

309. Let  $ABCD$  be the parallelogram whose sides  $AB$  and  $AD$  are represented by the vectors  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$  respectively. Then, if  $\vec{a}$  is a unit vector parallel to  $\overrightarrow{AC}$ , then  $\vec{a}$  equal to
- a)  $\frac{1}{3}(3\hat{i} - 6\hat{j} - 2\hat{k})$       b)  $\frac{1}{3}(3\hat{i} + 6\hat{j} + 2\hat{k})$       c)  $\frac{1}{7}(3\hat{i} - 6\hat{j} - 3\hat{k})$       d)  $\frac{1}{7}(3\hat{i} + 6\hat{j} - 2\hat{k})$
310. The value of  $b$  such that the scalar product of the vector  $\hat{i} + \hat{j} + \hat{k}$  with the unit vector parallel to the sum of the vectors  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $b\hat{i} + 2\hat{j} + 3\hat{k}$  is one, is
- a)  $-2$       b)  $-1$       c)  $0$       d)  $1$
311. If  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar vectors and  $x\vec{a} + y\vec{b} + z\vec{c} = 0$ , then
- a) At least of one of  $x, y, z$  is zero  
b)  $x, y, z$  are necessarily zero  
c) None of them are zero  
d) None of these
312. The ratio in which  $\hat{i} + 2\hat{j} + 3\hat{k}$  divides the join of  $-2\hat{i} + 3\hat{j} + 5\hat{k}$  and  $7\hat{i} - \hat{k}$ , is
- a)  $1 : 2$       b)  $2 : 3$       c)  $3 : 4$       d)  $1 : 4$
313. For any three vectors  $\vec{a}, \vec{b}, \vec{c}$  the expression  $(\vec{a} - \vec{b}) \cdot \{(\vec{b} - \vec{c}) \times (\vec{c} - \vec{a})\}$  equals
- a)  $[\vec{a}\vec{b}\vec{c}]$       b)  $2[\vec{a}\vec{b}\vec{c}]$       c)  $[\vec{a}\vec{b}\vec{c}]^2$       d) None of these
314. The point of intersection of the lines  $\vec{r} = 7\hat{i} + 10\hat{j} + 3\hat{k} + s(2\hat{i} + 3\hat{j} + 4\hat{k})$  and  $\vec{r} = 3\hat{i} + 5\hat{j} + 7\hat{k} + t(\hat{i} + 2\hat{j} + 3\hat{k})$  is
- a)  $\hat{i} + \hat{j} - \hat{k}$       b)  $2\hat{i} - \hat{j} + 4\hat{k}$       c)  $\hat{i} - \hat{j} + \hat{k}$       d)  $\hat{i} + \hat{j} + \hat{k}$
315. Let  $\vec{p}$  and  $\vec{q}$  be the position vectors of  $P$  and  $Q$  respectively, with respect to  $O$  and  $|\vec{p}| = p, |\vec{q}| = q$ . The points  $R$  and  $S$  divide  $PQ$  internally and externally in the ratio  $2 : 3$  respectively. If  $OR$  and  $OS$  are perpendicular, then
- a)  $9p^2 = 4q^2$       b)  $4p^2 = 9q^2$       c)  $9p = 4q$       d)  $4p = 9q$
316. If  $\vec{a} = \hat{i} + \hat{j}$  and  $\vec{b} = 2\hat{i} - \hat{k}$  are two vectors, then the point of intersection of two lines  $\vec{r} \times \vec{a} = \vec{b} \times \vec{a}$  and  $\vec{r} \times \vec{b} = \vec{a} \times \vec{b}$  is
- a)  $\hat{i} + \hat{j} - \hat{k}$       b)  $\hat{i} - \hat{j} + \hat{k}$       c)  $3\hat{i} + \hat{j} - \hat{k}$       d)  $3\hat{i} - \hat{j} + \hat{k}$
317. If  $\vec{A} \times (\vec{B} \times \vec{C}) = \vec{B} \times (\vec{C} \times \vec{A})$  and  $[\vec{A} \vec{B} \vec{C}] \neq 0$ , then  $\vec{A} \times (\vec{B} \times \vec{C})$  is equal to
- a)  $\vec{0}$       b)  $\vec{A} \times \vec{B}$       c)  $\vec{B} \times \vec{C}$       d)  $\vec{C} \times \vec{A}$
318. If  $\vec{a}$  and  $\vec{b}$  are two vectors, then the equality  $|\vec{a} + \vec{b}| = |\vec{a}| + |\vec{b}|$  holds
- a) Only if  $\vec{a} = \vec{b} = \vec{0}$   
b) For all  $\vec{a}, \vec{b}$   
c) Only if  $\vec{a} = \lambda\vec{b}, \lambda > 0$  or  $\vec{a} = \vec{b} = \vec{0}$   
d) None of these
319. Let  $\vec{a} = \hat{i} - \hat{k}, \vec{b} = x\hat{i} + \hat{j} + (1 - x)\hat{k}$  and  $\vec{c} = y\hat{i} + x\hat{j} + (1 + x - y)\hat{k}$ . Then  $[\vec{a}, \vec{b}, \vec{c}]$  depends on
- a) neither  $x$  nor  $y$       b) both  $x$  and  $y$       c) only  $x$       d) only  $y$
320. If the position vectors of three points  $A, B, C$  are respectively  $\hat{i} + \hat{j} + \hat{k}, 2\hat{i} + 3\hat{j} - 4\hat{k}$  and  $7\hat{i} + 4\hat{j} + 9\hat{k}$ , then the unit vector perpendicular to the plane of triangle  $ABC$  is
- a)  $31\hat{i} - 18\hat{j} - 9\hat{k}$       b)  $\frac{31\hat{i} - 38\hat{j} - 9\hat{k}}{\sqrt{2486}}$       c)  $\frac{31\hat{i} + 38\hat{j} + 9\hat{k}}{\sqrt{2486}}$       d) None of these
321. For any three vectors  $\vec{a}, \vec{b}$  and  $\vec{c}, (\vec{a} - \vec{b}) \cdot (\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})$  is equal to
- a)  $2\vec{a} \cdot (\vec{b} \times \vec{c})$       b)  $[\vec{a} \vec{b} \vec{c}]$       c)  $[\vec{a} \vec{b} \vec{c}]^2$       d)  $0$
322. If  $\vec{a}, \vec{b}, \vec{c}$  are unit coplanar vectors, then  $[2\vec{a} - \vec{b} \ 2\vec{b} - \vec{c} \ 2\vec{c} - \vec{a}]$  is equal to
- a)  $1$       b)  $0$       c)  $-\sqrt{3}$       d)  $\sqrt{3}$
323. If  $\vec{a}$  and  $\vec{b}$  are two unit vectors inclined to  $x$ -axis at angles  $30^\circ$  and  $120^\circ$ , then  $|\vec{a} + \vec{b}|$  equals

a)  $\sqrt{\frac{2}{3}}$

b)  $\sqrt{2}$

c)  $\sqrt{3}$

d) 2

324. If the vectors  $\hat{i} - 2x\hat{j} + 3y\hat{k}$  and  $\hat{i} + 2x\hat{j} - 3y\hat{k}$  perpendicular, then the locus of  $(x, y)$  is  
 a) A circle                      b) An ellipse                      c) A hyperbola                      d) None of these

325. Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be non-zero vectors such that  $(\vec{a} \times \vec{b}) \times \vec{c} = -\frac{1}{4}|\vec{b}||\vec{c}|\vec{a}$ . If  $\theta$  is the acute angle between vectors  $\vec{b}$  and  $\vec{c}$ , then the angle between  $\vec{a}$  and  $\vec{c}$  is equal to

a)  $\frac{2\pi}{3}$

b)  $\frac{\pi}{4}$

c)  $\frac{\pi}{3}$

d)  $\frac{\pi}{2}$

326. A vector perpendicular to both the vectors  $\hat{i} + \hat{j} + \hat{k}$  and  $\hat{i} + \hat{j}$  is  
 a)  $\hat{i} + \hat{j}$                       b)  $\hat{i} - \hat{j}$                       c)  $c(\hat{i} - \hat{j})$ ,  $c$  is a scalar                      d) None of these

327. If  $\vec{a}, \vec{b}, \vec{c}$  are non-collinear vectors such that  $\vec{a} + \vec{b}$  is parallel to  $\vec{c}$  and  $\vec{c} + \vec{a}$  is parallel to  $\vec{b}$ , then  
 a)  $\vec{a} + \vec{b} = \vec{c}$   
 b)  $\vec{a}, \vec{b}, \vec{c}$  taken in order from the sides of a triangle  
 c)  $\vec{b} + \vec{c} = \vec{a}$   
 d) None of these

328. A force of magnitude  $\sqrt{6}$  acting along the line joining the points  $A(2, -1, 1)$  and  $B(3, 1, 2)$  displaces a particle from  $A$  to  $B$ . The work done by the force is  
 a) 6                      b)  $6\sqrt{6}$                       c)  $\sqrt{6}$                       d) 12

329. A unit vector  $\vec{a}$  makes an angle  $\frac{\pi}{4}$  with  $z$ -axis, if  $\vec{a} + \hat{i} + \hat{j}$  is a unit vector, then  $\vec{a}$  is equal to  
 a)  $\frac{\hat{i}}{2} + \frac{\hat{j}}{2} + \frac{\hat{k}}{2}$                       b)  $\frac{\hat{i}}{2} + \frac{\hat{j}}{2} - \frac{\hat{k}}{\sqrt{2}}$                       c)  $-\frac{\hat{i}}{2} - \frac{\hat{j}}{2} + \frac{\hat{k}}{\sqrt{2}}$                       d)  $\frac{\hat{i}}{2} - \frac{\hat{j}}{2} - \frac{\hat{k}}{\sqrt{2}}$

330. If  $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 144$  and  $|\vec{a}| = 4$  then  $|\vec{b}|$  is equal to  
 a) 12                      b) 3                      c) 8                      d) 4

331. If  $\vec{a}$  is non-zero vector of modulus  $|\vec{a}|$  and  $m$  is a non-zero scalar, then  $m\vec{a}$  is a unit vector, if  
 a)  $m = \pm 1$                       b)  $m = |\vec{a}|$                       c)  $m = \frac{1}{|\vec{a}|}$                       d)  $m = \pm 2$

332. If the constant forces  $2\hat{i} - 5\hat{j} + 6\hat{k}$  and  $-\hat{i} + 2\hat{j} - \hat{k}$  act on a particle due to which it is displaced from a point  $A(4, -3, -2)$  to a point  $B(6, 1, -3)$ , then the work done by the forces is  
 a) 15 units                      b) -15 units                      c) 9 units                      d) -9 units

333. If  $P, Q, R$  are three points with respective position vectors  $\hat{i} + \hat{j}, \hat{i} - \hat{j}$  and  $a\hat{i} + b\hat{j} + c\hat{k}$ . The points  $P, Q, R$  are collinear, if  
 a)  $a = b = c = 1$                       b)  $a = b = c = 0$                       c)  $a = 1, b, c \in R$                       d)  $a = 1, c = 0, b \in R$

334. The projection of the vector  $\vec{a} = 4\hat{i} - 3\hat{j} + 2\hat{k}$  on the axis making equal acute angles with the coordinate axes is  
 a) 3                      b)  $\sqrt{3}$                       c)  $\frac{3}{\sqrt{3}}$                       d) None of these

335. The value of  $[2\hat{i} \ 3\hat{j} - 5\hat{k}]$  is equal to  
 a) -30                      b) -25                      c) 0                      d) 11

336.  $(\vec{a} \times \vec{b}) \times (\vec{a} \times \vec{c}) \cdot \vec{d}$  equals  
 a)  $[\vec{a}\vec{b}\vec{c}](\vec{b} \cdot \vec{d})$                       b)  $[\vec{a}\vec{b}\vec{c}](\vec{a} \cdot \vec{d})$                       c)  $[\vec{a}\vec{b}\vec{c}](\vec{c} \cdot \vec{d})$                       d) None of these

337. If the constant force  $2\hat{i} - 5\hat{j} + 6\hat{k}$  and  $-\hat{i} + 2\hat{j} - \hat{k}$  act on a particle due to which it is displaced from a point  $A(4, -3, -2)$  to a point  $B(6, 1, -3)$  then the work done by the force is  
 a) 10 units                      b) -10 units                      c) 9 units                      d) None of these

338. If forces of magnitudes 6 and 7 units acting in the directions  $\hat{i} - 2\hat{j} + 2\hat{k}$  and  $2\hat{i} - 3\hat{j} - 6\hat{k}$  respectively act on a particle which is displaced from the point  $P(2, -1, -3)$  to  $Q(5, -1, 1)$ , then the work done by the forces is  
a) 4 units                      b) -4 units                      c) 7 units                      d) -7 units
339.  $[\vec{b} \times \vec{c} \quad \vec{c} \times \vec{a} \quad \vec{a} \times \vec{b}]$  is equal to  
a)  $[\vec{a} \vec{b} \vec{c}]$                       b)  $2[\vec{a} \vec{b} \vec{c}]$                       c)  $[\vec{a} \vec{b} \vec{c}]^2$                       d)  $\vec{a} \times (\vec{b} \times \vec{c})$
340.  $ABCD$  is a quadrilateral,  $P, Q$  are the mid points of  $\overline{BC}$  and  $\overline{AD}$ , then  $\overline{AB} + \overline{DC}$  is equal to  
a)  $3\overline{QP}$                       b)  $\overline{QP}$                       c)  $4\overline{QP}$                       d)  $2\overline{QP}$
341. If  $D, E, F$  are respectively the mid-points of  $AB, AC$  and  $BC$  respectively in a  $\Delta ABC$ , then  $\overline{BE} + \overline{AF} =$   
a)  $\overline{DC}$                       b)  $\frac{1}{2}\overline{BF}$                       c)  $2\overline{BF}$                       d)  $\frac{3}{2}\overline{BF}$
342.  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular unit vectors, then  $|\vec{a} + \vec{b} + \vec{c}|$  is equal to  
a)  $\sqrt{3}$                       b) 3                      c) 1                      d) 0
343. Let  $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} + 3\hat{j} - \hat{k}$  and  $\vec{c} = d\hat{i} + \hat{j} + (2d - 1)\hat{k}$ . If  $\vec{c}$  is parallel to the plane of the vectors  $\vec{a}$  and  $\vec{b}$ , then  $11d =$   
a) 2                      b) 1                      c) -1                      d) 0
344. If  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar vectors and  $\vec{p}, \vec{q}, \vec{r}$ , are reciprocal vectors, then  $(l\vec{a} + m\vec{b} + n\vec{c}) \cdot (l\vec{p} + m\vec{q} + n\vec{r})$  is  
a)  $l + m + n$                       b)  $l^3 + m^3 + n^3$                       c)  $l^2 + m^2 + n^2$                       d) None of these
345. If  $\vec{a} \cdot \vec{b} \cdot \vec{c}$  are unit vectors, then  $|\vec{a} - \vec{b}|^2 + |\vec{b} - \vec{c}|^2 + |\vec{c} - \vec{a}|^2$  does not exceed  
a) 4                      b) 9                      c) 8                      d) 6
346. A constant force  $\vec{F} = 2\hat{i} - 3\hat{j} + 2\hat{k}$  is acting on a particle such that the particle is displaced from the point  $(1, 2, 3)$  to the point  $(3, 4, 5)$ . The work done by the force is  
a) 2                      b) 3                      c) 4                      d) 5
347. The value of  $a$ , for which the points  $A, B, C$  with position vectors  $2\hat{i} - \hat{j} + \hat{k}, \hat{i} - 3\hat{j} - 5\hat{k}$  and  $a\hat{i} - 3\hat{j} + \hat{k}$  respectively are the vertices of a right triangle with  $C = \frac{\pi}{2}$  are  
a) -2 and -1                      b) -2 and 1                      c) 2 and -1                      d) 2 and 1
348. If  $(\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$ , then  
a)  $\vec{b} \times (\vec{c} \times \vec{a}) = \vec{0}$                       b)  $\vec{a} \times (\vec{b} \times \vec{c}) = \vec{0}$                       c)  $\vec{c} \times \vec{a} = \vec{a} \times \vec{b}$                       d)  $\vec{c} \times \vec{b} = \vec{b} \times \vec{a}$
349. If  $\vec{a} + \vec{b} \neq \vec{0}$  and  $\vec{c}$  is a non-zero vector, then  $(\vec{a} + \vec{b}) \times \{\vec{c} - (\vec{a} + \vec{b})\}$  is equal to  
a)  $\vec{a} + \vec{b}$                       b)  $(\vec{a} + \vec{b}) \times \vec{c}$                       c)  $\lambda \vec{c}$ , where  $\lambda \neq 0$                       d)  $\lambda(\vec{a} \times \vec{b}), \lambda \neq 0$
350. If a force  $\vec{F} = 3\hat{i} + 2\hat{j} - 4\hat{k}$  is acting at the point  $P(1, -1, 2)$  then the magnitude of moment of  $\vec{F}$  about the point  $Q(2, -1, 3)$  is  
a)  $\sqrt{57}$                       b)  $\sqrt{39}$                       c) 12                      d) 17
351. If  $|\vec{a}| = |\vec{b}| = 1$  and  $|\vec{a} + \vec{b}| = \sqrt{3}$ , then the value of  $(3\vec{a} - 4\vec{b}) \cdot (2\vec{a} + 5\vec{b})$  is  
a) -21                      b)  $-\frac{21}{2}$                       c) 21                      d)  $\frac{21}{2}$
352. If  $\hat{a}, \hat{b}, \hat{c}$  are three unit vectors such that  $\hat{b}$  and  $\hat{c}$  are non-parallel and  $\hat{a} \times (\hat{b} \times \hat{c}) = \frac{1}{2}\hat{b}$ , then the angle between  $\hat{a}$  and  $\hat{c}$  is  
a)  $30^\circ$                       b)  $45^\circ$                       c)  $60^\circ$                       d)  $90^\circ$
353. If the vectors  $3\hat{i} + \lambda\hat{j} + \hat{k}$  and  $2\hat{i} - \hat{j} + 8\hat{k}$  are perpendicular, then  $\lambda$  is equal to  
a) -14                      b) 7                      c) 14                      d)  $1/7$
354. The equation of the plane perpendicular to the line  $\frac{x-1}{1} = \frac{y-2}{-1} = \frac{z+1}{2}$  and passing through the point  $(2, 3, 1)$  is

- a)  $\vec{r} \cdot (\hat{i} + \hat{j} + 2\hat{k}) = 1$     b)  $\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 1$     c)  $\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 7$     d)  $\vec{r} \cdot (\hat{i} + \hat{j} - 2\hat{k}) = 10$
355.  $(\vec{a} - \vec{b}) \cdot \{(\vec{b} - \vec{c}) \times (\vec{c} - \vec{a})\}$  is equal to  
a)  $2\vec{a} \cdot \vec{b} \times \vec{c}$     b)  $\vec{a} \cdot \vec{b} \times \vec{c}$     c) 0    d)  $\vec{a} \cdot \vec{b}$
356. If  $\hat{n}_1, \hat{n}_2$  are two unit vectors and  $\theta$  is the angle between them, then  $\cos \theta/2 =$   
a)  $\frac{1}{2}|\hat{n}_1 + \hat{n}_2|$     b)  $\frac{1}{2}|\hat{n}_1 - \hat{n}_2|$     c)  $\frac{1}{2}(\hat{n}_1 \cdot \hat{n}_2)$     d)  $\frac{|\hat{n}_1 \times \hat{n}_2|}{2|\hat{n}_1| |\hat{n}_2|}$
357. Let  $ABCD$  be the parallelogram whose sides  $AB$  and  $AD$  are represented by the vectors  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$  respectively. Then if  $\vec{a}$  is a unit vector parallel to  $\vec{AC}$ , then  $\vec{a}$  is equal to  
a)  $(3\hat{i} - 6\hat{j} - 2\hat{k})/3$     b)  $(3\hat{i} + 6\hat{j} + 2\hat{k})/3$     c)  $(3\hat{i} - 6\hat{j} - 3\hat{k})/7$     d)  $(3\hat{i} + 6\hat{j} - 2\hat{k})/7$
358. If the points with position vectors  $60\hat{i} + 3\hat{j}$ ,  $40\hat{i} - 8\hat{j}$  and  $a\hat{i} - 52\hat{j}$  are collinear, then  $a$  is equal to  
a)  $-40$     b)  $-20$     c)  $20$     d)  $40$
359. If  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar vectors such that  $\vec{a} + \vec{b} + \vec{c} = \alpha\vec{d}$  and  $\vec{b} + \vec{c} + \vec{d} = \beta\vec{a}$ , then  $\vec{a} + \vec{b} + \vec{c} + \vec{d}$  is equal to  
a)  $\vec{0}$     b)  $\alpha\vec{d}$     c)  $\beta\vec{b}$     d)  $(\alpha + \beta)\vec{c}$
360. The unit vector perpendicular to  $\hat{i} - \hat{j}$  and coplanar with  $\hat{i} + 2\hat{j}$  and  $\hat{i} + 3\hat{j}$  is  
a)  $\frac{2\hat{i} - 5\hat{j}}{\sqrt{29}}$     b)  $2\hat{i} + 5\hat{j}$     c)  $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j})$     d)  $\hat{i} + \hat{j}$
361. If  $\vec{r} \cdot \vec{a} = \vec{r} \cdot \vec{b} = \vec{r} \cdot \vec{c} = 0$  for some non-zero vector  $\vec{r}$ , then the value of  $[\vec{a}\vec{b}\vec{c}]$ , is  
a) 2    b) 3    c) 0    d) None of these
362. If the angle between  $\hat{i} + \hat{k}$  and  $\hat{i} + \hat{j} + a\hat{k}$  is  $\frac{\pi}{3}$ , then the value of  $a$  is  
a) 0 or 2    b)  $-4$  or 0    c) 0 or  $-2$     d) 2 or  $-2$
363. A vector which makes equal angles with the vectors  $\frac{1}{3}(\hat{i} - 2\hat{j} + 2\hat{k})$ ,  $\frac{1}{5}(-4\hat{i} - 3\hat{k})$ , and  $\hat{j}$ , is  
a)  $5\hat{i} + \hat{j} + 5\hat{k}$     b)  $-5\hat{i} + \hat{j} + 5\hat{k}$     c)  $-5\hat{i} + \hat{j} + 5\hat{k}$     d)  $5\hat{i} + \hat{j} - 5\hat{k}$
364. Which one of the following vectors is of magnitude 6 and perpendicular to both  $\vec{a} = 2\hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} - 2\hat{j} + 2\hat{k}$ ?  
a)  $2\hat{i} - \hat{j} - 2\hat{k}$     b)  $2(2\hat{i} - \hat{j} + 2\hat{k})$     c)  $3(2\hat{i} - \hat{j} - 2\hat{k})$     d)  $2(2\hat{i} - \hat{j} - 2\hat{k})$
365. In a right angled triangle  $ABC$ , the hypotenuse  $Ab = p$ , then  $\vec{AB} \cdot \vec{AC} + \vec{BC} \cdot \vec{BA} + \vec{CA} \cdot \vec{CB}$  is equal to  
a)  $2p^2$     b)  $\frac{p^2}{2}$     c)  $p^2$     d) None of these
366. Which one of the following is not correct?  
a) If  $\vec{p} \cdot \vec{a} = \vec{p} \cdot \vec{b} = \vec{p} \cdot \vec{c}$  for some non-zero vector  $\vec{p}$  then  $\vec{a}, \vec{b}, \vec{c}$  are coplanar  
b) The vectors  $\hat{i} + 3\hat{j}$ ,  $2\hat{i} + \hat{k}$  and  $\hat{j} + \hat{k}$  are coplanar  
c) The vector  $\vec{a} \times (\vec{b} \times \vec{c})$  is coplanar with  $\vec{a}$  and  $\vec{b}$   
d) If  $\vec{a}, \vec{b}$  are unit vectors and angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{3}$ , then  $|\vec{a} + \vec{b}| < 1$
367. The length of the shortest distance between the two lines  
 $\vec{r} = (-3\hat{i} + 6\hat{j}) + s(-4\hat{i} + 3\hat{j} + 2\hat{k})$  and  $\vec{r} = (-2\hat{i} + 7\hat{k}) + t(-4\hat{i} + \hat{j} + \hat{k})$  is  
a) 7 units    b) 13 units    c) 8 units    d) 9 units
368. A vector perpendicular to the plane containing the points  $A(1, -1, 2)$ ,  $B(2, 0, -1)$ ,  $C(0, 2, 1)$  is  
a)  $4\hat{i} + 8\hat{j} - 4\hat{k}$     b)  $8\hat{i} + 4\hat{j} + 4\hat{k}$     c)  $3\hat{i} + \hat{j} + 2\hat{k}$     d)  $\hat{i} + \hat{j} - \hat{k}$
369. If  $\vec{a}$  and  $\vec{b}$  are unit vectors such that  $[\vec{a}\vec{b}\vec{a} \times \vec{b}] = \frac{1}{4}$ , then angle between  $\vec{a}$  and  $\vec{b}$  is  
a)  $\frac{\pi}{3}$     b)  $\frac{\pi}{4}$     c)  $\frac{\pi}{6}$     d)  $\frac{\pi}{2}$
370. If  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$ , then a value of  $\lambda$  for which  $\vec{a} + \lambda\vec{b}$  is perpendicular to  $\vec{a} - \lambda\vec{b}$ , is  
a)  $\frac{9}{16}$     b)  $\frac{3}{4}$     c)  $\frac{3}{2}$     d)  $\frac{4}{3}$
371.  $(\vec{x} - \vec{y}) \times (\vec{x} + \vec{y}) = \dots$  where  $\vec{x}, \vec{y} \in R^3$

- a)  $2(\vec{x} \times \vec{y})$                       b)  $|\vec{x}|^2 - |\vec{y}|^2$                       c)  $\frac{1}{2}(\vec{x} \times \vec{y})$                       d) None of these
372. If the vectors  $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$ ,  $\vec{b} = 2\hat{i} + 4\hat{j} + \hat{k}$  and  $\vec{c} = \lambda\hat{i} + \hat{j} + \mu\hat{k}$  are mutually orthogonal, then  $(\lambda, \mu)$  is equal to  
a)  $(-3, 2)$                       b)  $(2, -3)$                       c)  $(-2, 3)$                       d)  $(3, -2)$
373. Given that  $\vec{a} = (1, 1, 1)$ ,  $\vec{c} = (0, 1, -1)$  and  $\vec{a} \cdot \vec{b} = 3$ . If  $\vec{a} \times \vec{b} = \vec{c}$ , then  $\vec{b} =$   
a)  $(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$                       b)  $(\frac{2}{3}, \frac{2}{3}, \frac{4}{3})$                       c)  $(\frac{5}{3}, \frac{2}{3}, \frac{2}{3})$                       d) None of these
374. If  $\hat{a}$ ,  $\hat{b}$  and  $\hat{c}$  are three unit vectors such that  $\hat{a} + \hat{b} + \hat{c}$  is also a unit vector and  $\theta_1, \theta_2$  and  $\theta_3$  are the angles between the vectors  $\hat{a}, \hat{b}; \hat{b}, \hat{c}$  and  $\hat{c}, \hat{a}$  respectively, then among  $\theta_1, \theta_2$  and  $\theta_3$   
a) All are acute angles  
b) All are right angles  
c) At least one is obtuse angle  
d) None of these
375. Given vectors  $\vec{x} = 3\hat{i} - 6\hat{j} - \hat{k}$ ,  $\vec{y} = \hat{i} + 4\hat{j} - 3\hat{k}$  and  $\vec{z} = 3\hat{i} + 4\hat{j} + 12\hat{k}$ , then the projection of  $\vec{x} \times \vec{y}$  on vector  $\vec{z}$  is  
a) 14                      b) -14                      c) 12                      d) 15
376. If the vectors  $\vec{a}$  and  $\vec{b}$  are mutually perpendicular, then  $\vec{a} \times \{ \vec{a} \times \{ \vec{a} \times (\vec{a} \times \vec{b}) \} \}$  is equal to  
a)  $|\vec{a}|^2 \vec{b}$                       b)  $|\vec{a}|^3 \vec{b}$                       c)  $|\vec{a}|^4 \vec{b}$                       d) None of these
377. Let  $G$  be the centroid of  $\Delta ABC$ . If  $\vec{AB} = \vec{a}$ ,  $\vec{AC} = \vec{b}$ , then the  $\vec{AG}$ , in terms of  $\vec{a}$  and  $\vec{b}$  is  
a)  $\frac{2}{3}(\vec{a} + \vec{b})$                       b)  $\frac{1}{6}(\vec{a} + \vec{b})$                       c)  $\frac{1}{3}(\vec{a} + \vec{b})$                       d)  $\frac{1}{2}(\vec{a} + \vec{b})$
378. The moment of the couple formed by the forces  $5\hat{i} + \hat{k}$  and  $-5\hat{i} - \hat{k}$  acting at the point  $(9, -1, 2)$  and  $(3, -2, 1)$  respectively is  
a)  $-\hat{i} + \hat{j} + 5\hat{k}$                       b)  $\hat{i} - \hat{j} - 5\hat{k}$                       c)  $2\hat{i} - 2\hat{j} - 10\hat{k}$                       d)  $-2\hat{i} + 2\hat{j} + 10\hat{k}$
379. The value of  $c$  so that for all real  $x$ , then vectors  $ocx \hat{i} - 6\hat{j} + 3\hat{k}$ ,  $x\hat{i} + 2\hat{j} + 2cx\hat{k}$  make an obtuse angle are  
a)  $c < 0$                       b)  $0 < c < \frac{4}{3}$                       c)  $-\frac{4}{3} < c < 0$                       d)  $c > 0$
380. If  $\theta$  be the angle between the vectors  $\vec{a} = 2\hat{i} + 2\hat{j} - \hat{k}$  and  $\vec{b} = 6\hat{i} - 3\hat{j} + 2\hat{k}$ , then  
a)  $\cos \theta = \frac{4}{21}$                       b)  $\cos \theta = \frac{3}{19}$                       c)  $\cos \theta = \frac{2}{19}$                       d)  $\cos \theta = \frac{5}{21}$
381. The vectors  $2\hat{i} + 3\hat{j} - 4\hat{k}$  and  $a\hat{i} + b\hat{j} + c\hat{k}$  are perpendicular when  
a)  $a = 2, b = 3, c = -4$                       b)  $a = 4, b = 4, c = 5$                       c)  $a = 4, b = 4, c = -2$                       d) None of these
382. If  $\vec{\alpha} = x(\vec{a} \times \vec{b}) + y(\vec{b} \times \vec{c}) + z(\vec{c} \times \vec{a})$  and  $[\vec{a} \vec{b} \vec{c}] = \frac{1}{8}$ , then  $x + y + z$  is equal to  
a)  $8 \vec{\alpha} \cdot (\vec{a} + \vec{b} + \vec{c})$                       b)  $\vec{\alpha} \cdot (\vec{a} + \vec{b} + \vec{c})$                       c)  $8 (\vec{a} + \vec{b} + \vec{c})$                       d) None of these
383. If vectors  $3\hat{i} + \hat{j} - 5\hat{k}$  and  $a\hat{i} + b\hat{j} - 15\hat{k}$  are collinear, then  
a)  $a = 3, b = 1$                       b)  $a = 9, b = 1$                       c)  $a = 3, b = 3$                       d)  $a = 9, b = 3$
384. Let  $\vec{a}$  and  $\vec{b}$  be two unit vectors such that angle between them is  $60^\circ$ . Then,  $|\vec{a} - \vec{b}|$  is equal to  
a)  $\sqrt{5}$                       b)  $\sqrt{3}$                       c) 0                      d) 1
385. The point collinear with  $(1, -2, -3)$  and  $(2, 0, 0)$  among the following is  
a)  $(0, 4, 6)$                       b)  $(0, -4, -5)$                       c)  $(0, -4, -6)$                       d)  $(0, -4, 6)$
386. If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then the vectors  $(\vec{a} + \vec{b}) \times (\vec{a} \times \vec{b})$  is parallel to the vector  
a)  $\vec{a} - \vec{b}$                       b)  $\vec{a} + \vec{b}$                       c)  $2\vec{a} - \vec{b}$                       d)  $2\vec{a} + \vec{b}$
387. If  $\theta$  is the angle between the lines  $AB$  and  $AC$  where  $A, B$  and  $C$  are the three points with coordinates  $(1, 2, -1), (2, 0, 3), (3, -1, 2)$  respectively, then  $\sqrt{462} \cos \theta$  is equal to  
a) 20                      b) 10                      c) 30                      d) 40
388. Let  $\vec{v} = 2\hat{i} + \hat{j} - \hat{k}$  and  $\vec{w} = \hat{i} + 3\hat{k}$ , If  $\vec{u}$  is a unit vector, then maximum value of the scalar triple product  $[\vec{u} \vec{v} \vec{w}]$  is

- a)  $-1$                                       b)  $\sqrt{10} + \sqrt{6}$                                       c)  $\sqrt{59}$                                       d)  $\sqrt{60}$
389. Each of the angle between vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  is equal to  $60^\circ$ . If  $|\vec{a}| = 4, |\vec{b}| = 2$  and  $|\vec{c}| = 6$ , then the modulus of  $\vec{a} + \vec{b} + \vec{c}$ , is  
a) 10                                      b) 15                                      c) 12                                      d) None of these
390. A force of magnitude 5 unit acting along the vector  $2\hat{i} - 2\hat{j} + \hat{k}$  displaces the point of applications from  $(1,2,3)$  to  $(5,3,7)$  then the work done is  
a)  $50/7$  unit                                      b)  $50/3$  unit                                      c)  $25/3$  unit                                      d)  $25/4$  unit
391. The equation of the plane passing through three non-collinear points  $\vec{a}, \vec{b}, \vec{c}$  is  
a)  $\vec{r} \cdot (\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}) = 0$                                       b)  $\vec{r} \cdot (\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}) = [\vec{a} \vec{b} \vec{c}]$   
c)  $\vec{r} \cdot (\vec{a} \times (\vec{b} \times \vec{c})) = [\vec{a} \vec{b} \vec{c}]$                                       d)  $\vec{r} \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$
392. If a vector  $\vec{r}$  of magnitude  $3\sqrt{6}$  is directed along the bisector of the angle between the vectors  $\vec{a} = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $\vec{b} = -2\hat{i} - \hat{j} + 2\hat{k}$ , then  $\vec{r} =$   
a)  $\hat{i} - 7\hat{j} + 2\hat{k}$                                       b)  $\hat{i} + 7\hat{j} - 2\hat{k}$                                       c)  $-\hat{i} + 7\hat{j} + 2\hat{k}$                                       d)  $\hat{i} - 7\hat{j} - 2\hat{k}$
393. If the point whose position vectors are  $2\hat{i} + \hat{j} + \hat{k}, 6\hat{i} - \hat{j} + 2\hat{k}$  and  $14\hat{i} - 5\hat{j} + p\hat{k}$  are collinear, then the value of  $p$  is  
a) 2                                      b) 4                                      c) 6                                      d) 8
394. Let  $\vec{a} \cdot \vec{b}$  and  $\vec{c}$  be non-zero vectors such that  
 $(\vec{a} \times \vec{b}) \times \vec{c} = \frac{1}{3} |\vec{b}| |\vec{c}| \vec{a}$   
If  $\theta$  is the acute angle between the vectors  $\vec{b}$  and  $\vec{c}$  then  $\sin \theta$  equals  
a)  $\frac{1}{3}$                                       b)  $\frac{\sqrt{2}}{3}$                                       c)  $\frac{2}{3}$                                       d)  $\frac{2\sqrt{2}}{3}$
395. Let  $ABC$  be a triangle, the position vectors of whose vertices are respectively  $7\hat{i} + 10\hat{k}, -\hat{i} + 6\hat{j} + 6\hat{k}$  and  $-4\hat{i} + 9\hat{j} + 6\hat{k}$  Then, the  $\Delta ABC$  is  
a) Isosceles                                      b) Equilateral  
c) Right angled isosceles                                      d) None of these
396. If  $C$  is the middle point of  $AB$  and  $P$  is any point outside  $AB$ , then  
a)  $P\vec{A} + P\vec{B} = P\vec{C}$                                       b)  $P\vec{A} + P\vec{B} = 2P\vec{C}$                                       c)  $P\vec{A} + P\vec{B} + P\vec{C} = \vec{0}$                                       d)  $P\vec{A} + P\vec{B} + 2P\vec{C} = \vec{0}$
397. If  $\vec{a}, \vec{b}$  are any two vvwctors, then  $(2\vec{a} + 3\vec{b}) \times (5\vec{a} + 7\vec{b}) + \vec{a} \times \vec{b}$  is equal to  
a)  $\vec{0}$                                       b) 0                                      c)  $\vec{a} \times \vec{b}$                                       d)  $\vec{b} \times \vec{a}$
398. The moment about the point  $M(-2, 4, -6)$  of the force represented in magnitude and position by  $AB$  where the points  $A$  and  $B$  have the coordinates  $(1, 2, -3)$  and  $(3, -4, 2)$  respectively is  
a)  $8\hat{i} - 9\hat{j} - 14\hat{k}$                                       b)  $2\hat{i} - 6\hat{j} + 5\hat{k}$                                       c)  $-3\hat{i} + 2\hat{j} - 3\hat{k}$                                       d)  $-5\hat{i} + 8\hat{j} - 8\hat{k}$
399. If the position vectors of  $A, B$  and  $C$  are respectively  $2\hat{i} - \hat{j} + \hat{k}, \hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  then  $\cos^2 A$  is equal to  
a) 0                                      b)  $\frac{6}{41}$                                       c)  $\frac{35}{41}$                                       d) 1
400. If  $\vec{r} \cdot \vec{a} = \vec{r} \cdot \vec{b} = \vec{r} \cdot \vec{c} = 0$  where  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar, then  
a)  $\vec{r} \perp \vec{c} \times \vec{a}$                                       b)  $\vec{r} \perp \vec{a} \times \vec{b}$                                       c)  $\vec{r} \perp \vec{b} \times \vec{c}$                                       d)  $\vec{r} = \vec{0}$
401. If  $\vec{a}, \vec{b}, \vec{c}$  be three non-coplanar vectors and  $\vec{p}, \vec{q}, \vec{r}$  constitute the corresponding reciprocal system of vectors then for any arbitrary vector  $\vec{\alpha}$   
a)  $\vec{\alpha} = (\vec{\alpha} \cdot \vec{a})\vec{a} + (\vec{\alpha} \cdot \vec{b})\vec{b} + (\vec{\alpha} \cdot \vec{c})\vec{c}$                                       b)  $\vec{\alpha} = (\vec{\alpha} \cdot \vec{p})\vec{p} + (\vec{\alpha} \cdot \vec{q})\vec{q} + (\vec{\alpha} \cdot \vec{r})\vec{r}$   
c)  $\vec{\alpha} = (\vec{\alpha} \cdot \vec{p})\vec{a} + (\vec{\alpha} \cdot \vec{q})\vec{b} + (\vec{\alpha} \cdot \vec{r})\vec{c}$                                       d) None of the above
402. The vector  $\vec{a} \times (\vec{b} \times \vec{c})$  is coplanar with the vectors  
a)  $\vec{b}, \vec{c}$                                       b)  $\vec{a}, \vec{b}$                                       c)  $\vec{a}, \vec{c}$                                       d)  $\vec{a}, \vec{b}, \vec{c}$
403. If  $\vec{b}$  is a unit vector, then  $(\vec{a} \cdot \vec{b})\vec{b} + \vec{b} \times (\vec{a} \times \vec{b})$  is

- a)  $|\vec{a}|^2\vec{b}$                       b)  $|\vec{a} \cdot \vec{b}|\vec{a}$                       c)  $\vec{a}$                       d)  $\vec{b}$
404. If  $\sum_{i=1}^n |\vec{a}_i| = \vec{0}$ , where  $|\vec{a}_i| = 1 \forall i$ , then the value of  $\sum_{1 \leq i < j \leq n} \vec{a}_i \cdot \vec{a}_j$  is  
a)  $n^2$                       b)  $-n^2$                       c)  $n$                       d)  $-\frac{n}{2}$
405. If the vector  $3\hat{i} - 2\hat{j} - 5\hat{k}$  is perpendicular to  $c\hat{k} - \hat{j} + 6\hat{i}$  then  $c$  is equal to  
a) 3                      b) 4                      c) 5                      d) 6
406. If  $\vec{a} \times \vec{b} = \vec{0}$  and  $\vec{a} \cdot \vec{b} = 0$ , then  
a)  $\vec{a} \perp \vec{b}$                       b)  $\vec{a} \parallel \vec{b}$                       c)  $\vec{a} = \vec{0}$  and  $\vec{b} = \vec{0}$                       d)  $\vec{a} = \vec{0}$  or  $\vec{b} = \vec{0}$
407. If  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$  are adjacent side of a parallelogram, then the lengths of its diagonals are  
a)  $7, \sqrt{69}$                       b)  $6, \sqrt{59}$                       c)  $5, \sqrt{65}$                       d)  $5, \sqrt{55}$
408. Let  $\vec{a}, \vec{b}, \vec{c}$  be unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ . Which of the following is correct?  
a)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} = \vec{0}$                       b)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$   
c)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{a} \times \vec{c} = \vec{0}$                       d)  $\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a}$  are mutually perpendicular
409. If  $G$  is the centre of a regular hexagon  $ABCDEF$ , then  $\vec{AB} + \vec{AC} + \vec{AD} + \vec{AE} + \vec{AF} =$   
a)  $3\vec{AG}$                       b)  $2\vec{AG}$                       c)  $6\vec{AG}$                       d)  $4\vec{AG}$
410. I. Two non-zero. Non-collinear vectors are linearly independent.  
II. Any three coplanar vectors are linearly dependent. Which of the above statements is /are true?  
a) Only I                      b) Only II                      c) Both I and II                      d) Neither I nor II
411. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are unit coplanar vectors, then  
 $[2\vec{a} - 3\vec{b} \ 7\vec{b} - 9\vec{c} \ 12\vec{c} - 23\vec{a}]$  is equal to  
a) 0                      b)  $1/2$                       c) 24                      d) 32
412.  $[\vec{a} + \vec{b} \ \vec{b} + \vec{c} \ \vec{c} + \vec{a}] = [\vec{a} \ \vec{b} \ \vec{c}]$ , then  
a)  $[\vec{a} \ \vec{b} \ \vec{c}] = 1$                       b)  $\vec{a}, \vec{b}, \vec{c}$  are coplanar  
c)  $[\vec{a} \ \vec{b} \ \vec{c}] = -1$                       d)  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular
413. If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = \sqrt{37}, |\vec{b}| = 3, |\vec{c}| = 4$ , then the angle between  $\vec{b}$  and  $\vec{c}$   
a)  $30^\circ$                       b)  $45^\circ$                       c)  $60^\circ$                       d)  $90^\circ$
414. A unit vector coplanar with  $\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + 2\hat{j} + \hat{k}$ , and perpendicular to  $\hat{i} + \hat{j} + \hat{k}$  is  
a)  $\left(\frac{\hat{j} - \hat{k}}{\sqrt{2}}\right)$                       b)  $\left(\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}\right)$                       c)  $\left(\frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}}\right)$                       d)  $\left(\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{6}}\right)$
415. The projection of the vector  $\hat{i} + \hat{j} + \hat{k}$  along the vector of  $\hat{j}$ , is  
a) 1                      b) 0                      c) 2                      d) -1
416. Volume of the parallelopiped having vertices at  $O \equiv (0,0,0), A \equiv (2, -2,4),$   
 $B \equiv (5, -4,4)$  and  $C \equiv (1, -2,4)$   
a) 5 cu units                      b) 10 cu units                      c) 15 cu units                      d) 20 cu units
417. The area of parallelogram constructed on the vectors  $\vec{a} = \vec{p} + 2\vec{q}$  and  $\vec{b} = 2\vec{p} + \vec{q}$ , where  $\vec{p}$  and  $\vec{q}$  are unit vectors forming an angle of  $30^\circ$  is  
a)  $3/2$                       b)  $5/2$                       c)  $7/2$                       d) None of these
418. If  $\vec{a}$  is a vector perpendicular to the vectors  $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{c} = -2\hat{i} + 4\hat{j} + \hat{k}$  and satisfies the condition  $\vec{a} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = -6$ , then  $\vec{a} =$   
a)  $5\hat{i} + \frac{7}{2}\hat{j} - 4\hat{k}$                       b)  $10\hat{i} + 7\hat{j} - 8\hat{k}$                       c)  $5\hat{i} - \frac{7}{2}\hat{j} + 4\hat{k}$                       d) None of these
419. The projection of  $\vec{a} = 3\hat{i} - \hat{j} + 5\hat{k}$  on  $\vec{b} = 2\hat{i} + 3\hat{j} + \hat{k}$  is  
a)  $\frac{8}{\sqrt{35}}$                       b)  $\frac{9}{\sqrt{39}}$                       c)  $\frac{8}{\sqrt{14}}$                       d)  $\sqrt{14}$
420. Let  $ABCDEF$  be a regular hexagon and  $\vec{AB} = \vec{a}, \vec{BC} = \vec{b}, \vec{CD} = \vec{c}$ , then  $\vec{AE}$  is equal to  
a)  $\vec{a} + \vec{b} + \vec{c}$                       b)  $\vec{b} + \vec{c}$                       c)  $\vec{a} + \vec{b}$                       d)  $\vec{a} + \vec{c}$







a)  $\frac{\pi}{4}$

b)  $\frac{\pi}{3}$

c)  $\frac{\pi}{2}$

d)  $\frac{3\pi}{2}$

466. Let  $\vec{a}, \vec{b}, \vec{c}$  three non-zero vectors such that no two of which are collinear and the vector  $\vec{a} + \vec{b}$  is collinear with  $\vec{c}$  and  $\vec{b} + \vec{c}$  is collinear with  $\vec{a}$ . Then,  $\vec{a} + \vec{b} + \vec{c} =$

a)  $\vec{a}$

b)  $\vec{b}$

c)  $\vec{c}$

d)  $\vec{0}$

467. The value of  $[\vec{a}\vec{b} + \vec{c}\vec{a} + \vec{b} + \vec{c}]$  is

a)  $[\vec{a}\vec{b}\vec{c}]$

b) 0

c)  $2[\vec{a}\vec{b}\vec{c}]$

d)  $\vec{a} \times (\vec{b} \times \vec{c})$

468. If the points with position vectors  $60\hat{i} + 3\hat{j}, 40\hat{i} - 8\hat{j}$  and  $a\hat{i} - 52\hat{j}$  are collinear, then  $a =$

a) -40

b) 40

c) 20

d) 30

469. Let  $\vec{a} = 2\hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 2\hat{j} - \hat{k}$  and a unit vector  $\vec{c}$  be coplanar. If  $\vec{c}$  is perpendicular to  $\vec{a}$  and  $\vec{c}$  is equal to

a)  $\pm \frac{1}{\sqrt{2}}(-\hat{j} + \hat{k})$

b)  $\pm \frac{1}{\sqrt{3}}(-\hat{i} - \hat{j} - \hat{k})$

c)  $\pm \frac{1}{\sqrt{5}}(\hat{i} - 2\hat{j})$

d)  $\pm \frac{1}{\sqrt{3}}(\hat{i} - \hat{j} - \hat{k})$

470. If the vectors  $\vec{a} = 2\hat{i} + \hat{j} + 4\hat{k}, \vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$  and  $\vec{c} = 2\hat{i} - 3\hat{j} - \lambda\hat{k}$  are coplanar, then the value of  $\lambda$  is equal to

a) 2

b) 1

c) 3

d) -1

471. The vectors

$$\vec{u} = (al + a_1l_1)\hat{i} + (am + a_1m_1)\hat{j} + (an + a_1n_1)\hat{k},$$

$$\vec{v} = (bl + b_1l_1)\hat{i} + (bm + b_1m_1)\hat{j} + (bn + b_1n_1)\hat{k},$$

$$\vec{w} = (cl + c_1l_1)\hat{i} + (cm + c_1m_1)\hat{j} + (cn + c_1n_1)\hat{k}$$

a) Form an equilateral triangle

b) Are coplanar

c) Are collinear

d) Are mutually perpendicular

472. If  $A, B, C, D$  are any four points in space, then  $|\vec{AB} \times \vec{CD} + \vec{BC} \times \vec{AD} + \vec{CA} \times \vec{BD}|$  is equal to

a)  $2\Delta$

b)  $4\Delta$

c)  $3\Delta$

d)  $5\Delta$

473. If  $\vec{a}$  lies in the plane of vectors  $\vec{b}$  and  $\vec{c}$ , then which of the following is correct?

a)  $[\vec{a}\vec{b}\vec{c}] = 0$

b)  $[\vec{a}\vec{b}\vec{c}] = 1$

c)  $[\vec{a}\vec{b}\vec{c}] = 3$

d)  $[\vec{b}\vec{c}\vec{a}] = 1$

474. What is the value of  $(\vec{d} + \vec{a}) \cdot [\vec{a} \times \{\vec{b} \times (\vec{c} \times \vec{d})\}]$ ?

a)  $(\vec{d} \cdot \vec{a}) \cdot [\vec{b} \vec{c} \vec{d}]$

b)  $(\vec{a} \cdot \vec{d}) \cdot [\vec{b} \vec{c} \vec{d}]$

c)  $(\vec{b} \cdot \vec{d}) \cdot [\vec{a} \vec{c} \vec{d}]$

d)  $(\vec{b} \cdot \vec{d}) \cdot [\vec{a} \vec{d} \vec{c}]$

475. A parallelogram is constructed on the vectors  $\vec{a} = 3\vec{\alpha} - \vec{\beta}, \vec{b} = \vec{\alpha} + 3\vec{\beta}$ . If  $|\vec{\alpha}| = |\vec{\beta}| = 2$  and the angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{3}$ , then the angle of a diagonal of the parallelogram are

a)  $4\sqrt{5}, 4\sqrt{3}$

b)  $4\sqrt{3}, 4\sqrt{7}$

c)  $4\sqrt{7}, 4\sqrt{5}$

d) None of these

476. If the vectors  $\hat{i} - 2\hat{j} + 3\hat{k}, -2\hat{i} + 3\hat{j} - 4\hat{k}, \lambda\hat{i} - \hat{j} + 2\hat{k}$  are linearly dependent, then the value of  $\lambda$  is equal to

a) 0

b) 1

c) 2

d) 3

477. For any vector  $\vec{a}$ , the value of  $(\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2$  is equal to

a)  $4\vec{a}^2$

b)  $2\vec{a}^2$

c)  $\vec{a}^2$

d)  $3\vec{a}^2$

478. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 2\hat{i} - 4\hat{k}, \vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$  are coplanar, then the value of  $\lambda$  is

a)  $\frac{5}{2}$

b)  $\frac{3}{5}$

c)  $\frac{7}{3}$

d) None of these

479. If the position vectors of  $P$  and  $Q$  are  $\hat{i} + 3\hat{j} - 7\hat{k}$  and  $5\hat{i} - 2\hat{j} + 4\hat{k}$  then the cosine of the angle between  $\vec{PQ}$  and  $y$ -axis is

a)  $\frac{5}{\sqrt{162}}$

b)  $\frac{4}{\sqrt{162}}$

c)  $-\frac{5}{\sqrt{162}}$

d)  $\frac{11}{\sqrt{162}}$

480. The value of 'a' so that volume of parallelepiped formed by  $\hat{i} + a\hat{j} + \hat{k}, \hat{j} + a\hat{k}$  and  $a\hat{i} + \hat{k}$  becomes minimum, is

a) -3

b) 3

c)  $1/\sqrt{3}$

d)  $\sqrt{3}$

481. If  $C$  is the mid point of  $AB$  and  $P$  is any point outside  $AB$ , then  
a)  $\overrightarrow{PA} + \overrightarrow{PB} = \overrightarrow{PC}$       b)  $\overrightarrow{PA} + \overrightarrow{PB} + 2\overrightarrow{PC} = \vec{0}$       c)  $\overrightarrow{PA} + \overrightarrow{PB} - 2\overrightarrow{PC} = \vec{0}$       d)  $\overrightarrow{PA} + \overrightarrow{PB} + 2\overrightarrow{PC} = \vec{0}$
482. The vector equation of the line passing through the points  $(3,2,1)$  and  $(-2,1,3)$  is  
a)  $\vec{r} = 3\hat{i} + 2\hat{j} + \hat{k} + \lambda(-5\hat{i} - \hat{j} + 2\hat{k})$       b)  $\vec{r} = 3\hat{i} + 2\hat{j} + \hat{k} + \lambda(-5\hat{i} + \hat{j} + \hat{k})$   
c)  $\vec{r} = -2\hat{i} + \hat{j} + 3\hat{k} + \lambda(5\hat{i} + \hat{j} + 2\hat{k})$       d)  $\vec{r} = -2\hat{i} + \hat{j} + \hat{k} + \lambda(5\hat{i} + \hat{j} + 2\hat{k})$
483. The angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{5\pi}{6}$  and the projection of  $\vec{a}$  in the direction of  $\vec{b}$  is  $\frac{-6}{\sqrt{3}}$  then  $|\vec{a}|$  is equal to  
a) 6      b)  $\sqrt{3}/2$       c) 12      d) 4
484. When a right handed rectangular cartesian system  $OXYZ$  rotated about  $z$ -axis through  $\pi/4$  in the counter-clock-wise sense it is found that a vector  $\vec{r}$  has the components  $2\sqrt{2}, 3\sqrt{2}$  and 4. The components of  $\vec{a}$  in the  $OXYZ$  coordinate system are  
a) 5, -1, 4      b) 5, -1,  $4\sqrt{2}$       c) -1, -5,  $4\sqrt{2}$       d) None of these
485. If  $\vec{x} \cdot \vec{a} = \vec{x} \cdot \vec{b} = \vec{x} \cdot \vec{c} = 0$  where  $\vec{x}$  is a non-zero vector. Then,  $[\vec{a} \times \vec{b} \cdot \vec{b} \times \vec{c} \cdot \vec{c} \times \vec{a}]$  is equal to  
a)  $[\vec{x} \vec{a} \vec{b}]^2$       b)  $[\vec{x} \vec{b} \vec{c}]^2$       c)  $[\vec{x} \vec{c} \vec{a}]^2$       d) 0
486. If  $ABCDEF$  is regular hexagon, then  $\overrightarrow{AD} + \overrightarrow{EB} + \overrightarrow{FC}$  is equal to  
a) 0      b)  $2\overrightarrow{AB}$       c)  $3\overrightarrow{AB}$       d)  $4\overrightarrow{AB}$
487. The shortest distance between the straight lines through the points  
 $A_1 = (6,2,2)$  and  $A_2 = (-4,0,-1)$  in the directions of  $(1,-2,2)$  and  $(3,-2,-2)$  is  
a) 6      b) 8      c) 12      d) 9
488. A unit vector perpendicular to the plane of  $\vec{a} = 2\hat{i} - 6\hat{j} - 3\hat{k}$  and  $\vec{b} = 4\hat{i} + 3\hat{j} - \hat{k}$  is  
a)  $\frac{4\hat{i} + 3\hat{j} - \hat{k}}{\sqrt{26}}$       b)  $\frac{2\hat{i} - 6\hat{j} - 3\hat{k}}{7}$       c)  $\frac{3\hat{i} - 2\hat{j} + 6\hat{k}}{7}$       d)  $\frac{2\hat{i} - 3\hat{j} - 6\hat{k}}{7}$
489. If  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  are the position vectors of points  $A, B, C, D$  such that no three of them are collinear and  $\vec{a} + \vec{c} = \vec{b} + \vec{d}$ , then  $ABCD$  is a  
a) Rhombus      b) Rectangle      c) Square      d) Parallelogram
490. If  $D, E, F$  are respectively the mid point of  $AB, AC$  and  $BC$  in  $\Delta ABC$ , then  $\overrightarrow{BE} + \overrightarrow{AF}$  is equal to  
a)  $\overrightarrow{DC}$       b)  $\frac{1}{2}\overrightarrow{BF}$       c)  $2\overrightarrow{BF}$       d)  $\frac{3}{2}\overrightarrow{BF}$
491. Let  $\vec{a}$  and  $\vec{b}$  be two unit vectors such that angle between them is  $60^\circ$ . Then,  $|\vec{a} - \vec{b}|$  is equal to  
a)  $\sqrt{5}$       b)  $\sqrt{3}$       c) 0      d) 1
492. If  $2\vec{a} + 3\vec{b} + \vec{c} = \vec{0}$ , then  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$  is equal to  
a)  $6(\vec{b} \times \vec{c})$       b)  $3(\vec{b} \times \vec{c})$       c)  $2(\vec{b} \times \vec{c})$       d)  $\vec{0}$
493. If  $\vec{a}, \vec{b}, \vec{c}$  are the three vectors mutually perpendicular to each other and  
 $|\vec{a}| = 1, |\vec{b}| = 3$  and  $|\vec{c}| = 5$ , then  $[\vec{a} - 2\vec{b} \cdot \vec{b} - 3\vec{c} \cdot \vec{c} - 4\vec{a}]$  is equal to  
a) 0      b) -24      c) 3600      d) -215
494. If the area of the parallelogram with  $\vec{a}$  and  $\vec{b}$  as two adjacent side is 15 sq units, then the area of the parallelogram having  $3\vec{a} + 2\vec{b}$  and  $\vec{a} + 3\vec{b}$  as two adjacent sides in sq units is  
a) 120      b) 105      c) 75      d) 45
495. If  $(\vec{a} \times \vec{b}) + (\vec{a} \cdot \vec{b})^2 = 144$  and  $|\vec{a}| = 4$ , then  $|\vec{b}| =$   
a) 16      b) 8      c) 3      d) 12
496. If the vectors  $\vec{c}, \vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$  and  $\vec{b} = \hat{j}$  are such that  $\vec{a}, \vec{c}$  and  $\vec{b}$  form a right handed system, then  $\vec{c}$  is  
a)  $z\hat{i} - x\hat{k}$       b)  $\vec{0}$       c)  $y\hat{i}$       d)  $-z\hat{i} + x\hat{k}$
497. The vectors  $2\hat{i} - m\hat{j} + 3m\hat{k}$  and  $(1+m)\hat{i} - 2m\hat{j} + \hat{k}$  include an acute angle for  
a)  $m = -1/2$   
b)  $m \in [-2, -1/2]$   
c)  $m \in R$

- d)  $m \in (-\infty, -2) \cup (-1/2, \infty)$
498. If  $|\vec{a}| + 3, |\vec{a}| = 4, |\vec{c}| = 5$  and  $\vec{a}, \vec{b}, \vec{c}$  are such that each is perpendicular to the sum of other two, then  $|\vec{a} + \vec{b} + \vec{c}|$  is
- a)  $5\sqrt{2}$                       b)  $\frac{5}{\sqrt{2}}$                       c)  $10\sqrt{2}$                       d)  $10\sqrt{3}$
499. For any three vectors  $\vec{a}, \vec{b}, \vec{c}$ , the vector  $(\vec{b} \times \vec{c}) \times \vec{a}$  equals
- a)  $(\vec{a} \cdot \vec{b})\vec{c} - (\vec{b} \cdot \vec{c})\vec{a}$       b)  $(\vec{a} \cdot \vec{b})\vec{c} - (\vec{a} \cdot \vec{c})\vec{b}$       c)  $(\vec{b} \cdot \vec{a})\vec{c} - (\vec{c} \cdot \vec{a})\vec{b}$       d) None of these
500. The vector  $\cos \alpha \cos \beta \hat{i} + \cos \alpha \sin \beta \hat{j} + \sin \alpha \hat{k}$  is a
- a) Null vector                      b) Unit vector                      c) Constant vector                      d) None of these
501. Let  $\vec{u}, \vec{v}, \vec{w}$  be such that  $|\vec{u}| = 1, |\vec{v}| = 2, |\vec{w}| = 3$ . If the projection  $\vec{v}$  along  $\vec{u}$  is equal to that of  $\vec{w}$  along  $\vec{u}$  and  $\vec{v}, \vec{w}$  are perpendicular to each other, then  $|\vec{u} - \vec{v} + \vec{w}|$  are equals
- a) 2                      b)  $\sqrt{7}$                       c)  $\sqrt{14}$                       d) 14
502. Let  $\vec{a}, \vec{b}, \vec{c}$  be the position vectors of the vertices  $A, B, C$  respectively of  $\Delta ABC$ . The vector area of  $\Delta ABC$  is
- a)  $\frac{1}{2}\{\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b})\}$       b)  $\frac{1}{2}\{\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}\}$   
c)  $\frac{1}{2}\{\vec{a} + \vec{b} + \vec{c}\}$                       d)  $\frac{1}{2}(\vec{b} \cdot \vec{c})\vec{a} + (\vec{c} \cdot \vec{a})\vec{b} + (\vec{a} \cdot \vec{b})\vec{c}$
503. If  $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \times \vec{c}$ , where  $\vec{a}, \vec{b}$  and  $\vec{c}$  are any three vectors such that  $\vec{a} \cdot \vec{b} \neq 0, \vec{b} \cdot \vec{c} \neq 0$ , then  $\vec{a}$  and  $\vec{c}$  are
- a) inclined at angle of  $\frac{\pi}{6}$  between them                      b) Perpendicular  
c) Parallel                      d) inclined at an angle of  $\frac{\pi}{3}$  between them
504. A unit vector in the plane of  $\hat{i} + 2\hat{j} + \hat{k}$  and  $\hat{i} + \hat{j} + 2\hat{k}$  and perpendicular to  $2\hat{i} + \hat{j} + \hat{k}$  is
- a)  $\hat{j} - \hat{k}$                       b)  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$                       c)  $\frac{\hat{j} + \hat{k}}{\sqrt{2}}$                       d)  $\frac{\hat{j} - \hat{k}}{\sqrt{2}}$
505. The unit vectors  $\vec{a}$  and  $\vec{b}$  are perpendicular, and the unit vector  $\vec{c}$  is inclined at an angle  $\theta$  to both  $\vec{a}$  and  $\vec{b}$ . If  $\vec{c} = \alpha\vec{a} + \beta\vec{b} + \gamma(\vec{a} \times \vec{b})$ , then which one of the following is incorrect?
- a)  $\alpha \neq \beta$                       b)  $\gamma^2 = 1 - 2\alpha^2$                       c)  $\gamma^2 = -\cos 2\theta$                       d)  $\beta^2 = \frac{1 + \cos 2\theta}{2}$
506. A vector  $\vec{c}$  of magnitude  $5\sqrt{6}$  directed along the bisector of the angle between  $\vec{a} = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $\vec{b} = -2\hat{i} - \hat{j} + 2\hat{k}$ , is
- a)  $\pm \frac{5}{3}(2\hat{i} + 7\hat{j} + \hat{k})$                       b)  $\pm \frac{3}{5}(\hat{i} + 7\hat{j} + 2\hat{k})$                       c)  $\pm \frac{5}{3}(\hat{i} - 2\hat{j} + 7\hat{k})$                       d)  $\pm \frac{5}{3}(\hat{i} - 7\hat{j} + 2\hat{k})$
507. If the vectors  $\vec{a} = 2\hat{i} + 3\hat{j} + 6\hat{k}$  and  $\vec{b}$  are collinear and  $|\vec{b}| = 21$ , then  $\vec{b}$  is equal to
- a)  $\pm(2\hat{i} + 3\hat{j} + 6\hat{k})$                       b)  $\pm 3(2\hat{i} + 3\hat{j} + 6\hat{k})$                       c)  $(\hat{i} + \hat{j} + \hat{k})$                       d)  $\pm 21(2\hat{i} + 3\hat{j} + 6\hat{k})$
508. A parallelogram is constructed on the vectors  $\vec{a} = 3\vec{p} - \vec{q}, \vec{b} = \vec{p} + 3\vec{q}$  and also given that  $|\vec{p}| = |\vec{q}| = 2$ . If the vectors  $\vec{p}$  and  $\vec{q}$  are inclined at an angle  $\pi/3$ , then the ratio of the lengths of the diagonals of the parallelogram is
- a)  $\sqrt{6}:\sqrt{2}$                       b)  $\sqrt{3}:\sqrt{5}$                       c)  $\sqrt{7}:\sqrt{3}$                       d)  $\sqrt{6}:\sqrt{5}$
509. If  $[2\vec{a} + 4\vec{b}\vec{c}\vec{d}] = \lambda[\vec{a}\vec{c}\vec{d}] + \mu[\vec{b}\vec{c}\vec{d}]$ , then  $\lambda + \mu =$
- a) 6                      b) -6                      c) 10                      d) 8
510. If  $A, B$  and  $C$  are the vertices of a triangle whose position vectors are  $\vec{a}, \vec{b}$  and  $\vec{c}$  respectively  $G$  is the centroid of the  $\Delta ABC$ , then  $\vec{GA} + \vec{GB} + \vec{GC}$  is
- a)  $\vec{0}$                       b)  $\vec{a} + \vec{b} + \vec{c}$                       c)  $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$                       d)  $\frac{\vec{a} - \vec{b} - \vec{c}}{3}$
511.  $A, B$  have position vectors  $\vec{a}, \vec{b}$  relative to the origin  $O$  and  $X, Y$  divide  $\vec{AB}$  internally and externally respectively in the ratio  $2 : 1$ . Then,  $\vec{XY} =$

a)  $\frac{3}{2}(\vec{b} - \vec{a})$                       b)  $\frac{4}{3}(\vec{a} - \vec{b})$                       c)  $\frac{5}{6}(\vec{b} - \vec{a})$                       d)  $\frac{4}{3}(\vec{b} - \vec{a})$

512. If  $\vec{a} = (2, 1, -1)$ ,  $\vec{b} = (1, -1, 0)$ ,  $\vec{c} = (5 - 1, 1)$ , then unit vector parallel to  $\vec{a} + \vec{b} - \vec{c}$  but in opposite direction is

a)  $\frac{1}{3}(2\hat{i} - \hat{j} + 2\hat{k})$                       b)  $\frac{1}{2}(2\hat{i} - \hat{j} + 2\hat{k})$                       c)  $\frac{1}{3}(2\hat{i} - \hat{j} - 2\hat{k})$                       d) None of these

513. The number of vectors of unit length perpendicular to the two vectors

$\vec{a} = (1, 1, 0)$  and  $\vec{b} = (0, 1, 1)$  is

a) One                      b) Two                      c) Three                      d) Infinite

514. A vector which is a linear combination of the vectors  $3\hat{i} + 4\hat{j} + 5\hat{k}$  and  $6\hat{i} - 7\hat{j} - 3\hat{k}$  and is perpendicular to the vector  $\hat{i} + \hat{j} - \hat{k}$  is

a)  $3\hat{i} - 11\hat{j} - 8\hat{k}$                       b)  $-3\hat{i} + 11\hat{j} + 87\hat{k}$                       c)  $-9\hat{i} + 3\hat{j} - 2\hat{k}$                       d)  $9\hat{i} - 3\hat{j} + 2\hat{k}$

515. If  $\vec{x}$  and  $\vec{y}$  are unit vectors and  $\vec{x} \cdot \vec{y} = 0$ , then

a)  $|\vec{x} + \vec{y}| = 1$                       b)  $|\vec{x} + \vec{y}| = \sqrt{3}$                       c)  $|\vec{x} + \vec{y}| = 2$                       d)  $|\vec{x} + \vec{y}| = \sqrt{2}$

516. If the volume of a parallelepiped with  $\vec{a} \times \vec{b}$ ,  $\vec{b} \times \vec{c}$ ,  $\vec{c} \times \vec{a}$  as coterminous edges is 9 cu units, then the volume of the parallelepiped with

$(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c})$ ,  $(\vec{b} \times \vec{c}) \times (\vec{c} \times \vec{a})$ ,  $(\vec{c} \times \vec{a}) \times (\vec{a} \times \vec{b})$  as coterminous edges is

a) 9 cu units                      b) 729 cu units                      c) 81 cu units                      d) 27 cu units

517. The non-zero vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are related by  $\vec{a} = 8\vec{b}$  and  $\vec{c} = -7\vec{b}$ . Then, the angle between  $\vec{a}$  and  $\vec{c}$  is

a)  $\pi$                       b) 0                      c)  $\frac{\pi}{4}$                       d)  $\frac{\pi}{2}$

518. For any three non-zero vectors  $\vec{r}_1, \vec{r}_2$  and  $\vec{r}_3$ ,  $\begin{vmatrix} \vec{r}_1 \cdot \vec{r}_1 & \vec{r}_1 \cdot \vec{r}_2 & \vec{r}_1 \cdot \vec{r}_3 \\ \vec{r}_2 \cdot \vec{r}_1 & \vec{r}_2 \cdot \vec{r}_2 & \vec{r}_2 \cdot \vec{r}_3 \\ \vec{r}_3 \cdot \vec{r}_1 & \vec{r}_3 \cdot \vec{r}_2 & \vec{r}_3 \cdot \vec{r}_3 \end{vmatrix} = 0$ , Then, which of the following is false?

- a) All the three vectors are parallel to one and the same plane                      b) All the three vectors are linearly dependent  
c) This system of equation has a non-trivial solution                      d) All the three vectors are perpendicular to each other

519. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j}$ ,  $\vec{c} = \hat{i}$  and  $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda\vec{a} + \mu\vec{b}$ , then  $\lambda + \mu$  is equal to

a) 0                      b) 1                      c) 2                      d) 3

520. Let  $\vec{a}, \vec{b}, \vec{c}$  be three vector such that  $\vec{a} \neq \vec{0}$  and  $\vec{a} \times \vec{b} = 2\vec{a} \times \vec{c}$ ,  $|\vec{a}| = |\vec{c}| = 1$ ,  $|\vec{b}| = 4$  and  $|\vec{b} \times \vec{c}| = \sqrt{15}$ . If  $\vec{b} - 2\vec{c} = \lambda\vec{a}$ , then  $\lambda$  is equal to

a) 1                      b)  $\pm 4$                       c) 3                      d) -2

521. If  $\vec{r} \cdot \vec{a} = 0$ ,  $\vec{r} \cdot \vec{b} = 0$  and  $\vec{r} \cdot \vec{c} = 0$  for some non-zero vector  $\vec{r}$ . Then, the value of  $[\vec{a} \ \vec{b} \ \vec{c}]$  is

a) 0                      b)  $\frac{1}{2}$                       c) 1                      d) 2

522. If  $\vec{a}, \vec{b}, \vec{c}$  are any three mutually perpendicular vectors of equal magnitude  $a$ , then  $|\vec{a} + \vec{b} + \vec{c}|$  is equal to

a)  $a$                       b)  $\sqrt{2} a$                       c)  $\sqrt{3} a$                       d)  $2a$

523. A unit vector perpendicular to both the vectors  $\hat{i} + \hat{j}$  and  $\hat{j} + \hat{k}$  is

a)  $\frac{-\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$                       b)  $\frac{-\hat{i} + \hat{j} - \hat{k}}{3}$                       c)  $\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$                       d)  $\frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$

524. Let,  $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$ ,  $\vec{c} = \hat{i} + \hat{j} - \hat{k}$ . A vector coplanar to  $\vec{a}$  and  $\vec{b}$  has a projection along  $\vec{c}$  of magnitude  $\frac{1}{\sqrt{3}}$ , then the vector is

a)  $4\hat{i} - \hat{j} + 4\hat{k}$                       b)  $4\hat{i} + \hat{j} - 4\hat{k}$                       c)  $2\hat{i} + \hat{j} + \hat{k}$                       d) None of these

525. Let  $\vec{u}$  and  $\vec{v}$  are unit vectors such that  $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$  and  $\vec{w} \times \vec{u} = \vec{v}$ , then the value of  $[\vec{u} \ \vec{v} \ \vec{w}]$  is

a) 1                      b) -1                      c) 0                      d) None of these

526. The position vectors of the points  $A, B, C$  are  $2\hat{i} + \hat{j} - \hat{k}$ ,  $3\hat{i} - 2\hat{j} + \hat{k}$  and  $\hat{i} + 4\hat{j} - 3\hat{k}$  respectively. These points
- Form an isosceles triangle
  - Form a right triangle
  - Are collinear
  - Form a scalene triangle
527. If  $\vec{a} = \hat{i} - \hat{j} - \hat{k}$  and  $\vec{b} = \lambda\hat{i} - 3\hat{j} + \hat{k}$  and the orthogonal projection of  $\vec{b}$  on  $\vec{a}$  is  $\frac{4}{3}(\hat{i} - \hat{j} - \hat{k})$  then  $\lambda$  is equal to
- 0
  - 2
  - 12
  - 1
528. If three points  $A, B$  and  $C$  have position vectors  $(1, x, 3)$ ,  $(3, 4, 7)$  and  $(y, -2, -5)$  respectively and, if they are collinear, then  $(x, y)$  is equal to
- $(2, -3)$
  - $(-2, 3)$
  - $(2, 3)$
  - $(-2, -3)$
529.  $\vec{OA}$  and  $\vec{OB}$  are two vectors of magnitude 5 and 6 respectively. If  $\angle BOA = 60^\circ$ , then  $\vec{OA} \cdot \vec{OB}$  is equal to
- 0
  - 15
  - 15
  - $15\sqrt{3}$
530. If  $\vec{a}$  and  $\vec{b}$  are two unit vectors inclined at an angle  $\theta$  such that  $\vec{a} + \vec{b}$  is a unit vector, then  $\theta$  is equal to
- $\frac{\pi}{3}$
  - $\frac{\pi}{4}$
  - $\frac{\pi}{2}$
  - $\frac{2\pi}{3}$
531.  $\vec{AB} \times \vec{AC} = 2\hat{i} - 4\hat{j} + 4\hat{k}$ , then the area of  $\Delta ABC$  is
- 3 sq units
  - 4 sq units
  - 16 sq units
  - 9 sq units
532. If the vectors  $\vec{c}, \vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$  and  $\vec{b} = \hat{j}$  are such that  $\vec{a}, \vec{c}$  and  $\vec{b}$  form a right handed system, then  $\vec{c}$  is
- $z\hat{i} - x\hat{k}$
  - $\vec{0}$
  - $y\hat{j}$
  - $-z\hat{i} - x\hat{k}$
533. Let  $\vec{a}, \vec{b}, \vec{c}$  be the vectors such that  $\vec{a} \neq \vec{0}$  and  $\vec{a} \times \vec{b} = 2\vec{a} \times \vec{c}$ ,  $|\vec{a}| = |\vec{c}| = 1$ ,  $|\vec{b}| = 4$  and  $|\vec{b} \times \vec{c}| = \sqrt{15}$ . If  $\vec{b} - 2\vec{c} = \lambda \vec{a}$ , then  $\lambda$  is equal to
- 1
  - 4
  - 3
  - 2
534. The position vectors of  $P$  and  $Q$  are respectively  $\vec{a}$  and  $\vec{b}$ . If  $R$  is a point on  $\vec{PQ}$  such that  $\vec{PR} = 5\vec{PQ}$ , then the position vector of  $R$ , is
- $5\vec{b} - 4\vec{a}$
  - $5\vec{b} + 4\vec{a}$
  - $4\vec{a} - 5\vec{b}$
  - $4\vec{b} + 5\vec{a}$
535. The vector  $\vec{c}$  is perpendicular to the vectors  $\vec{a} = (2, -3, 1)$ ,  $\vec{b} = (1, -2, 3)$  and satisfies the condition  $\vec{c} \cdot (\hat{i} + 2\hat{j} - 7\hat{k})$ . Then,  $\vec{c} =$
- $7\hat{i} + 5\hat{j} + \hat{k}$
  - $-7\hat{i} - 5\hat{j} - \hat{k}$
  - $\hat{i} + \hat{j} - \hat{k}$
  - None of these
536. If  $ABCD$  is a quadrilateral, then  $\vec{BA} + \vec{BC} + \vec{CD} + \vec{DA} =$
- $2\vec{BA}$
  - $2\vec{AB}$
  - $2\vec{AC}$
  - $2\vec{BC}$
537. The vector equation of the sphere whose centre is the point  $(1, 0, 1)$  and radius is 4, is
- $|\vec{r} - (\hat{i} + \hat{k})| = 4$
  - $|\vec{r} + (\hat{i} + \hat{k})| = 4^2$
  - $|\vec{r} \cdot (\hat{i} + \hat{k})| = 4$
  - $|\vec{r} \cdot (\hat{i} + \hat{k})| = 4^2$
538. If three concurrent edges of a parallelepiped of volume  $V$  represent vectors  $\vec{a}, \vec{b}, \vec{c}$  then the volume of the parallelepiped whose three concurrent edges are the three concurrent diagonals of the three faces of the given parallelepiped, is
- $V$
  - $2V$
  - $3V$
  - None of these
539. A unit vector in  $xy$ -plane makes an angle of  $45^\circ$  with the vector  $\hat{i} + \hat{j}$  and an angle of  $60^\circ$  with the vector  $3\hat{i} - 4\hat{j}$  is
- $\hat{i}$
  - $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$
  - $\frac{\hat{i} - \hat{j}}{\sqrt{2}}$
  - None of these
540. The equation  $\vec{r}^2 - 2\vec{r} \cdot \vec{c} + h = 0$ ,  $|\vec{c}| > \sqrt{h}$ , represent
- Circle
  - Ellipse
  - Cone
  - Sphere
541. The points with position vectors  $10\hat{i} + 3\hat{j}$ ,  $12\hat{i} - 5\hat{j}$  and  $a\hat{i} + 11\hat{j}$  are collinear if the value of  $a$  is
- 8
  - 4
  - 8
  - 12





## MATHS ( QUESTION BANK )

### 10.VECTOR ALGEBRA

#### : ANSWER KEY :

1)	d	2)	d	3)	c	4)	b	141)	d	142)	c	143)	d	144)	c
5)	c	6)	d	7)	d	8)	a	145)	a	146)	a	147)	d	148)	b
9)	c	10)	c	11)	c	12)	a	149)	a	150)	d	151)	b	152)	b
13)	b	14)	b	15)	c	16)	b	153)	a	154)	b	155)	b	156)	c
17)	b	18)	a	19)	c	20)	d	157)	a	158)	d	159)	b	160)	c
21)	a	22)	c	23)	d	24)	a	161)	c	162)	a	163)	c	164)	b
25)	d	26)	b	27)	b	28)	a	165)	a	166)	d	167)	d	168)	a
29)	d	30)	d	31)	c	32)	a	169)	b	170)	d	171)	c	172)	b
33)	a	34)	d	35)	a	36)	a	173)	d	174)	c	175)	d	176)	c
37)	b	38)	d	39)	d	40)	a	177)	a	178)	b	179)	c	180)	b
41)	c	42)	d	43)	a	44)	c	181)	c	182)	c	183)	c	184)	b
45)	d	46)	b	47)	a	48)	b	185)	a	186)	a	187)	b	188)	a
49)	b	50)	d	51)	a	52)	c	189)	d	190)	a	191)	a	192)	b
53)	b	54)	a	55)	c	56)	a	193)	b	194)	c	195)	b	196)	d
57)	b	58)	c	59)	d	60)	c	197)	a	198)	a	199)	a	200)	b
61)	b	62)	c	63)	a	64)	a	201)	a	202)	c	203)	c	204)	b
65)	d	66)	c	67)	a	68)	b	205)	a	206)	d	207)	b	208)	a
69)	b	70)	c	71)	d	72)	d	209)	d	210)	d	211)	c	212)	d
73)	a	74)	b	75)	d	76)	a	213)	b	214)	c	215)	b	216)	a
77)	b	78)	d	79)	b	80)	b	217)	d	218)	d	219)	a	220)	d
81)	b	82)	b	83)	c	84)	a	221)	b	222)	a	223)	c	224)	a
85)	b	86)	c	87)	a	88)	c	225)	d	226)	b	227)	c	228)	c
89)	a	90)	d	91)	a	92)	d	229)	b	230)	a	231)	a	232)	c
93)	c	94)	a	95)	d	96)	b	233)	b	234)	d	235)	a	236)	c
97)	d	98)	d	99)	a	100)	a	237)	c	238)	d	239)	d	240)	c
101)	d	102)	b	103)	b	104)	b	241)	a	242)	d	243)	b	244)	c
105)	b	106)	c	107)	c	108)	a	245)	b	246)	a	247)	d	248)	b
109)	a	110)	a	111)	c	112)	b	249)	b	250)	d	251)	c	252)	c
113)	d	114)	b	115)	b	116)	b	253)	b	254)	b	255)	d	256)	c
117)	b	118)	d	119)	d	120)	d	257)	a	258)	d	259)	d	260)	d
121)	c	122)	a	123)	a	124)	d	261)	c	262)	c	263)	b	264)	b
125)	a	126)	a	127)	c	128)	a	265)	d	266)	d	267)	b	268)	d
129)	a	130)	a	131)	a	132)	a	269)	c	270)	b	271)	d	272)	c
133)	c	134)	d	135)	a	136)	d	273)	b	274)	a	275)	b	276)	b
137)	a	138)	c	139)	c	140)	d	277)	b	278)	a	279)	d	280)	a

281) a	282) a	283) a	284) c	429) a	430) c	431) c	432) d
285) a	286) b	287) a	288) a	433) a	434) a	435) c	436) d
289) b	290) a	291) a	292) c	437) c	438) d	439) a	440) d
293) a	294) d	295) a	296) c	441) a	442) a	443) b	444) b
297) a	298) a	299) c	300) b	445) d	446) a	447) a	448) a
301) d	302) c	303) a	304) b	449) b	450) a	451) b	452) c
305) a	306) a	307) c	308) c	453) d	454) c	455) a	456) c
309) d	310) d	311) b	312) a	457) c	458) d	459) c	460) a
313) d	314) d	315) a	316) c	461) a	462) c	463) b	464) a
317) a	318) c	319) a	320) b	465) b	466) d	467) b	468) a
321) d	322) b	323) b	324) b	469) a	470) b	471) b	472) b
325) d	326) c	327) b	328) a	473) a	474) c	475) b	476) a
329) c	330) b	331) c	332) b	477) b	478) d	479) c	480) c
333) d	334) b	335) a	336) b	481) c	482) a	483) d	484) d
337) d	338) a	339) c	340) d	485) d	486) d	487) d	488) c
341) a	342) a	343) c	344) c	489) d	490) a	491) d	492) b
345) b	346) a	347) d	348) a	493) d	494) b	495) c	496) a
349) b	350) a	351) b	352) c	497) d	498) a	499) b	500) b
353) c	354) b	355) c	356) a	501) c	502) b	503) c	504) d
357) d	358) a	359) a	360) c	505) a	506) d	507) b	508) a
361) c	362) b	363) b	364) d	509) a	510) a	511) d	512) a
365) c	366) d	367) d	368) b	513) b	514) b	515) d	516) c
369) c	370) b	371) a	372) a	517) a	518) a	519) a	520) b
373) c	374) c	375) b	376) c	521) a	522) c	523) d	524) a
377) a	378) b	379) c	380) a	525) a	526) a	527) b	528) a
381) b	382) a	383) d	384) d	529) b	530) d	531) a	532) a
385) c	386) a	387) a	388) c	533) b	534) a	535) a	536) a
389) a	390) b	391) b	392) a	537) a	538) b	539) b	540) d
393) b	394) d	395) c	396) b	541) c	542) a	543) a	544) d
397) a	398) a	399) c	400) d	545) a	546) b	547) c	548) a
401) c	402) a	403) c	404) d	549) a	550) c	551) a	552) b
405) b	406) d	407) a	408) b	553) b	554) d	555) c	556) a
409) c	410) c	411) a	412) b	557) d	558) c	559) b	560) d
413) c	414) a	415) a	416) b	561) a	562) c	563) c	564) c
417) a	418) a	419) c	420) b	565) a	566) a	567) b	
421) d	422) c	423) a	424) a				
425) a	426) b	427) a	428) c				

## MATHS ( QUESTION BANK )

### 10. VECTOR ALGEBRA

#### : HINTS AND SOLUTIONS :

1 (d)

Let the unit vector in  $xy$ -plane be  $\vec{a} = x\hat{i} + y\hat{j}$ .

$$\therefore \cos 45^\circ = \frac{(x\hat{i} + y\hat{j})(\hat{i} + \hat{j})}{\sqrt{x^2 + y^2}\sqrt{1^2 + 1^2}}$$

$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{x + y}{\sqrt{2}\sqrt{x^2 + y^2}}$$

$$\Rightarrow 1 = \frac{x + y}{\sqrt{x^2 + y^2}}$$

$$\Rightarrow x + y = \sqrt{x^2 + y^2}$$

Since,  $\vec{a}$  is a unit vector.

$$\therefore |\vec{a}| = \sqrt{x^2 + y^2} = 1$$

$$\Rightarrow x + y = 1 \quad \dots(i)$$

$$\text{Again } \cos 60^\circ = \frac{(x\hat{i} + y\hat{j}) \cdot (3\hat{i} - 4\hat{j})}{\sqrt{x^2 + y^2}\sqrt{3^2 + 4^2}}$$

$$\Rightarrow \frac{1}{2} = \frac{3x - 4y}{1 \cdot 5} \Rightarrow \frac{5}{2} = 3x - 4y$$

$$5 = 6x - 8y \quad \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$x = \frac{13}{14}, y = \frac{1}{14}$$

$$\therefore \vec{a} = \frac{1}{14}(13\hat{i} + \hat{j})$$

No value in the given options satisfies the above relations.

Thus, option (d) is correct.

2 (d)

$$\text{Given, } |\vec{a} + \vec{b}| < 1$$

$$\Rightarrow \sqrt{1 + 1 + 2 \cos 2\alpha} < 1$$

$$\Rightarrow \sqrt{2(1 + \cos 2\alpha)} < 1$$

$$\Rightarrow \sqrt{4 \cos^2 \alpha} < 1$$

$$\Rightarrow |\cos \alpha| < \frac{1}{2}$$

$$\Rightarrow \frac{\pi}{3} < \alpha < \frac{2\pi}{3} \quad (\because 0 \leq \alpha \leq \pi)$$

3 (c)

Given equation can be rewritten as

$$\vec{r} = 3\hat{j} + (\hat{i} + 2\hat{k})s + (-2\hat{i} - \hat{j} + \hat{k})t$$

which is a plane passing through  $\vec{a} = 3\hat{j}$  and

parallel to the vectors  $\vec{b} = \hat{i} + 2\hat{k}$  and

$$\vec{c} = -2\hat{i} - \hat{j} + \hat{k}.$$

Therefore, it is perpendicular to the vector

$$\vec{n} = \vec{b} \times \vec{c} = 2\hat{i} - 5\hat{j} - \hat{k}$$

Hence, its vector equation is  $(\vec{r} - \vec{a}) \cdot \vec{n} = 0$

$$\Rightarrow \vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} - 5\hat{j} - \hat{k}) = 3\hat{j} \cdot (2\hat{i} - 5\hat{j} - \hat{k})$$

$$\Rightarrow 2x - 5y - z + 15 = 0$$

4 (b)

$$\because \vec{a} \cdot \vec{b} = 18 \text{ and } |\vec{b}| = 5$$

$\therefore$  Vector component of  $\vec{a}$  along  $\vec{b}$

$$= \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b} = \frac{18}{25} (3\hat{j} + 4\hat{k})$$

5 (c)

Given that,  $(\vec{F}) = 2\hat{i} + \hat{j} - \hat{k}$  and its position vector  $2\hat{i} - \hat{j}$ .

The position vector of a force about origin ( $\vec{r}$ ) =  $(2\hat{i} - \hat{j})$ .

$\therefore$  Moment of the force about origin

$$= \vec{r} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 0 \\ 2 & 1 & -1 \end{vmatrix} = \hat{i} + 2\hat{j} + 4\hat{k}$$

7 (d)

$$\text{Since, } \vec{a} \cdot (\vec{b} + \vec{c}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} = 0$$

$$\text{Similarly, } \vec{b} \cdot \vec{c} + \vec{b} \cdot \vec{a} = 0$$

$$\text{and } \vec{c} \cdot \vec{a} + \vec{c} \cdot \vec{b} = 0$$

$$\therefore \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c} = 0 \dots(i)$$

$$\text{Given, } |\vec{a} + \vec{b}| = 6$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = 36 \dots(ii)$$

$$\text{Similarly, } |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{b} \cdot \vec{c} = 64 \dots(iii)$$

$$\text{and } |\vec{c}|^2 + |\vec{a}|^2 + 2\vec{c} \cdot \vec{a} = 100 \dots(iv)$$

On adding Eqs. (ii),(iii) and (iv), we get

$$2|\vec{a}|^2 + 2|\vec{b}|^2 + 2|\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 200$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 = 100 \dots (v) \text{ [from Eqs. (i)]}$$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 100 \text{ [from Eqs. (i) and (v)]}$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = 10$$

8 (a)

It is given that  $|\vec{a}| = |\vec{b}| = |\vec{c}| = \lambda$  (say) and  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular vectors. Therefore,

$$|\vec{a} + \vec{b} + \vec{c}| = \sqrt{3} \lambda$$

Let  $\theta$  be the angle which  $\vec{a} + \vec{b} + \vec{c}$  makes with  $\vec{a}$ .

Then,

$$\cos \theta = \frac{\vec{a}(\vec{a} + \vec{b} + \vec{c})}{|\vec{a}||\vec{a} + \vec{b} + \vec{c}|} = \frac{|\vec{a}|^2}{|\vec{a}||\vec{a} + \vec{b} + \vec{c}|}$$

$$\Rightarrow \cos \theta = \frac{\lambda^2}{\lambda(\sqrt{3}\lambda)} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \cos^{-1}(1/\sqrt{3})$$

9 (c)

The resultant of forces  $3 \vec{OA}$  and  $5 \vec{OB}$  is  $8 \vec{OC}$ , where  $C$  divides  $AB$  in the ratio  $5 : 3$  i.e.  $3AC = 5CB$

10 (c)

The equation of a line passing through the centre  $(\hat{j} + 2\hat{k})$  and normal to the given plane is

$$\vec{r} = \hat{j} + 2\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k}) \dots (i)$$

This meets the plane at a point for which we must have

$$[(\hat{j} + 2\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 2\hat{k})] \cdot (\hat{i} + 2\hat{j} + 2\hat{k}) = 15$$

$$\Rightarrow 6 + \lambda(9) = 15 \Rightarrow \lambda = 1$$

$\therefore$  From Eq. (i),

$$\vec{r} = \hat{i} + 3\hat{j} + 4\hat{k}$$

$\therefore$  Coordinates of the centre of the circle are  $(1, 3, 4)$

12 (a)

$$\text{Let } \vec{a} = \hat{i} + 2\hat{j} - \hat{k}, \vec{b} = \hat{i} + \hat{j} + \hat{k}$$

$$\text{and } \vec{c} = \hat{i} - \hat{j} + \lambda\hat{k}$$

$$\text{Since, volume of tetrahedron} = \frac{1}{6} [\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow \frac{2}{3} = \frac{1}{6} \begin{vmatrix} 1 & 2 & -1 \\ 1 & 1 & 1 \\ 1 & -1 & \lambda \end{vmatrix}$$

$$\Rightarrow \frac{2}{3} = \frac{1}{6} [1(\lambda + 1) - 2(\lambda - 1) - 1(-1 - 1)]$$

$$\Rightarrow 4 = [-\lambda + 5]$$

$$\Rightarrow \lambda = 1$$

13 (b)

Given equation represents a plane.

15 (c)

$$\therefore \vec{\alpha} \times \vec{\beta} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & -1 \\ -1 & 2 & -4 \end{vmatrix} = -10\hat{i} + 9\hat{j} + 7\hat{k}$$

$$\therefore (\vec{\alpha} \times \vec{\beta}) \cdot (\vec{\alpha} \times \vec{\gamma}) = \begin{vmatrix} -10 & 9 & 7 \\ 2 & 3 & -1 \\ 1 & 1 & 1 \end{vmatrix}$$

$$= -10(3 + 1) - 9(2 + 1) + 7(2 - 3)$$

$$= -74$$

Alternate

$$(\vec{\alpha} \times \vec{\beta}) \cdot (\vec{\alpha} \times \vec{\gamma}) = \begin{vmatrix} \vec{\alpha} \cdot \vec{\alpha} & \vec{\alpha} \cdot \vec{\gamma} \\ \vec{\beta} \cdot \vec{\alpha} & \vec{\beta} \cdot \vec{\gamma} \end{vmatrix}$$

$$= \begin{vmatrix} 14 & 4 \\ 8 & -3 \end{vmatrix} = -42 - 32$$

$$= -74$$

16 (b)

Given planes are

$$\vec{r} \cdot (\hat{i} - 3\hat{j} + \hat{k}) = 1 \dots (i)$$

$$\text{and } \vec{r} \cdot (2\hat{i} + 5\hat{j} - 3\hat{k}) = 2 \dots (ii)$$

Now,


$$(\hat{i} - 3\hat{j} + \hat{k}) \times (2\hat{i} + 5\hat{j} - 3\hat{k}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 1 \\ 2 & 5 & -3 \end{vmatrix}$$

$$= 4\hat{i} + 5\hat{j} + 11\hat{k}$$

Hence, line of intersection of the planes is parallel to the vector  $4\hat{i} + 5\hat{j} + 11\hat{k}$ .

17 (b)

$$\text{Given, } \vec{AD} + \vec{EB} + \vec{FC} = \lambda \vec{ED}$$



$$\Rightarrow (\vec{AE} + \vec{ED}) + (\vec{ED} + \vec{DB}) + 2\vec{ED} = \lambda \vec{ED}$$

$$\Rightarrow 4\vec{ED} + (\vec{AE} + \vec{DB}) = \lambda \vec{ED}$$

$$\Rightarrow 4\vec{ED} = \lambda \vec{ED} \quad (\because \vec{AE} = -\vec{DB})$$

Alternate

$$\text{Now, } \vec{AD} + \vec{EB} + \vec{FC} = 2(\vec{OD} + \vec{EO} + \vec{ED})$$

$$= 2(\vec{ED} + \vec{ED}) = 4\vec{ED} \therefore \lambda = 4$$

18 (a)

$$\text{Let } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\text{and } \vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

$$\text{Now, } [\vec{a} \vec{b} \hat{i}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ 1 & 0 & 0 \end{vmatrix}$$

$$= a_1(0 - 0) - a_2(0 - b_3) + a_3(0 - b_2)$$

$$= a_2b_3 - a_3b_2$$

$$\therefore 2[\vec{a} \vec{b} \hat{i}] \hat{i} = 2[a_2b_3 - a_3b_2] \hat{i}$$

$$\begin{aligned} \text{Similarly, } 2[\vec{a} \vec{b} \hat{j}] \hat{j} &= 2[a_3 b_1 - a_1 b_3] \hat{j} \\ \text{and } 2[\vec{a} \vec{b} \hat{k}] \hat{k} &= 2[a_1 b_2 - a_2 b_1] \hat{k} \\ \therefore 2[\vec{a} \vec{b} \hat{i}] \hat{i} + 2[\vec{a} \vec{b} \hat{j}] \hat{j} + 2[\vec{a} \vec{b} \hat{k}] \hat{k} + [\vec{a} \vec{b} \vec{a}] \\ &= 2[(a_2 b_3 - a_3 b_2) \hat{i} + (a_3 b_1 - a_1 b_3) \hat{j} \\ &\quad + (a_1 b_2 - a_2 b_1) \hat{k}] \\ &= (\vec{a} \times \vec{b}) \end{aligned}$$

19 (c)

$$\text{Given that, } \vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{a} \cdot \vec{b} = 1 \text{ and } \vec{a} \times \vec{b} = \hat{j} - \hat{k}$$

$$\begin{aligned} \text{As we know } \vec{a}(\vec{a} \times \vec{b}) &= (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} \\ \Rightarrow (\hat{i} + \hat{j} + \hat{k}) \times (\hat{j} - \hat{k}) &= (\hat{i} + \hat{j} + \hat{k}) - (\sqrt{3})^2 \vec{b} \\ \Rightarrow -2\hat{i} + \hat{j} + \hat{k} &= \hat{i} + \hat{j} + \hat{k} - 3\vec{b} \\ \Rightarrow 3\vec{b} &= 3\hat{i} \\ \Rightarrow \vec{b} &= \hat{i} \end{aligned}$$

20 (d)

Given,  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar vectors and  $\vec{p}, \vec{q}, \vec{r}$  defined by the relations

$$\vec{p} = \frac{\vec{b} \times \vec{c}}{[\vec{a} \vec{b} \vec{c}]}, \vec{q} = \frac{\vec{c} \times \vec{a}}{[\vec{a} \vec{b} \vec{c}]} \text{ and } \vec{r} = \frac{\vec{a} \times \vec{b}}{[\vec{a} \vec{b} \vec{c}]}$$

$$\therefore \vec{a} \cdot \vec{p} = \frac{\vec{a} \cdot \vec{b} \times \vec{c}}{[\vec{a} \vec{b} \vec{c}]} = \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{[\vec{a} \vec{b} \vec{c}]} = 1$$

$$\text{and } \vec{a} \cdot \vec{q} = \vec{a} \cdot \frac{\vec{c} \times \vec{a}}{[\vec{a} \vec{b} \vec{c}]} = \frac{\vec{a} \cdot (\vec{c} \times \vec{a})}{[\vec{a} \vec{b} \vec{c}]} = 0$$

$$\text{Similarly, } \vec{b} \cdot \vec{q} = \vec{c} \cdot \vec{r} = 1$$

$$\text{and } \vec{a} \cdot \vec{r} = \vec{b} \cdot \vec{p} = \vec{c} \cdot \vec{q} = \vec{c} \cdot \vec{p} = \vec{b} \cdot \vec{r} = 0$$

$$\begin{aligned} \therefore (\vec{a} + \vec{b}) \cdot \vec{p} + (\vec{b} + \vec{c}) \cdot \vec{q} + (\vec{c} + \vec{a}) \cdot \vec{r} \\ = \vec{a} \cdot \vec{p} + \vec{b} \cdot \vec{p} + \vec{b} \cdot \vec{q} + \vec{c} \cdot \vec{q} + \vec{c} \cdot \vec{r} + \vec{a} \cdot \vec{r} \\ = 1 + 1 + 1 = 3 \end{aligned}$$

21 (a)

$$\text{Given, } m_1 = |\vec{a}_1| = \sqrt{2^2 + (-1)^2 + (1)^2} = \sqrt{6}$$

$$m_2 = |\vec{a}_2| = \sqrt{3^2 + (-4)^2 + (-4)^2} = \sqrt{41}$$

$$m_3 = |\vec{a}_3| = \sqrt{1^2 + 1^2 + (-1)^2} = \sqrt{3}$$

$$\text{and } m_4 = |\vec{a}_4| = \sqrt{(-1)^2 + (3)^2 + (1)^2} = \sqrt{11}$$

$$\therefore m_3 < m_1 < m_4 < m_2$$

22 (c)

$$\text{Given, } [\lambda(\vec{a} + \vec{b}) \lambda^2 \vec{b} \lambda \vec{c}] = [\vec{a} \vec{b} + \vec{c} \vec{b}]$$

$$\begin{aligned} \Rightarrow \begin{vmatrix} \lambda(a_1 + b_1) & \lambda(a_2 + b_2) & \lambda(a_3 + b_3) \\ \lambda^2 b_1 & \lambda^2 b_2 & \lambda^2 b_3 \\ \lambda c_1 & \lambda c_2 & \lambda c_3 \end{vmatrix} \\ = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 + c_1 & b_2 + c_2 & b_3 + c_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \end{aligned}$$

$$\begin{aligned} \Rightarrow \lambda^4 \begin{vmatrix} a_1 + b_1 & a_2 + b_2 & a_3 + b_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 + c_1 & b_2 + c_2 & b_3 + c_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \end{aligned}$$

[applying  $R_1 \rightarrow R_1 - R_2$  in LHS and  $R_2 \rightarrow R_2 - R_3$  in RHS]

$$\Rightarrow \lambda^4 \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = - \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$\Rightarrow \lambda^4 = -1$$

Hence, no real value of  $\lambda$  exists.

23 (d)

Since the given points lie in a plane.

$$\therefore \begin{vmatrix} a & a & c \\ 1 & 0 & 1 \\ c & c & b \end{vmatrix} = 0$$

Applying  $C_1 \rightarrow C_1 - C_2$

$$\Rightarrow \begin{vmatrix} 0 & a & c \\ 1 & 0 & 1 \\ 0 & c & b \end{vmatrix} = 0$$

$$\Rightarrow -1(ab - c^2) = 0$$

$$\Rightarrow c^2 = ab$$

Hence,  $c$  is GM of  $a$  and  $b$ .

24 (a)

We have,

$$\vec{P} = A\vec{C} + \vec{B}D$$

$$\Rightarrow \vec{p} = A\vec{C} + B\vec{C} + \vec{C}D$$

$$\Rightarrow \vec{p} = A\vec{C} + \lambda \vec{A}D + \vec{C}D$$

$$\Rightarrow \vec{p} = \lambda \vec{A}D + (A\vec{C} + \vec{C}D)$$

$$\Rightarrow \vec{p} = \lambda \vec{A}D + \vec{A}D = (\lambda + 1)\vec{A}D$$

$$\therefore \vec{p} = \mu \vec{A}D \Rightarrow \mu = \lambda + 1$$

25

(d)

We have,

$$\vec{a} + \vec{b} + \vec{c} = 0$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 0$$

$$\Rightarrow (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$$

$$\Rightarrow 2(\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c}) = -\{|\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2\}$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{29}{2}$$

26 (b)

Since,  $(\vec{a} + \lambda \vec{b}) \cdot (\vec{a} - \lambda \vec{b}) = 0$

$$\Rightarrow (\vec{a})^2 - \lambda^2 (\vec{b})^2 = 0$$

$$\Rightarrow \lambda^2 \frac{(\vec{a})^2}{(\vec{b})^2} = \left(\frac{3}{4}\right)^2$$

$$\Rightarrow \lambda = \frac{3}{4}$$

27 (b)

$$\begin{aligned} \text{Let } \vec{a} &= a_1\hat{i} + a_2\hat{j} + a_3\hat{k} \\ \therefore \vec{u} &= \hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k}) \\ &= \hat{i} \times (-a_2\hat{k} + a_3\hat{j}) + \hat{j} \times (a_1\hat{k} - a_3\hat{i}) + \hat{k} \\ &\quad \times (-a_1\hat{j} + a_2\hat{i}) \\ &= a_2\hat{j} + a_3\hat{k} + a_1\hat{i} + a_3\hat{k} + a_1\hat{i} + a_2\hat{j} \\ &= 2\vec{a} \end{aligned}$$

29 (d)

$$\begin{aligned} \text{Since, } |\vec{a} + \vec{b}|^2 &= |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta \\ \Rightarrow (\sqrt{7})^2 &= (3\sqrt{3})^2 + 4^2 + 2(3\sqrt{3})(4)\cos\theta \\ \Rightarrow 7 &= 27 + 16 + 24\sqrt{3}\cos\theta \\ \Rightarrow \cos\theta &= -\sqrt{3}/2 \\ \Rightarrow \theta &= 150^\circ \end{aligned}$$

30 (d)

$$\begin{aligned} \therefore \vec{b} \times \vec{c} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -5 \\ 3 & 5 & -1 \end{vmatrix} = 23\hat{i} - 14\hat{j} - \hat{k} \\ \therefore \vec{a} \times (\vec{b} \times \vec{c}) &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & -1 \\ 23 & -14 & -1 \end{vmatrix} \\ &= -17\hat{i} - 21\hat{j} - 97\hat{k} \end{aligned}$$

31 (c)

$$\begin{aligned} \text{We have,} \\ |\vec{a}| = |\vec{b}| = |\vec{c}| = 1 \text{ and } \vec{a} \perp \vec{b} \perp \vec{c} \\ \Rightarrow \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0 \\ \therefore |\vec{a} + \vec{b} + \vec{c}|^2 &= (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) \\ &= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{a} \cdot \vec{b} + 2\vec{b} \cdot \vec{c} + 2\vec{c} \cdot \vec{a} = 3 \\ \Rightarrow |\vec{a} + \vec{b} + \vec{c}| &= \sqrt{3} \end{aligned}$$

32 (a)

$$\begin{aligned} \text{Let } \vec{a} &= x\hat{i} + y\hat{j} + z\hat{k} \\ \therefore (\vec{a} \cdot \hat{i})\hat{i} &= [(x\hat{i} + y\hat{j} + z\hat{k}) \cdot \hat{i}]\hat{i} = x\hat{i} \\ \text{Similarly, } (\vec{a} \cdot \hat{j})\hat{j} &= y\hat{j}, (\vec{a} \cdot \hat{k})\hat{k} = z\hat{k} \\ \therefore (\vec{a} \cdot \hat{i})\hat{i} + (\vec{a} \cdot \hat{j})\hat{j} + (\vec{a} \cdot \hat{k})\hat{k} &= x\hat{i} + y\hat{j} + z\hat{k} = \vec{a} \end{aligned}$$

33 (a)

$$\begin{aligned} \text{Let } \vec{a} &= x\hat{i} + y\hat{j} + z\hat{k} \\ \therefore \vec{a} \cdot \hat{i} &= (x\hat{i} + y\hat{j} + z\hat{k}) \cdot \hat{i} = x \\ \vec{a} \cdot (\hat{i} + \hat{j}) &= (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + \hat{j}) = x + y \\ \text{and } \vec{a} \cdot (\hat{i} + \hat{j} + \hat{k}) &= (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k}) = x + y + z \\ \therefore \text{Given that, } \vec{a} \cdot \hat{i} &= \vec{a} \cdot (\hat{i} + \hat{j}) = \vec{a} \cdot (\hat{i} + \hat{j} + \hat{k}) \\ \Rightarrow x &= x + y = x + y + z \\ \text{Take } x &= x + y \Rightarrow y = 0 \\ \text{and } x + y &= x + y + z \Rightarrow z = 0 \\ \Rightarrow x &\text{ has any real values.} \\ \text{Now, take } x &= 1 \therefore \vec{a} = \hat{i} \end{aligned}$$

34 (d)

$$\begin{aligned} \text{Let } \vec{c} &= \vec{a} + \lambda + \vec{b} = (1 + \lambda)\hat{i} + (1 - \lambda)\hat{j} + (\lambda - 1)\hat{k} \\ \text{Also, } \vec{c} \cdot \vec{a} &= 0 \\ \Rightarrow [(1 + \lambda)\hat{i} + (1 - \lambda)\hat{j} + (\lambda - 1)\hat{k}] \cdot [\hat{i} + \hat{j} - \hat{k}] &= 0 \\ \Rightarrow 1 + \lambda + 1 - \lambda - \lambda + 1 &= 0 \\ \Rightarrow \lambda &= 3 \\ \therefore \vec{c} &= 4\hat{i} - 2\hat{j} + 2\hat{k} \\ \Rightarrow \vec{c} &= \pm \frac{2\hat{i} - \hat{j} + \hat{k}}{\sqrt{6}} \end{aligned}$$

35 (a)

$$\begin{aligned} \text{The cartesian form of an equation of planes are} \\ x + 3y - z = 0 \text{ and } y + 2z = 0 \\ \text{The line of intersection of two planes is} \\ (x + 3y - z) + \lambda(y + 2z) = 0 \dots(i) \\ \text{Since, it is passing through } (-1, -1, -1) \\ \therefore (-1 - 3 + 1) + \lambda(-1 - 2) = 0 \\ \Rightarrow \lambda = -1 \\ \text{On putting the value of } \lambda \text{ in Eq. (i), we get} \\ x + 2y - 3z = 0 \end{aligned}$$

Hence, vector equation of plane is

$$\vec{r} \cdot (\hat{i} + 2\hat{j} - 3\hat{k}) = 0$$

39 (d)

$$\begin{aligned} \vec{a} \times [\vec{a} \times (\vec{a} \times \vec{b})] &= \vec{a} \times \{(\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}\} \\ &= 0 - (\vec{a} \cdot \vec{a})(\vec{a} \times \vec{b}) \\ &= (\vec{a} \cdot \vec{a})(\vec{b} \times \vec{a}) \end{aligned}$$

40 (a)

$$\begin{aligned} \begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{c} \cdot \vec{a} & \vec{c} \cdot \vec{b} & \vec{c} \cdot \vec{c} \end{vmatrix} \\ = \begin{vmatrix} a_1^2 + a_2^2 + a_3^2 & a_1b_1 + a_2b_2 + a_3b_3 & a_1c_1 \\ a_1b_1 + a_2b_2 + a_3b_3 & b_1^2 + b_2^2 + b_3^2 & b_1c_1 \\ a_1c_1 + a_2c_2 + a_3c_3 & b_1c_1 + b_2c_2 + b_3c_3 & c_1^2 \end{vmatrix} \\ = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ = [\vec{a} \vec{b} \vec{c}]^2 \end{aligned}$$

41 (c)

$$\begin{aligned} \text{Given, } \vec{a} \cdot \vec{b} &= 12 \\ \Rightarrow |\vec{a}||\vec{b}|\cos\theta &= 12 \\ \Rightarrow 10 \times 2 \times \cos\theta &= 12 \\ \Rightarrow \cos\theta &= \frac{3}{5} \\ \therefore \sin\theta &= \sqrt{1 - \cos^2\theta} = \sqrt{1 - \frac{9}{25}} = \frac{4}{5} \end{aligned}$$

Now,

$$|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}|\sin\theta = 10 \times 2 \times \frac{4}{5} = 16$$

42 (d)

We have,

$$|\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) \dots(i)$$

It is given that  $\vec{a} \perp (\vec{b} + \vec{c})$ ,  $\vec{b} \perp (\vec{c} + \vec{a})$  and  $\vec{c} \perp (\vec{a} + \vec{b})$

$$\therefore \vec{a} \cdot (\vec{b} + \vec{c}) = 0, \vec{b} \cdot (\vec{c} + \vec{a}) = 0 \text{ and } \vec{c} \cdot (\vec{a} + \vec{b}) = 0$$

$$\Rightarrow 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = 0$$

$$\therefore |\vec{a} + \vec{b} + \vec{c}|^2 = 16 + 16 + 25 + 0 \quad [\text{From (i)}]$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = \sqrt{57}$$

43 (a)

$$\text{Since, } \vec{a} + \vec{b} = \vec{c} \Rightarrow (\vec{a} + \vec{b})^2 = \vec{c}^2$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta = |\vec{c}|^2$$

$$\Rightarrow 2(1 + \cos\theta) = 1 \Rightarrow \cos\theta = -\frac{1}{2} \quad [\because |\vec{a}| = |\vec{b}| = |\vec{c}| = 1, \text{ given}]$$

$$\text{Now, } |\vec{a} + \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|\cos\theta$$

$$= 1 + 1 + 2 \cdot \frac{1}{2} = 3$$

$$\Rightarrow |\vec{a} + \vec{b}| = \sqrt{3}$$

44 (c)

We have,

$$\vec{a} \cdot \vec{b} \geq 0 \Rightarrow |\vec{a}||\vec{b}|\cos\theta \geq 0 \Rightarrow 0 \leq \theta \leq \frac{\pi}{2}$$

45 (d)

Since the vectors  $2\hat{i} + 3\hat{j}$  and  $5\hat{i} + 6\hat{j}$  have (1, 1) as initial point. Therefore, their terminal points are (3, 4) and (6, 7) respectively. The equation of the line joining these two points is  $x - y + 1 = 0$ . The terminal point of  $8\hat{i} + \lambda\hat{j}$  is  $(9, \lambda + 1)$ . Since the vectors terminate on the same straight line.

$$\text{Therefore, point } (9, (\lambda + 1)) \text{ lies on } x - y + 1 = 0 \Rightarrow 9 - (\lambda + 1) + 1 = 0 \Rightarrow \lambda = 9$$

47 (a)

$$\text{Let } \vec{A} = 2\hat{i} + 3\hat{j} - \hat{k} \dots(i)$$

$$\vec{B} = \hat{i} + 2\hat{j} - 3\hat{k} \dots(ii)$$

$$\vec{C} = 3\hat{i} + 4\hat{j} - 2\hat{k} \dots(iii)$$

$$\vec{D} = \hat{i} - \lambda\hat{j} + 6\hat{k} \dots(iv)$$

From Eq. (i) and (ii), we get

$$\vec{AB} = -\hat{i} - \hat{j} + 4\hat{k}$$

$\therefore$  From Eq. (i) and (iii), we get

$$\vec{AC} = \hat{i} + \hat{j} - \hat{k}$$

Similarly, from Eqs.(i) and (iv), we get

$$\vec{AD} = -\hat{i} - (\lambda - 3)\hat{j} + 7\hat{k}$$

Now, using condition of coplanarity

$$\begin{vmatrix} -1 & -1 & 4 \\ 1 & 1 & -1 \\ -1 & -(\lambda + 3) & 7 \end{vmatrix} = 0$$

Applying  $R_1 \rightarrow R_1 + R_2$ , we get

$$\begin{vmatrix} 0 & 0 & 3 \\ 1 & 1 & -1 \\ -1 & -(\lambda + 3) & 7 \end{vmatrix} = 0$$

$$\Rightarrow -\lambda - 2 = 0 \Rightarrow \lambda = -2$$

48 (b)

$$\text{Since, } |\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = |\vec{a}|^2 |\vec{b}|^2$$

$$\Rightarrow (10)^2 + |\vec{a} \cdot \vec{b}|^2 = (3)^2 \cdot (4)^2$$

$$\Rightarrow |\vec{a} \cdot \vec{b}|^2 = 44$$

49 (b)

Let  $\vec{a}, \vec{b}$  be the sides of the given parallelogram.

Then, its diagonals are  $\vec{a} + \vec{b}$  and  $\pm(\vec{a} - \vec{b})$

We have,

$$\vec{a} + \vec{b} = 3\hat{i} + \hat{j} - 2\hat{k} \text{ and } \vec{a} - \vec{b} = \pm(\hat{i} - 3\hat{j} + 4\hat{k})$$

$$\Rightarrow \vec{a} = 2\hat{i} - \hat{j} + \hat{k}, \vec{b} = \hat{i} + 2\hat{j} - 3\hat{k}$$

$$\text{or } \vec{a} = \hat{i} + 2\hat{j} - 3\hat{k} \text{ and } \vec{b} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\Rightarrow |\vec{a}| = \sqrt{6}, |\vec{b}| = \sqrt{14} \text{ or } |\vec{a}| = \sqrt{14}, |\vec{b}| = \sqrt{6}$$

50 (d)

$$\text{We have, } \vec{a} \times \vec{b} = \vec{c}$$

$$\Rightarrow \vec{c} \text{ is perpendicular to } \vec{a} \text{ and } \vec{b} \text{ and } \vec{b} \times \vec{c} = \vec{a}$$

$$\Rightarrow \vec{a} \text{ is perpendicular to } \vec{b} \text{ and } \vec{c}$$

$$\Rightarrow \vec{a}, \vec{b}, \vec{c} \text{ are mutually perpendicular.}$$

$$\text{Again } \vec{a} \times \vec{b} = \vec{c}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = |\vec{c}|$$

$$= |\vec{a}||\vec{b}| \cdot \sin 90^\circ = |\vec{c}|$$

$$\Rightarrow |\vec{a}||\vec{b}| = |\vec{c}| \dots(i)$$

$$\text{Also, } \vec{b} \times \vec{c} = \vec{a}$$

$$|\vec{b}||\vec{c}| \cdot \sin 90^\circ = |\vec{a}|$$

$$|\vec{b}||\vec{c}| = |\vec{a}| \dots(ii)$$

From Eqs. (i) and (ii), we get

$$|\vec{b}|^2 |\vec{c}| = |\vec{c}|$$

$$\therefore |\vec{b}|^2 = 1 \quad (\because |\vec{c}| \neq 0)$$

$$\Rightarrow |\vec{b}| = 1$$

$$\Rightarrow |\vec{a}| = |\vec{c}|$$

51 (a)

$$\vec{a} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & -1 \\ 1 & -1 & -1 \end{vmatrix}$$

$$= \hat{i}(-1 - 1) - \hat{j}(0 + 1) + \hat{k}(0 - 1)$$

$$= -2\hat{i} - \hat{j} + \hat{k}$$

Given,

$$\vec{a} \times \vec{b} + \vec{c} = \vec{0}$$

$$\begin{aligned} \Rightarrow \vec{a} \times (\vec{a} \times \vec{b}) + \vec{a} \times \vec{c} &= \vec{0} \\ \Rightarrow (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} &= -\vec{a} \times \vec{c} \\ \Rightarrow 3\vec{a} - 2\vec{b} &= -\vec{a} \times \vec{c} \\ \Rightarrow \vec{b} &= \frac{3\vec{a} + \vec{a} \times \vec{c}}{2} \\ \Rightarrow \vec{b} &= \frac{3\hat{j} - 3\hat{k} - 2\hat{i} - \hat{j} - \hat{k}}{2} \\ &= \frac{-2\hat{i} + 2\hat{j} - 4\hat{k}}{2} = -\hat{i} + \hat{j} - 2\hat{k} \end{aligned}$$

53 (b)

$$\begin{aligned} \text{Given } \vec{a} + \vec{b} &= -\vec{c} \\ \Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta &= |\vec{c}|^2 \\ \Rightarrow 9 + 25 + 2 \cdot 3 \cdot 5 \cos\theta &= 49 \\ \Rightarrow \cos\theta &= \frac{1}{2} \\ \Rightarrow \theta &= \frac{\pi}{3} \end{aligned}$$

54 (a)

$$\begin{aligned} 3\vec{p} + \vec{q} - 2\vec{r} &= 3(\hat{i} + \hat{j}) + (4\hat{k} - \hat{j}) - 2(\hat{i} + \hat{k}) \\ &= \hat{i} + 2\hat{j} + 2\hat{k} \\ \therefore \text{Unit vector in the direction of } 3\vec{p} + \vec{q} - 2\vec{r} \\ &= \frac{1}{3}(\hat{i} + 2\hat{j} + 2\hat{k}) \end{aligned}$$

55 (c)

$$\begin{aligned} \text{Solving the two equations for } \vec{X} \text{ and } \vec{Y}, \text{ we get} \\ \vec{X} &= \frac{1}{3}(\hat{i} + 3\hat{j}) \text{ and } \vec{Y} = \frac{1}{3}(\hat{i} - 3\hat{j}) \\ \therefore \cos\theta &= \frac{\vec{X} \cdot \vec{Y}}{|\vec{X}||\vec{Y}|} \Rightarrow \cos\theta = -\frac{4}{5} \end{aligned}$$

56 (a)

$$\begin{aligned} |\vec{p} + \vec{q}| &= 6 \\ \Rightarrow |\vec{p} + \vec{q}|^2 &= 36 \\ \Rightarrow p^2 + q^2 + 2\vec{p} \cdot \vec{q} &= 36 \\ \text{Similarly, } q^2 + r^2 + 2\vec{q} \cdot \vec{r} &= 48 \\ \text{and } r^2 + p^2 + 2\vec{r} \cdot \vec{p} &= 16 \\ \text{adding all, we get} \\ 2(p^2 + q^2 + r^2 + \vec{p} \cdot \vec{q} + \vec{q} \cdot \vec{r} + \vec{r} \cdot \vec{p}) \\ \Rightarrow 2(p^2 + q^2 + r^2) &= 100 \quad (\because \vec{p} \cdot \vec{q} + \vec{q} \cdot \vec{r} + \vec{r} \cdot \vec{p} = 0) \\ \Rightarrow p^2 + q^2 + r^2 &= 50 \\ \Rightarrow |\vec{p} + \vec{q} + \vec{r}|^2 &= 50 \\ \Rightarrow |\vec{p} + \vec{q} + \vec{r}| &= 5\sqrt{2} \end{aligned}$$

57 (b)

$$\begin{aligned} \text{In triangles } OAC \text{ and } OBD, \text{ we have} \\ \vec{OA} + \vec{OC} &= 2\vec{OM} \text{ and } \vec{OB} + \vec{OD} = 2\vec{OM} \\ \Rightarrow \vec{OA} + \vec{OB} + \vec{OC} + \vec{OD} &= 4\vec{OM} \end{aligned}$$

58 (c)

The work done is given by

$$\begin{aligned} W = \vec{F} \cdot \vec{d} &= (2\hat{i} - \hat{j} - \hat{k}) \cdot (3\hat{i} + 2\hat{j} - 5\hat{k}) \\ &= 9 \text{ units} \end{aligned}$$

59 (d)

$$\begin{aligned} [\hat{i} \hat{k} \hat{j}] + [\hat{k} \hat{j} \hat{i}] + [\hat{j} \hat{k} \hat{i}] \\ = [\hat{i} \hat{k} \hat{j}] + [\hat{i} \hat{k} \hat{j}] - [\hat{i} \hat{k} \hat{j}] \\ = [\hat{i} \hat{k} \hat{j}] = \hat{i} \cdot (\hat{k} \times \hat{j}) \\ = \hat{i} \cdot (-\hat{i}) = -1 \end{aligned}$$

60 (c)

$$\begin{aligned} \text{Given, } \vec{a} \cdot \vec{b} &= \vec{a} \cdot \vec{c} = 0 \\ \text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 &= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) \\ &= (1)^2 + (1)^2 + (1)^2 + 2\left(0 + |\vec{b}||\vec{c}|\cos\frac{\pi}{3} + 0\right) \\ &= 3 + 2 \times 1 \times 1 \times \frac{1}{2} = 4 \\ \Rightarrow |\vec{a} + \vec{b} + \vec{c}| &= \pm 2 \end{aligned}$$

61 (b)

$$\begin{aligned} \text{Let } \vec{AB} = \vec{a} &= 3\vec{\alpha} - \vec{\beta}, \vec{BC} = \vec{b} = \vec{\alpha} + 3\vec{\beta} \\ \text{Diagonal } \vec{AC} &= \vec{AB} + \vec{BC} = \vec{a} + \vec{b} \\ \Rightarrow |\vec{AC}| &= |\vec{a} + \vec{b}| \\ \Rightarrow |\vec{AC}| &= |4\vec{\alpha} + 2\vec{\beta}| \\ \Rightarrow |\vec{AC}|^2 &= 16\vec{\alpha}^2 + 4\vec{\beta}^2 + 16\vec{\alpha} \cdot \vec{\beta} \\ \Rightarrow |\vec{AC}|^2 &= 64 + 16 + 16|\vec{\alpha}||\vec{\beta}|\cos\frac{\pi}{3} \\ \Rightarrow |\vec{AC}|^2 &= 80 + 16 \times 4 \times \frac{1}{2} = 112 \\ \Rightarrow |\vec{AC}| &= 4\sqrt{7} \\ \text{Other diagonal is } |\vec{BD}| &= |\vec{a} - \vec{b}| \\ \Rightarrow |\vec{BD}|^2 &= |2\vec{\alpha} - 4\vec{\beta}|^2 \\ &= 4|\vec{\alpha}|^2 + 16|\vec{\beta}|^2 - 16|\vec{\alpha}||\vec{\beta}|\cos\frac{\pi}{3} \\ &= 64 + 16 - 16 \times 4 \times \frac{1}{2} = 48 \\ \Rightarrow |\vec{BD}| &= \sqrt{48} = 4\sqrt{3} \end{aligned}$$

62 (c)

$$\begin{aligned} \text{Given that, } |\vec{a}| &= |\vec{b}| \\ \text{Now, } (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) &= \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{a} - \vec{b} \cdot \vec{a} - \vec{b} \cdot \vec{b} \\ &= 0 \quad (\because |\vec{a}| = |\vec{b}|) \end{aligned}$$

63 (a)

$$\begin{aligned} \text{We have,} \\ \vec{a} \cdot \vec{b} &= \vec{a} \cdot \vec{c} \text{ and } \vec{a} \times \vec{b} = \vec{a} \times \vec{c} \\ \Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) &= 0 \text{ and } \vec{a} \times (\vec{b} - \vec{c}) = 0 \\ \Rightarrow (\vec{b} - \vec{c} = 0 \text{ or, } \vec{b} - \vec{c} \perp \vec{a}) \text{ and } (\vec{b} - \vec{c} \\ &= 0 \text{ or, } \vec{b} - \vec{c} \parallel \vec{a}) \\ \Rightarrow \vec{b} - \vec{c} &= 0 \Rightarrow \vec{b} = \vec{c} \end{aligned}$$

67 (a)

We know that, any vector  $\vec{a}$  can be uniquely expressed in terms of three non-coplanar vectors  $\hat{i}, \hat{j}$  and  $\hat{k}$ , we get

$$\vec{a} = x\hat{i} + y\hat{j} + z\hat{k} \text{ multiply in succession by } \hat{i}, \hat{j} \text{ and } \hat{k}, \text{ we get}$$

$$x = \vec{a} \cdot \hat{i}, y = \vec{a} \cdot \hat{j}, z = \vec{a} \cdot \hat{k}$$

$$\therefore (\vec{a} \cdot \hat{i})\hat{i} + (\vec{a} \cdot \hat{j})\hat{j} + (\vec{a} \cdot \hat{k})\hat{k} = x\hat{i} + y\hat{j} + z\hat{k} = \vec{a}$$

69 (b)

$$\text{Let } \vec{b} = \hat{i} \text{ and } \vec{c} = \hat{j}$$

$$\therefore |\vec{b} \times \vec{c}| = |\hat{k}| = 1$$

$$\text{Let } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\text{Now, } \vec{a} \cdot \vec{b} = \vec{a} \cdot \hat{i} = a_1, \vec{a} \cdot \vec{c} = \vec{a} \cdot \hat{j} = a_2$$

$$\text{and } \vec{a} \cdot \frac{\vec{b} \times \vec{c}}{|\vec{b} \times \vec{c}|} = \vec{a} \cdot \hat{k} = a_3$$

$$\therefore (\vec{a} \cdot \vec{b})\vec{b} + (\vec{a} \cdot \vec{c})\vec{c} + \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{|\vec{b} \times \vec{c}|} \cdot (\vec{b} \times \vec{c})$$

$$= a_1\vec{b} + a_2\vec{c} + a_3(\vec{b} \times \vec{c})$$

$$= a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = \vec{a}$$

70 (c)

Let the unit vector  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$  is perpendicular to  $\hat{i} - \hat{j}$ ,

then we get

$$\frac{(\hat{i} + \hat{j}) \cdot (\hat{i} - \hat{j})}{\sqrt{2}} = \frac{1 - 1}{\sqrt{2}} = 0$$

$\therefore \frac{\hat{i} + \hat{j}}{\sqrt{2}}$  is the required unit vector.

71 (d)

Let the unit vector be  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

$$\text{Since, } \vec{r} \cdot (3\hat{i} + \hat{j} + 2\hat{k}) = 0 \text{ and } \vec{r} \cdot (2\hat{i} - 2\hat{j} + 4\hat{k}) = 0$$

$$\Rightarrow 3x + y + 2z = 0 \text{ and } 2x - 2y + 4z = 0$$

On solving, we get  $x = 1, y = -1$  and  $z = -1$

$$\therefore \text{Required unit vector} = \frac{\hat{i} - \hat{j} - \hat{k}}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{\hat{i} - \hat{j} - \hat{k}}{\sqrt{3}}$$

72 (d)

The position vector of the vertices  $A, B, C$  of  $\Delta ABC$  are  $7\hat{j} + 10\hat{k}, -\hat{i} + 6\hat{j} + 6\hat{k}$  and  $-4\hat{i} + 9\hat{j} + 6\hat{k}$  respectively.

$$\therefore \vec{AB} = -\hat{i} - \hat{j} - 4\hat{k}, \vec{BC} = -3\hat{i} + 3\hat{j}$$

$$\text{And } \vec{CA} = 4\hat{i} - 2\hat{j} - 4\hat{k}$$

$$\Rightarrow |\vec{AB}| = \sqrt{(-1)^2 + (-1)^2 + (-4)^2} = \sqrt{18} = 3\sqrt{2}$$

$$|\vec{BC}| = \sqrt{(-3)^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$$

$$\text{and } |\vec{CA}| = \sqrt{4^2 + (-2)^2 + (-4)^2} = \sqrt{36} = 6$$

It is clear from these values that

$$|\vec{AB}|^2 + |\vec{BC}|^2 = |\vec{CA}|^2$$

Hence,  $\Delta ABC$  is right angled and isosceles also.

(b)

For collinearity,  $\cos x \hat{i} + \sin x \hat{j} = \lambda(x\hat{i} + \sin x \hat{j})$

$$\Rightarrow \cos x = x$$

$$\text{Let } f(x) = \cos x - x$$

$$\Rightarrow f'(x) = -\sin x - 1 < 0$$

$f(x)$  is decreasing function and for  $x \geq \frac{\pi}{3}, f(x) < 0$

and for  $\frac{\pi}{3} < x < \frac{\pi}{6}, f(x) > 0$ .

Hence, unique solution exist.

75 (d)

Let the required unit vector be  $\vec{r} = a\hat{i} + b\hat{j}$

$$\text{Then, } |\vec{r}| = 1$$

$$\Rightarrow a^2 + b^2 = 1 \dots(i)$$

Since,  $\vec{r}$  makes an angle of  $45^\circ$  with  $\hat{i} + \hat{j}$  and an angle of  $60^\circ$  with  $3\hat{i} - 4\hat{j}$ , therefore

$$\cos \frac{\pi}{4} = \frac{\vec{r} \cdot (\hat{i} + \hat{j})}{|\vec{r}| |\hat{i} + \hat{j}|}$$

$$\text{and } \cos \frac{\pi}{3} = \frac{\vec{r} \cdot (3\hat{i} - 4\hat{j})}{|\vec{r}| |3\hat{i} - 4\hat{j}|}$$

$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{a + b}{\sqrt{2}}$$

$$\text{and } \frac{1}{2} = \frac{3a - 4b}{5}$$

$$\Rightarrow a + b = 1$$

$$\text{and } 3a - 4b = \frac{5}{2}$$

$$\Rightarrow a = \frac{13}{14}, b = \frac{1}{14}$$

$$\therefore \vec{r} = \frac{13}{14}\hat{i} + \frac{1}{14}\hat{j}$$

77 (b)

Since, volume of parallelopiped = 34

$$\therefore \begin{vmatrix} 4 & 5 & 1 \\ 0 & -1 & 1 \\ 3 & 9 & p \end{vmatrix} = 34$$

$$\Rightarrow 4(-p - 9) - 5(-3) + 1(3) = 34$$

$$\Rightarrow -4p - 36 + 15 + 3 = 34$$

$$\Rightarrow 4p = -52$$

$$\Rightarrow p = -13$$

78 (d)

$$\frac{(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2}{2\vec{a}^2\vec{b}^2} = \frac{\vec{a}^2\vec{b}^2\sin^2\theta + \vec{a}^2\vec{b}^2\cos^2\theta}{2\vec{a}^2\vec{b}^2} = \frac{\cos^2\theta + \sin^2\theta}{2} = \frac{1}{2}$$

79 (b)

Let two vectors are  $\vec{a}$  and  $\vec{b}$

$$\text{Given, } |\vec{a} \times \vec{b}| = \sqrt{3} |\vec{a} \cdot \vec{b}|$$

$$\Rightarrow |\vec{a}| \cdot |\vec{b}| \sin \theta = \sqrt{3} |\vec{a}| |\vec{b}| \cos \theta$$


$$\Rightarrow \tan \theta = \sqrt{3}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

82 (b)

We have,

$$\vec{AC} = \vec{a} + \vec{b}, \vec{AD} = 2\vec{b}$$



In  $\triangle ADE$ , we have

$$\vec{AD} = \vec{DE} = \vec{AE} \Rightarrow 2\vec{b} - \vec{a} = \vec{AE} \Rightarrow \vec{EA} = \vec{a} - 2\vec{b}$$

In  $\triangle ACD$ , we have

$$\vec{AC} + \vec{CD} = \vec{AD} \Rightarrow \vec{a} + \vec{b} + \vec{CD} = 2\vec{b} \Rightarrow \vec{CD} = \vec{b} - \vec{a}$$

$$\therefore \vec{FA} = -\vec{CD} = \vec{a} - \vec{b}$$

Hence,  $\vec{AC} + \vec{AD} + \vec{EA} + \vec{FA}$

$$= \vec{a} + \vec{b} + 2\vec{b} + \vec{a} - 2\vec{b} + \vec{a} - \vec{b} = 3\vec{a} = 3\vec{AB}$$

83 (c)

We have,

$$(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c}) = \{(\vec{a} \times \vec{b}) \cdot \vec{c}\}\vec{b} - \{(\vec{a} \times \vec{b}) \cdot \vec{b}\}\vec{c} = [\vec{a} \vec{b} \vec{c}]\vec{b}$$

$$(\vec{b} \times \vec{c}) \times (\vec{c} \times \vec{a})$$

$$= \{(\vec{b} \times \vec{c}) \cdot \vec{a}\} \cdot \vec{c} - \{(\vec{b} \times \vec{c}) \cdot \vec{c}\}\vec{a}$$

$$= [\vec{b} \vec{c} \vec{a}]\vec{c}$$

and,

$$(\vec{c} \times \vec{a}) \times (\vec{a} \times \vec{b}) = \{(\vec{c} \times \vec{a}) \cdot \vec{b}\}\vec{a} - \{(\vec{c} \times \vec{a}) \cdot \vec{a}\}\vec{b} = [\vec{c} \vec{a} \vec{b}]\vec{a}$$

$$\therefore [(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c})](\vec{b} \times \vec{c}) \times (\vec{c} \times \vec{a})(\vec{c} \times \vec{a}) \times (\vec{a} \times \vec{b})]$$

$$= [[\vec{a} \vec{b} \vec{c}]\vec{a} [\vec{a} \vec{b} \vec{c}]\vec{b} [\vec{a} \vec{b} \vec{c}]\vec{c}]$$

$$= [\vec{a} \vec{b} \vec{c}]^3 [\vec{a} \vec{b} \vec{c}] = [\vec{a} \vec{b} \vec{c}]^4$$

84 (a)

$$\text{Given, } \vec{a} = \lambda\hat{i} - 7\hat{j} + 3\hat{k}, \vec{b} = \lambda\hat{i} + \hat{j} + 2\lambda\hat{k}$$

$$\therefore \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|}$$

$$= \frac{\lambda^2 - 7 + 6\lambda}{\sqrt{\lambda^2 + 49 + 9\sqrt{\lambda^2 + 1 + 4\lambda^2}}} < 0$$

$$\Rightarrow (\lambda + 7)(\lambda - 1) < 0$$

$$\Rightarrow -7 < \lambda < 1$$

85 (b)

We have,

$$\vec{r} = \lambda_1 \vec{r}_1 + \lambda_2 \vec{r}_2 + \lambda_3 \vec{r}_3$$

$$\Rightarrow 2\vec{a} - 3\vec{b} + 4\vec{c}$$

$$= (\lambda_1 - \lambda_2 + \lambda_3)\vec{a}$$

$$+ (-\lambda_1 + \lambda_2 - \lambda_3)\vec{b}$$

$$+ (\lambda_1 + \lambda_2 + \lambda_3)\vec{c}$$

$$\Rightarrow \lambda_1 - \lambda_2 + \lambda_3 = 2, -\lambda_1 + \lambda_2 - \lambda_3 =$$

$$-3, \lambda_1 + \lambda_2 + \lambda_3 = 4$$

$[\because \vec{a}, \vec{b}, \vec{c}$  are non-coplanar]

$$\Rightarrow \lambda_1 = \frac{7}{2}, \lambda_2 = 1, \lambda_3 = -\frac{1}{2}$$

86 (c)

We have,

$$(\vec{a} - \vec{d}) \cdot (\vec{b} - \vec{c}) = (\vec{b} - \vec{d}) \cdot (\vec{c} - \vec{a}) = 0$$

$$\Rightarrow \vec{DA} \cdot \vec{BC} = 0 \text{ and } \vec{DB} \cdot \vec{AC} = 0$$

$$\Rightarrow AD \perp BC \text{ and } DB \perp AC$$

$$\Rightarrow D \text{ is the orthocenter of } \triangle ABD$$

87 (a)

$$\text{Given } \vec{OA} = 4\hat{i} + 3\hat{j} - \hat{k}$$

$$\vec{OB} = -3\hat{i} + \hat{j} + \hat{k}$$

$$\therefore \vec{AB} = \vec{OB} - \vec{OA} = -7\hat{i} - 2\hat{j} + 2\hat{k}$$

$$\therefore \vec{DE} = -\vec{AB} = 7\hat{i} + 2\hat{j} - 2\hat{k}$$

88 (c)

$$\text{Given, } |\vec{a} \times \vec{b}| = |\vec{a} \cdot \vec{b}|$$

$$\Rightarrow |\vec{a}| \cdot |\vec{b}| \sin \theta = |\vec{a}| |\vec{b}| \cos \theta$$

$$\Rightarrow \sin \theta = \cos \theta \Rightarrow \theta = \frac{\pi}{4}$$

89 (a)

Let the line joining the points with position

vectors  $-2\hat{i} + 3\hat{j} + 5\hat{k}$  and  $7\hat{i} - \hat{k}$  be

Divide in the ratio  $\lambda:1$  by  $\hat{i} + 2\hat{j} + 3\hat{k}$

$$\therefore \frac{\lambda(7\hat{i} - \hat{k}) + (-2\hat{i} + 3\hat{j} + 5\hat{k})}{\lambda + 1} = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\Rightarrow (7\lambda - 2)\hat{i} + 3\hat{j} + (5 - \lambda)\hat{k}$$

$$= (\lambda + 1)\hat{i} + 2(\lambda + 1)\hat{j} + 3(\lambda$$

$$+ 1)\hat{k}$$

On equating the coefficient of  $\hat{i}$ , we get

$$7\lambda - 2 = \lambda + 1 \Rightarrow \lambda = 2$$

Hence, required ratio =  $\lambda:1 = 2:1$

91 (a)

$$\text{Force } \vec{F} = \vec{AB} = (3 - 1)\hat{i} + (-4 - 2)\hat{j} + (2 + 3)\hat{k} = 2\hat{i} - 6\hat{j} + 5\hat{k}$$

Moment of force  $\vec{F}$  with respect to  $M = \vec{MA} \times \vec{F}$

$$\therefore \vec{MA} = (1 + 2)\hat{i} + (2 - 4)\hat{j} + (-3 + 6)\hat{k}$$

$$= 3\hat{i} - 2\hat{j} + 3\hat{k}$$

$$\text{Now, } \vec{MA} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -2 & 3 \\ 2 & -6 & 5 \end{vmatrix}$$

$$= \hat{i}(-10 + 18) + \hat{j}(6 - 15) + \hat{k}(-18 + 4)$$

$$= 8\hat{i} - 9\hat{j} - 14\hat{k}$$

92 (d)

$$\therefore \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \frac{5\pi}{6} = -\frac{|\vec{a}| |\vec{b}| \sqrt{3}}{2}$$

$$\therefore -\frac{6}{\sqrt{3}} = -\frac{|\vec{a}| |\vec{b}| \sqrt{3}}{2} \quad (\text{given condition})$$

$$\Rightarrow |\vec{a}| = \frac{6 \times 2}{3} = 4$$

93 (c)

Equation of straight line passing through the points

$$a_1\hat{i} + a_2\hat{j} + a_3\hat{k} \text{ and } b_1\hat{i} + b_2\hat{j} + b_3\hat{k} \text{ is}$$

$$a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + (b_1\hat{i} + b_2\hat{j} + b_3\hat{k})t$$

95 (d)

$$(3\vec{a} + \vec{b}) \cdot (\vec{a} - 4\vec{b}) = 3|\vec{a}|^2 - 11\vec{a} \cdot \vec{b} - 4|\vec{b}|^2$$

$$= 3 \cdot 36 - 11 \cdot 6 \cdot 8 \cos \pi - 4 \cdot 64 > 0$$

$\therefore$  Angle between  $\vec{a}$  and  $\vec{b}$  is acute angle.

$\therefore$  The longer diagonal is given by

$$\vec{\alpha} = (3\vec{a} + \vec{b}) + (\vec{a} - 4\vec{b}) = 4\vec{a} - 3\vec{b}$$

$$\text{Now, } |\vec{\alpha}|^2 = |4\vec{a} - 3\vec{b}|^2$$

$$= 16|\vec{a}|^2 + 9|\vec{b}|^2 - 24\vec{a} \cdot \vec{b}$$

$$= 16 \cdot 36 + 9 \cdot 64 - 24 \cdot 6 \cdot 8 \cos \pi$$

$$= 16 \times 144$$

$$|4\vec{a} - 3\vec{b}| = 48$$

96 (b)

$$\text{Given, } \vec{a} + \vec{b} = -\vec{c}$$

$$\Rightarrow \vec{a} \times \vec{c} + \vec{b} \times \vec{c} = \vec{0} \Rightarrow \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$$

$$\text{Similarly, } \vec{a} \times \vec{b} = \vec{b} \times \vec{c}$$

$$\text{Hence, } \vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$$

97 (d)

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}| \cos 90^\circ$$

$$25 + 25 - 2 \times 0 = 50$$

$$\Rightarrow |\vec{a} - \vec{b}| = 5\sqrt{2}$$

98 (d)

Given vectors are non-coplanar, if

$$\Delta = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} \neq 0$$

$$\text{Now, } \begin{vmatrix} a & a^2 & 1+a^3 \\ b & b^2 & 1+b^3 \\ c & c^2 & 1+c^3 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} a & a^2 & 1 \\ b & b^2 & 1 \\ c & c^2 & 1 \end{vmatrix} + \begin{vmatrix} a & a^2 & a^3 \\ b & b^2 & b^3 \\ c & c^2 & c^3 \end{vmatrix} = 0$$

$$\Rightarrow \Delta(1+abc) = 0 \Rightarrow abc = -1$$

99 (a)

$$\text{Let } \vec{A} = \vec{a} + \vec{b} = (\hat{i} + \hat{j} + \hat{k}) + (\hat{i} + 3\hat{j} + 5\hat{k})$$

$$= 2\hat{i} + 4\hat{j} + 6\hat{k}$$

$$\text{and } \vec{B} = \vec{b} + \vec{c} = (\hat{i} + 3\hat{j} + 5\hat{k}) + (7\hat{i} + 9\hat{j} + 11\hat{k})$$

$$= 8\hat{i} + 12\hat{j} + 16\hat{k}$$

$$\therefore \text{Area of parallelogram} = \frac{1}{2} ||\vec{A} \times \vec{B}||$$

$$= \frac{1}{2} \left\| \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 4 & 6 \\ 8 & 12 & 16 \end{vmatrix} \right\|$$

$$= \frac{1}{2} |-8\hat{i} + 16\hat{j} - 8\hat{k}|$$

$$= \sqrt{(-4)^2 + (8)^2 + (-4)^2}$$

$$= 4\sqrt{6} \text{ sq units}$$

100 (a)

Clearly,  $\vec{c}$  is a unit vector parallel to the vector  $\vec{a} \times (\vec{a} \times \vec{b})$

$$\text{i.e. } \vec{c} = \pm \frac{\vec{a} \times (\vec{a} \times \vec{b})}{|\vec{a} \times (\vec{a} \times \vec{b})|}$$

We have,

$$\vec{a} = \hat{i} + \hat{j} - \hat{k} \text{ and } \vec{b} = \hat{i} - \hat{j} + \hat{k}$$

$$\therefore \vec{a} \times (\vec{a} \times \vec{b}) = (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}$$

$$\Rightarrow \vec{a} \times (\vec{a} \times \vec{b}) = -\vec{a} - 3\vec{b} = -4\hat{i} + 2\hat{j} - 2\hat{k}$$

$$\therefore \vec{c} = \pm \frac{(-4\hat{i} + 2\hat{j} - 2\hat{k})}{\sqrt{16 + 4 + 4}} = \pm \frac{1}{\sqrt{6}} (-2\hat{i} + \hat{j} - \hat{k})$$

102 (b)

$$\text{Given, } \vec{a} + 2\vec{b} + 4\vec{c} = \vec{0}$$

$$\text{Now, } \vec{a} \times (\vec{a} + 2\vec{b} + 4\vec{c}) = \vec{0}$$

$$\Rightarrow 2(\vec{a} \times \vec{b}) + 4(\vec{a} \times \vec{c}) = \vec{0}$$

$$\Rightarrow \frac{(\vec{a} \times \vec{b})}{4} = \frac{(\vec{c} \times \vec{a})}{2} \quad \dots (i)$$

$$\text{Again, } \vec{b} \times (\vec{a} + 2\vec{b} + 4\vec{c}) = \vec{0}$$

$$\Rightarrow \vec{b} \times \vec{a} + 4(\vec{b} \times \vec{c}) = \vec{0}$$

$$\Rightarrow \vec{b} \times \vec{c} = (\vec{a} \times \vec{b})/4 \quad \dots (ii)$$

From Eqs. (i) and (ii)

$$(\vec{a} \times \vec{b})/4 = \vec{b} \times \vec{c} = (\vec{c} \times \vec{a})/2 = \vec{p}$$

$$\therefore \vec{a} \times \vec{b} = 4\vec{p}, \vec{b} \times \vec{c} = \vec{p}$$

$$\text{and } \vec{c} \times \vec{a} = 2\vec{p}$$

$$\therefore (\vec{a} \times \vec{b}) + (\vec{b} \times \vec{c}) + (\vec{c} \times \vec{a}) = 4\vec{p} + \vec{p} + 2\vec{p}$$

$$= 7\vec{p} = 7(\vec{b} \times \vec{c})$$

$$\therefore \lambda = 7$$

103 (b)

$$\therefore \vec{F}_1 = \frac{5(6\hat{i} + 2\hat{j} + 3\hat{k})}{7}, \vec{F}_2 = \frac{3(3\hat{i} - 2\hat{j} + 6\hat{k})}{7}$$

$$\vec{F}_3 = \frac{1(2\hat{i} - 3\hat{j} - 6\hat{k})}{7}$$

$$\text{And } \vec{F} = \vec{F}_1 + \vec{F}_2 + \vec{F}_3$$

$$= \frac{1}{7}(30\hat{i} + 10\hat{j} + 15\hat{k} + 9\hat{i} - 6\hat{j} + 18\hat{k} + 2\hat{i} - 3\hat{j} - 6\hat{k})$$

$$= \frac{1}{7}(41\hat{i} + \hat{j} + 27\hat{k})$$

and  $\overline{\mathbf{AB}} = 5\hat{i} - \hat{j} + \hat{k} - 2\hat{i} + \hat{j} + 3\hat{k}$   
 $= 3\hat{i} + 4\hat{k}$

$\therefore$  Work done  $= \frac{1}{7} [41\hat{i} + \hat{j} + 27\hat{k}] \cdot [3\hat{i} + 4\hat{k}]$   
 $= \frac{1}{7} [123 + 108] = 33$  unit

104 (b)

Let  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ . Then,

$$\hat{i} \times (\vec{r} \times \hat{i}) + \hat{j} \times (\vec{r} \times \hat{j}) + \hat{k} \times (\vec{r} \times \hat{k})$$

$$= (\hat{i} \cdot \hat{i})\vec{r} - (\hat{i} \cdot \vec{r})\hat{i} + (\hat{j} \cdot \hat{j})\vec{r} - (\hat{j} \cdot \vec{r})\hat{j} + (\hat{k} \cdot \hat{k})\vec{r} - (\hat{k} \cdot \vec{r})\hat{k}$$

$$= \vec{r} - x\hat{i} + \vec{r} - y\hat{j} + \vec{r} - z\hat{k}$$

$$= 3\vec{r} - (x\hat{i} + y\hat{j} + z\hat{k}) = 3\vec{r} - \vec{r} = 2\vec{r}$$

105 (b)

The equation of the plane through the line of intersection of given plane is

$$(\vec{r} \cdot \vec{a} - \lambda) + k(\vec{r} \cdot \vec{b} - \mu) = 0$$

or  $\vec{r} \cdot (\vec{a} + k\vec{b}) = \lambda + k\mu$  ..... (i)

this passes through the origin, therefore

$$0 \cdot (\vec{a} + k\vec{b}) = \lambda + k\mu$$

$$\Rightarrow k = -\frac{\lambda}{\mu}$$

On putting the value of  $k$  in Eq. (i), we get the equation of the required plane as

$$\vec{r} \cdot (\mu\vec{a} - \lambda\vec{b}) = 0$$

$$0 \Rightarrow \vec{r} \cdot (\lambda\vec{b} - \mu\vec{a}) = 0$$

106 (c)

By the properties of scalar triple product

$$[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = 2[\vec{a}, \vec{b}, \vec{c}]$$

$$\therefore k = 2$$

107 (c)

$$\vec{a} \cdot \vec{a} = 1 + 1 + 1 = 3$$

Using,

$$\vec{a} \times (\vec{a} \times \vec{b}) = (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}$$

$$\therefore (\hat{i} + \hat{j} + \hat{k}) \times (\hat{j} - \hat{k}) = (\hat{i} + \hat{j} + \hat{k}) - 3\vec{b}$$

$$\Rightarrow -2\hat{i} + \hat{j} + \hat{k} = \hat{i} + \hat{j} + \hat{k} - 3\vec{b}$$

$$\Rightarrow \vec{b} = \hat{i}$$

108 (a)

Vector perpendicular to face  $OAB$  is  $\vec{n}_1$

$$= \overline{\mathbf{OA}} \times \overline{\mathbf{OB}} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 1 \\ 2 & 1 & 3 \end{vmatrix}$$

$$= 5\hat{i} - \hat{j} - 3\hat{k}$$

Vector perpendicular to face  $ABC$  is  $\vec{n}_2$

$$= \overline{\mathbf{AB}} \times \overline{\mathbf{AC}} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ -2 & -1 & 1 \end{vmatrix}$$

$$= \hat{i} - 5\hat{j} - 3\hat{k}$$

$$\therefore \cos \theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|}$$

$$= \frac{5 \times 1 + (-1) \times (-5) + (-3) \times (-3)}{\sqrt{5^2 + (-1)^2 + (-3)^2} \sqrt{1^2 + (-5)^2 + (-3)^2}}$$

$$= \frac{5 + 5 + 9}{\sqrt{35} \sqrt{35}} = \frac{19}{35}$$

$$\Rightarrow \theta = \cos^{-1} \left( \frac{19}{35} \right)$$

109 (a)

Given,  $2\vec{a} + 3\vec{b} - 5\vec{c} = \vec{0}$

$$\Rightarrow \frac{2\vec{a} + 3\vec{b}}{5} = \vec{c}$$

$$\Rightarrow \frac{2\vec{a} + 3\vec{b}}{2 + 3} = \vec{c}$$

$$\Rightarrow \frac{\vec{a} + \frac{3}{2}\vec{b}}{1 + \frac{3}{2}} = \vec{c} \quad \dots \dots \dots (i)$$


Let  $\vec{c}$  divides  $\overline{\mathbf{AB}}$  in the ratio  $\lambda:1$

Then,  $\vec{c} = \frac{\lambda\vec{a} + \vec{b}}{1 + \lambda}$  ..... (ii)

On comparing Eqs.(i) and (ii), we get

$$\lambda = \frac{3}{2}$$

$\therefore$  Required ratio is 3:2 internally.



110 (a)

Let  $\overline{\mathbf{OA}} = \hat{i} + \hat{j} + \hat{k}$ ,  $\overline{\mathbf{OB}} = 5\hat{i} + 3\hat{j} - 3\hat{k}$  and  $\overline{\mathbf{OC}} = 2\hat{i} + 5\hat{j} + 9\hat{k}$

$$\therefore \overline{\mathbf{AB}} = 4\hat{i} + 2\hat{j} - 4\hat{k}, \overline{\mathbf{BC}} = -3\hat{i} + 2\hat{j} + 12\hat{k} \quad \text{and}$$

$$\overline{\mathbf{AC}} = \hat{i} + 4\hat{j} + 8\hat{k}$$

$$\Rightarrow AB = 6, BC = \sqrt{157}, AC = 9$$

$$\therefore \text{Perimeter of } \Delta ABC = 15 + \sqrt{157}$$

111 (c)

Given,  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\therefore |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = 0$$

$$\Rightarrow 25 + 16 + 9 + 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = 0$$

$$\Rightarrow 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = -50$$

$$\Rightarrow [\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = -25$$

113 (d)

It is given that  $\vec{a} + \vec{b}$  is collinear with  $\vec{c}$  and  $\vec{b} + \vec{c}$  is collinear with  $\vec{a}$

$\therefore \vec{a} + \vec{b} = \lambda \vec{c}$  and  $\vec{b} + \vec{c} = \mu \vec{a}$  for some scalars  $\lambda$  and  $\mu$

$$\Rightarrow \vec{b} + \vec{c} = \mu(\lambda \vec{c} - \vec{b}) \quad [\text{On eliminating } \vec{a}]$$

$$\Rightarrow (\mu + 1)\vec{b} + (1 - \mu\lambda)\vec{c} = \vec{0}$$

$\Rightarrow \mu + 1 = 0$  and  $\mu\lambda = 1$  [ $\because \vec{b}$  and  $\vec{c}$  are non-collinear]

$$\Rightarrow \mu = -1 \text{ and } \lambda = -1$$

$$\therefore \vec{a} + \vec{b} + \vec{c} = \vec{0} \quad [\text{Putting } \lambda = -1 \text{ in } \vec{a} + \vec{b} = \lambda \vec{c}]$$

114 (b)

$$\text{Let } \vec{r} = l(\vec{b} \times \vec{c}) + m(\vec{c} \times \vec{a}) + n(\vec{a} \times \vec{b})$$

$$\vec{r} \cdot \vec{a} = l[\vec{a} \cdot \vec{b} \times \vec{c}]$$

$$\Rightarrow l = 1$$

Similarly,  $m = 2, n = 3$

115 (b)

$$\text{Given, } |\vec{x}| = |\vec{y}| = |\vec{z}| = 2$$

$$\text{and } \vec{x} = -\vec{y} - \vec{z}$$

$$\Rightarrow |\vec{x}|^2 = |\vec{y}|^2 + |\vec{z}|^2 + 2|\vec{y}||\vec{z}|\cos\theta$$

$$\Rightarrow 4 = 4 + 4 + 2 \times 2 \times 2 \cos\theta$$

$$\Rightarrow \cos\theta = -\frac{1}{2}$$

$$\Rightarrow \theta = 120^\circ$$

$$\text{Now, } \operatorname{cosec}^2\theta + \cot^2\theta = \operatorname{cosec}^2 120^\circ + \cot^2 120^\circ$$

$$= \left(\frac{2}{\sqrt{3}}\right)^2 + \left(-\frac{1}{\sqrt{3}}\right)^2 = \frac{5}{3}$$

116 (b)

$$\text{Given, } \vec{a} \cdot \vec{b} = -|\vec{a}||\vec{b}|$$

$$\Rightarrow |\vec{a}||\vec{b}|\cos\theta = -|\vec{a}||\vec{b}|$$

$$\Rightarrow \cos\theta = -1$$

$$\Rightarrow \theta = 180^\circ$$

117 (b)

$$\text{Let } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\text{Given, } \vec{r} \times \vec{b} = \vec{c} \times \vec{b}$$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \times (\hat{i} + \hat{j} + \hat{k})$$

$$= (4\hat{i} - 3\hat{j} + 7\hat{k}) \times (\hat{i} + \hat{j} + \hat{k})$$

$$\Rightarrow (y - z)\hat{i} - (x - z)\hat{j} + (x - y)\hat{k}$$

$$= -10\hat{i} + 3\hat{j} + 7\hat{k}$$

$$\Rightarrow y - z = -10, -(x - z) = 3, x - y = 7$$

$$\Rightarrow y - z = -10, -x + z = 3, x - y = 7 \dots (i)$$

$$\text{and } \vec{r} \cdot \vec{a} = 0$$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + \hat{k})$$

$$\Rightarrow 2x + z = 0 \dots (ii)$$

From Eqs. (i) and (ii), we get

$$x = -1, y = -8, z = 2$$

$$\therefore \vec{r} \cdot \vec{b} = (-\hat{i} - 8\hat{j} + 2\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k})$$

$$= -1 - 8 + 2$$

$$= -7$$

118 (d)

Since, given vectors are coplanar so it can be written as

$$\vec{a} + \lambda \vec{b} + 3\vec{c} = x(-2\vec{a} + 3\vec{b} - 4\vec{c}) + y(\vec{a} - 3\vec{b} + 5\vec{c})$$

On comparing the coefficient of  $\vec{a}, \vec{b}$  and  $\vec{c}$  on both sides, we get

$$-2x + y = 1; 3x - 3y = \lambda \text{ and } -4x + 5y = 3$$

On solving, we get

$$x = -\frac{1}{3}, y = \frac{1}{3}, \lambda = -2$$

119 (d)

Since,  $\vec{A} + \vec{B}$  is collinear to  $\vec{C}$  and  $\vec{B} + \vec{C}$  is collinear to  $\vec{A}$

$$\therefore \vec{A} + \vec{B} = \lambda \vec{C} \text{ and } \vec{B} + \vec{C} = \mu \vec{A}$$

Where  $\lambda$  and  $\mu$  are scalars.

$$\Rightarrow \vec{A} + \vec{B} + \vec{C} = (\lambda + 1)\vec{C}$$

$$\text{and } \vec{A} + \vec{B} + \vec{C} = (\mu + 1)\vec{A}$$

$$\Rightarrow (\lambda + 1)\vec{C} = (\mu + 1)\vec{A}$$

If  $\lambda \neq -1$ , then

$$\vec{C} = \frac{\mu + 1}{\lambda + 1} \vec{A}$$

$\Rightarrow \vec{C}$  and  $\vec{A}$  are collinear.

This is a contradiction to the given condition.

$$\therefore \lambda = -1$$

$$\therefore \vec{A} + \vec{B} + \vec{C} = \vec{0}$$

120 (d)

$$|\vec{AB}| = \sqrt{(7-1)^2 + (-4+6)^2 + (7-10)^2} = 7$$

$$|\vec{BC}| = \sqrt{(1+1)^2 + (-6+3)^2 + (10-4)^2} = 7$$

$$|\vec{CD}| = \sqrt{(-1-5)^2 + (-3+1)^2 + (4-5)^2} = \sqrt{41}$$

$$\text{and } |\vec{DA}| = \sqrt{(5-7)^2 + (-1+4)^2 + (5-7)^2} = \sqrt{17}$$

121 (c)

We have,

$$|\vec{a} + \vec{b}|^2 + |\vec{a} - \vec{b}|^2 = 2\{|\vec{a}|^2 + |\vec{b}|^2\}$$

$$\Rightarrow 300 + |\vec{a} - \vec{b}|^2 = 2(49 + 121)$$

$$\Rightarrow |\vec{a} - \vec{b}| = 2\sqrt{10}$$

123 (a)

We know, if  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ , then

$$\cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|}$$

$$= \frac{(2\hat{i} + 2\hat{j} - \hat{k}) \cdot (6\hat{i} - 3\hat{j} + 2\hat{k})}{\sqrt{2^2 + 2^2 + (-1)^2} \sqrt{6^2 + (-3)^2 + 2^2}}$$

$$= \frac{12 - 6 - 2}{\sqrt{4 + 4 + 1} \sqrt{36 + 9 + 4}}$$

$$= \frac{4}{\sqrt{9} \sqrt{49}} = \frac{2}{21}$$

124 (d)

If  $\vec{a} + 2\vec{b}$  is collinear with  $\vec{c}$ , then

$$\vec{a} + 2\vec{b} = t\vec{c} \quad \dots(i)$$

Also, if  $\vec{b} + 3\vec{c}$  is collinear with  $\vec{a}$  then

$$\vec{b} + 3\vec{c} = \lambda\vec{a}$$

$$\Rightarrow \vec{b} = \lambda\vec{a} - 3\vec{c} \quad \dots(ii)$$

On putting the value of  $\vec{b}$  in Eq. (i), we get

$$\vec{a} + 2(\lambda\vec{a} - 3\vec{c}) = t\vec{c}$$

$$\Rightarrow (\vec{a} - 6\vec{c}) = t\vec{c} - 2\lambda\vec{a}$$

On comparing, we get  $1 = -2\lambda$  and  $-6 = t$

$$\Rightarrow \lambda = -\frac{1}{2} \text{ and } t = -6$$

From Eq. (i)

$$\vec{a} + 2\vec{b} = -6\vec{c}$$

$$\Rightarrow \vec{a} + 2\vec{b} + 6\vec{c} = \vec{0}$$

125 (a)

We have,

$$\vec{AB} = -\hat{i} - \hat{j} - 4\hat{k}, \vec{BC} = -3\hat{i} + 3\hat{j} \text{ and } \vec{CA} = 4\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\therefore |\vec{AB}| = |\vec{BC}| = 3\sqrt{2} \text{ and } |\vec{CA}| = 6$$

$$\text{Clearly, } |\vec{AB}|^2 + |\vec{BC}|^2 = |\vec{CA}|^2$$

Hence, the triangle is right angled isosceles triangle

127 (c)

Since, three vectors  $(\vec{a} + 2\vec{b} + 3\vec{c})$ ,  $(\lambda\vec{b} + 4\vec{c})$  and  $(2\lambda - 1)\vec{c}$  are non-coplanar

$$\therefore \begin{vmatrix} 1 & 2 & 3 \\ 0 & \lambda & 4 \\ 0 & 0 & 2\lambda - 1 \end{vmatrix} \neq 0$$

$$\Rightarrow (2\lambda - 1)(\lambda) \neq 0$$

$$\Rightarrow \lambda \neq 0, \frac{1}{2}$$

Hence, these three vectors are non-coplanar for all except two values of  $\lambda$ .

128 (a)

$$\text{Given } \vec{PR} = 5\vec{PQ}$$

It means  $R$  divides  $PQ$  externally in the ratio 5:4

$$\therefore \text{Position vector of } R = \frac{5\vec{b} - 4\vec{a}}{5 - 4}$$

$$= 5\vec{b} - 4\vec{a}$$

130 (a)

$$\text{Let } \vec{OA} = \hat{i} + 2\hat{j} + 3\hat{k} \text{ and } \vec{OB} = 2\hat{i} - \hat{j} + 4\hat{k}$$

Let point  $C(x_1, y_1, z_1)$  divide  $AB$  in the ratio 1:2

$$\therefore x_1 = \frac{2 + 2}{1 + 2} = \frac{4}{3}, \quad y_1 = \frac{-1 + 4}{1 + 2} = \frac{3}{3} = 1$$

$$\text{and } z_1 = \frac{4 + 6}{1 + 2} = \frac{10}{3}$$

Again let point  $D(x_2, y_2, z_2)$  divides  $AB$  in the ratio 2:1, then

$$x_2 = \frac{4 + 1}{2 + 1} = \frac{5}{3}, \quad y_2 = \frac{-2 + 2}{2 + 1} = 0$$

$$\text{and } z_2 = \frac{8 + 3}{2 + 1} = \frac{11}{3}$$

So, position vector of the points of trisection of  $AB$  are position vector of

$$C = -\frac{4}{3}\hat{i} + \hat{j} + \frac{10}{3}\hat{k}$$

and position vector of

$$D = \frac{5}{3}\hat{i} + \frac{11}{3}\hat{k}$$

131 (a)

Let  $\vec{a}, \vec{b}, \vec{c}$  be the position vectors  $A, B$  and  $C$  respectively. Then, the position vector of  $G$  is

$\frac{\vec{a} + \vec{b} + \vec{c}}{3}$  and the position vectors of  $D, E$  and  $F$  are

$\frac{\vec{b} + \vec{c}}{2}, \frac{\vec{c} + \vec{a}}{2}$  and  $\frac{\vec{a} + \vec{b}}{2}$  respectively

$$\therefore \vec{GD} + \vec{GE} + \vec{GF}$$

$$= \left( \frac{\vec{b} + \vec{c}}{2} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right) + \left( \frac{\vec{c} + \vec{a}}{2} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right)$$

$$+ \left( \frac{\vec{a} + \vec{b}}{2} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right)$$

$$= (\vec{a} + \vec{b} + \vec{c}) - (\vec{a} + \vec{b} + \vec{c}) = \vec{0}$$

132 (a)

Let  $\vec{OA} = \vec{a}, \vec{OB} = \vec{b}$  and  $\vec{OC} = \vec{c}$ , then

$$\vec{OD} = \frac{\vec{a} + \vec{b}}{2}, \vec{OE} = \frac{\vec{a} + \vec{c}}{2}, \vec{OF} = \frac{\vec{b} + \vec{c}}{2}$$

$$\text{Now, } \vec{AF} = \frac{1}{2}(\vec{b} + \vec{c}) - \vec{a}, \vec{BE} = \frac{1}{2}(\vec{a} + \vec{c}) - \vec{b}$$

$$\text{and } \vec{CD} = \frac{1}{2}(\vec{a} + \vec{b}) - \vec{c}$$

$$\therefore \vec{AF} + \vec{BE} = \frac{1}{2}(\vec{b} + \vec{c}) - \vec{a} + \frac{1}{2}(\vec{a} + \vec{c}) - \vec{b}$$


$$= \vec{c} - \frac{1}{2}(\vec{a} + \vec{b}) = \vec{DC}$$

133 (c)

We have,

$$\vec{a} \cdot \vec{b} = 0 \Rightarrow \vec{a} \perp \vec{b}$$

So, vectors  $\vec{a}, \vec{b}$  and  $\vec{a} + \vec{b}$  form a right angled triangle



In  $\Delta PQR$ , we have

$$\tan 30^\circ = \frac{|\vec{b}|}{|\vec{a}|} \Rightarrow |\vec{a}| = 3|\vec{b}|$$

134 (d)

We have,  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$

$$\therefore \vec{b} = \hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k}) \dots \dots (i)$$

$$\begin{aligned} \text{Now, } \hat{i} \times (\vec{a} \times \hat{i}) &= (\hat{i} \cdot \hat{i})\vec{a} - (\hat{i} \cdot \vec{a})\hat{i} \\ &= 1(\hat{i} + 2\hat{j} + 3\hat{k}) - (1)\hat{i} \\ &= 2\hat{j} + 3\hat{k} \end{aligned}$$

$$\text{Similarly, } \hat{j} \times (\vec{a} \times \hat{j}) = \hat{i} + 3\hat{k}$$

$$\text{and } \hat{k} \times (\vec{a} \times \hat{k}) = \hat{i} + 2\hat{j}$$

$\therefore$  From Eq. (i),

$$\begin{aligned} \vec{b} &= 2\hat{j} + 3\hat{k} + \hat{i} + 3\hat{k} + \hat{i} + 2\hat{j} \\ &= 2\hat{i} + 4\hat{j} + 6\hat{k} \\ \Rightarrow |\vec{b}| &= \sqrt{4 + 16 + 36} = 2\sqrt{14} \end{aligned}$$

135 (a)

The centroid of triangle

$$\begin{aligned} &= \frac{(a\hat{i} + b\hat{j} + c\hat{k}) + (b\hat{i} + c\hat{j} + a\hat{k}) + (c\hat{i} + a\hat{j} + b\hat{k})}{3} \\ &= \frac{a + b + c}{3}(\hat{i} + \hat{j} + \hat{k}) \end{aligned}$$

136 (d)

$$\text{Given, } |\vec{a} + \vec{b}| = 1, |\vec{a}| = |\vec{b}| = 1$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}| = 1$$

$$\Rightarrow 2|\vec{a}||\vec{b}| = -1 \dots (i)$$

$$\text{Now, } |\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|$$

$$= 1^2 + 1^2 - (-1) = 3 \quad [\text{from Eq. (i)}]$$

$$\Rightarrow |\vec{a} + \vec{b}| = \sqrt{3}$$

137 (a)

Since,  $\vec{a}$  and  $\vec{b}$  are collinear vectors.

$$\therefore \vec{a} = \lambda\vec{b}$$

$$\Rightarrow \hat{i} - \hat{j} = \lambda(-2\hat{i} + m\hat{j})$$

$$\Rightarrow 1 = -2\lambda, -1 = \lambda m$$

$$\Rightarrow \lambda = -\frac{1}{2}, m = -\frac{1}{\lambda}$$

$$\Rightarrow m = 2$$

138 (c)

Since,  $C$  is the mid point of  $A(2, -1)$  and  $B(-4, 3)$ .

$$\therefore \text{Coordinates of } C \text{ is } \left(\frac{2-4}{2}, \frac{-1+3}{2}\right) = (-1, 1)$$

$$\therefore \vec{OC} = -\hat{i} + \hat{j}$$

139 (c)

According to the given conditions, we have

$$\vec{a} \cdot \vec{b} > 0 \text{ and } \vec{b} \cdot \hat{j} < 0$$

$$\Rightarrow 2x^2 - 3x + 1 > 0 \text{ and } x < 0$$

$$\Rightarrow (x < 1/2 \text{ or } x > 1) \text{ and } x < 0 \Rightarrow x < 0$$

140 (d)

$$\frac{|(\vec{a} - \vec{c}) \times (\vec{b} - \vec{a})|}{(\vec{c} - \vec{a}) \cdot (\vec{b} - \vec{a})} = \frac{|\vec{AC} \times \vec{BA}|}{\vec{AC} \cdot \vec{BA}}$$

$$= \frac{||\vec{AC}|||\vec{BA}|\sin A\hat{n}}{||\vec{AC}|||\vec{BA}|\cos A} = \tan A$$

141 (d)

$$\text{Let, } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\therefore \vec{a} \cdot \hat{i} = 1 \Rightarrow a_1 = 1$$

$$\text{Since, } \vec{a} \cdot (2\hat{i} + \hat{j}) = 1$$

$$\Rightarrow 2a_1 + a_2 = 1$$

$$\Rightarrow a_2 = 1 - 2$$

$$\Rightarrow a_2 = -1$$

$$\text{and } \vec{a} \cdot (\hat{i} + \hat{j} + 3\hat{k}) = 1$$

$$\Rightarrow a_1 + a_2 + 3a_3 = 1$$

$$\Rightarrow 1 - 1 + 3a_3 = 1$$

$$\Rightarrow a_3 = \frac{1}{3}$$

$$\therefore \vec{a} = \hat{i} - \hat{j} + \frac{1}{3}\hat{k} = \frac{1}{3}(3\hat{i} - 3\hat{j} + \hat{k})$$

142 (c)

$$\text{Given, } \vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} - \hat{j} + 2\hat{k}$$

$$\text{and } \vec{c} = x\hat{i} + (x-2)\hat{j} - \hat{k}$$

Since,  $\vec{c}$  lies in the plane of vectors  $\vec{a}$  and  $\vec{b}$

therefore  $\vec{a}, \vec{b}$  and  $\vec{c}$  are coplanar.

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ x & (x-2) & -1 \end{vmatrix} = 0$$

$$\Rightarrow 1(1 - 2x + 4) - 1(-1 - 2x) + 1(x - 2 + x) = 0$$

$$\Rightarrow 5 - 2x + 1 + 2x + 2x - 2 = 0$$

$$\Rightarrow x = -2$$

143 (d)

$$\text{Let } \vec{P} = \hat{i} + \hat{j} - \hat{k}, \vec{Q} = 2\hat{i} + 3\hat{j}, \vec{R} = 5\hat{j} - 2\hat{k}$$

$$\text{and } \vec{S} = -\hat{j} + \hat{k}$$

$$\therefore \vec{PQ} = \hat{i} + 2\hat{j} + \hat{k}$$

$$\Rightarrow |\vec{PQ}| = \sqrt{6}$$

$$\vec{QR} = -2\hat{i} + 2\hat{j} - 2\hat{k}$$

$$\Rightarrow |\vec{QR}| = \sqrt{12}$$

$$\text{and } \vec{RS} = -6\hat{j} + 3\hat{k}$$

$$\Rightarrow |\overline{RS}| = \sqrt{45}$$

$$\text{and } \overline{SP} = \hat{i} + 2\hat{j} - 2\hat{k} \Rightarrow |\overline{SP}| = 3$$

Which are not satisfied the conditions of any of the following. Trapezium, rectangle and parallelogram.

144 (c)

Clearly,

$$\text{Required vector} = |\vec{b}|\hat{a} = \frac{|\vec{b}|}{|\vec{a}|}\vec{a} = \frac{7}{3}(\hat{i} + 2\hat{j} + 2\hat{k})$$

145 (a)

If  $I$  is incentre of  $\Delta ABC$ . Then,

$$I \text{ is } \frac{a\vec{a} + b\vec{b} + c\vec{c}}{a + b + c}$$

147 (d)

For a parallel  $\vec{a} \times \vec{b} = 0$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 3 \\ 4 & -\lambda & 6 \end{vmatrix} = 0$$

$$\begin{aligned} \Rightarrow \hat{i}(6 + 3\lambda) - \hat{j}(0) + \hat{k}(-2\lambda - 4) &= 0 \\ = 0 \cdot \hat{i} + 0 \cdot \hat{j} + 0 \cdot \hat{k} \\ \therefore 6 + 3\lambda = 0 \Rightarrow \lambda &= -2 \end{aligned}$$

148 (b)

Total force,

$$\vec{F} = \frac{5(6\hat{i} + 2\hat{j} + 3\hat{k})}{7} + \frac{3(3\hat{i} - 2\hat{j} + 6\hat{k})}{7} + \frac{1(2\hat{i} - 3\hat{j} - 6\hat{k})}{7}$$

$$= \frac{1}{7}(41\hat{i} + \hat{j} + 27\hat{k})$$

$$\text{and } \overline{AB} = 5\hat{i} - \hat{j} + \hat{k} - 2\hat{i} + \hat{j} + 3\hat{k} = 3\hat{i} + 4\hat{k}$$

$$\therefore \text{Work done} = \vec{F} \cdot \overline{AB}$$

$$= \frac{1}{7}[41\hat{i} + \hat{j} + 27\hat{k}] \cdot [3\hat{i} + 4\hat{k}]$$

$$= \frac{1}{7}[123 + 108] = 33 \text{ units}$$

150 (d)

Since vectors  $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}$  and  $\vec{b} = 4\hat{i} - \lambda\hat{j} + 6\hat{k}$  are parallel

$$\therefore \frac{2}{4} = \frac{1}{-\lambda} = \frac{3}{6} \Rightarrow \lambda = -2$$

151 (b)

If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are coplanar vectors, then  $2\vec{a} - \vec{b}, 2\vec{b} - \vec{c}$  and  $2\vec{c} - \vec{a}$  are also coplanar.

$$\therefore [2\vec{a} - \vec{b}, 2\vec{b} - \vec{c}, 2\vec{c} - \vec{a}] = 0$$

152 (b)

$$\text{Here, } |\vec{a}| = \sqrt{1 + 1 + (4)^2} = 3\sqrt{2}$$

$$\text{and } |\vec{b}| = \sqrt{1 + (-1)^2 + (4)^2} = 3\sqrt{2}$$

$$\therefore |\vec{a}| = |\vec{b}|$$

$$\text{Now, } (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = |\vec{a}|^2 - |\vec{b}|^2 = 0$$


Hence, angle between them is  $90^\circ$

153 (a)

Given,

$$\overline{OQ} = (1 - 3\mu)\hat{i} + (\mu - 1)\hat{j} + (5\mu + 2)\hat{k}$$

$$\overline{OP} = 3\hat{i} + 2\hat{j} + 6\hat{k} \text{ (where } O \text{ is origin)}$$



Now,

$$\overline{PQ} = (1 - 3\mu - 3)\hat{i} + (\mu - 1 - 2)\hat{j} + (5\mu + 2 - 6)\hat{k}$$

$$= (-2 - 3\mu)\hat{i} + (\mu - 3)\hat{j} + (5\mu - 4)\hat{k}$$

$\therefore \overline{PQ}$  is parallel to the plane  $x - 4y + 3z = 1$

$$\therefore -2 - 3\mu - 4\mu + 12 + 15\mu - 12 = 0$$

$$\Rightarrow 8\mu = 2$$

$$\Rightarrow \mu = \frac{1}{4}$$

154 (b)

$$\text{Let } \vec{A} = \hat{i} - 2\hat{j} + 3\hat{k}, \vec{B} = -2\hat{i} + 3\hat{j} - \hat{k}$$

$$\text{and } \vec{C} = 4\hat{i} - 7\hat{j} + 7\hat{k}$$

$$\therefore \overline{AB} = -3\hat{i} + 5\hat{j} - 4\hat{k}$$

$$\text{and } \overline{AC} = 3\hat{i} - 5\hat{j} + 4\hat{k}$$

$$\therefore \text{Area of } \Delta ABC = \frac{1}{2} \|\overline{AB} \times \overline{AC}\|$$

$$= \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -3 & 5 & -4 \\ 3 & -5 & 4 \end{vmatrix} = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -3 & 5 & -4 \\ 0 & 0 & 0 \end{vmatrix}$$

[operating  $R_2 \rightarrow R_2 + R_3$ ]

$$= \frac{1}{2}[0] = 0$$

155 (b)

We have,

$$\vec{r} \times \vec{a} = \vec{b} \times \vec{a} \text{ and } \vec{r} \times \vec{b} = \vec{a} \times \vec{b}$$

$$\Rightarrow (\vec{r} - \vec{b}) \times \vec{a} = 0 \text{ and } (\vec{r} - \vec{a}) \times \vec{b} = 0$$

$$\Rightarrow \vec{r} - \vec{b} \parallel \vec{a} \text{ and } \vec{r} - \vec{a} \parallel \vec{b}$$

$$\Rightarrow \vec{r} - \vec{b} = \lambda\vec{a} \text{ and } \vec{r} - \vec{a} = \mu\vec{b} \text{ for some } \lambda, \mu \in \mathbb{R}$$

$$\Rightarrow \vec{r} = \vec{b} + \lambda\vec{a} \text{ and } \vec{r} = \vec{a} + \mu\vec{b} \text{ for some } \lambda, \mu \in \mathbb{R}$$

$$\Rightarrow \vec{b} + \lambda\vec{a} = \vec{a} + \mu\vec{b}$$

$$\Rightarrow \lambda = \mu = 1 \quad [\because \vec{a}, \vec{b} \text{ are non-collinear}]$$


$$\therefore \vec{r} = \vec{a} + \vec{b}$$

156 (c)

$$\begin{aligned}
& |\vec{a} + \vec{b} + \vec{c}|^2 \\
&= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{a} \cdot \vec{b} + 2\vec{b} \cdot \vec{c} + 2\vec{c} \cdot \vec{a} \\
&\Rightarrow 0 = 1 + 1 + 1 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) \\
&[\because |\vec{a}| = |\vec{b}| = |\vec{c}| = 1, \text{ given}] \\
&\therefore \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{3}{2}
\end{aligned}$$

157 (a)

The volume of the parallelepiped with coterminous edges as  $\hat{a}, \hat{b}, \hat{c}$  is given by  $[\hat{a}, \hat{b}, \hat{c}] = \hat{a} \cdot (\hat{b} \times \hat{c})$



$$\text{Now, } [\hat{a}, \hat{b}, \hat{c}]^2 = \begin{vmatrix} \hat{a} \cdot \hat{a} & \hat{a} \cdot \hat{b} & \hat{a} \cdot \hat{c} \\ \hat{b} \cdot \hat{a} & \hat{b} \cdot \hat{b} & \hat{b} \cdot \hat{c} \\ \hat{c} \cdot \hat{a} & \hat{c} \cdot \hat{b} & \hat{c} \cdot \hat{c} \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 1/2 & 1/2 \\ 1/2 & 1 & 1/2 \\ 1/2 & 1/2 & 1 \end{vmatrix} = \frac{1}{2}$$

$$[\because |\vec{a}| = |\vec{b}| = |\vec{c}| = 1]$$

$$\Rightarrow [\hat{a}, \hat{b}, \hat{c}]^2 = \frac{1}{2}$$

Thus, the required volume of the parallelepiped

$$= \frac{1}{\sqrt{2}} \text{ cu unit}$$

158 (d)

We have,  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$

and  $\vec{b} = \hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k})$

$$= 3\vec{a} - \vec{a} = 2\vec{a}$$

$$= 2(\hat{i} + 2\hat{j} + 3\hat{k})$$

$$\Rightarrow |\vec{b}| = \sqrt{4 + 16 + 36} = \sqrt{56} = 2\sqrt{14}$$

159 (b)

Let,  $\vec{a} = 2p\hat{i} + \hat{j}$ ,  $\vec{b} = (p+1)\hat{i} + \hat{j}$

Given,  $|\vec{a}| = |\vec{b}| \Rightarrow 4p^2 + 1 = (p+1)^2 + 1$

$$\Rightarrow 3p^2 - 2p - 1 = 0 \Rightarrow p = 1, -\frac{1}{3}$$

160 (c)

Since  $\vec{r}_1, \vec{r}_2, \vec{r}_3$  are coplanar

$$\therefore [\vec{r}_1 \vec{r}_2 \vec{r}_3] = 0$$

$$\Rightarrow \begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0 \Rightarrow abc = a + b + c - 2 \dots (i)$$

$$\therefore \frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c}$$

$$\begin{aligned}
&= \frac{3 - 2(a+b+c) + ab + bc + ca}{1 - (a+b+c) + ab + bc + ca - abc} \\
&= \frac{3 - 2(a+b+c) + ab + bc + ca}{1 - (a+b+c) + ab + bc + ca - a - b - c + 2} \\
&= \frac{3 - 2(a+b+c) + ab + bc + ca}{3 - 2(a+b+c) + ab + bc + ca} = 1
\end{aligned}$$

161 (c)

Let projection be  $x$ , then

$$\vec{a} = \frac{x(\hat{i} + \hat{j})}{\sqrt{2}} + \frac{x(-\hat{i} + \hat{j})}{\sqrt{2}} + x\hat{k}$$

$$\therefore \vec{a} = \frac{2x\hat{j}}{\sqrt{2}} + x\hat{k}$$

$$\Rightarrow \vec{a} = \frac{\sqrt{2}}{\sqrt{3}}\hat{j} + \frac{\hat{k}}{\sqrt{3}}$$

162 (a)

$$\vec{PQ} = 6\hat{i} + \hat{j}$$

$$\vec{QR} = -\hat{i} + 3\hat{j}$$

$$\vec{RS} = -6\hat{i} - \hat{j}$$

$$\vec{SP} = \hat{i} - 3\hat{j}$$

$$|\vec{PQ}| = \sqrt{37} = |\vec{RS}|$$

$$|\vec{QR}| = \sqrt{10} = |\vec{SP}|$$

$$\vec{PQ} \cdot \vec{QR} = -6 + 3 = -3 \neq 0$$

$\vec{PQ}$  is not parallel to  $\vec{RS}$  and their magnitude are equal.

$\Rightarrow$  Quadrilateral  $PQRS$  must be a parallelogram, which is neither a rhombus nor a rectangle.

163 (c)

If  $\Delta = 0$ , then

$$\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{a} \cdot \vec{c} & \vec{b} \cdot \vec{c} & \vec{c} \cdot \vec{c} \end{vmatrix} = 0$$

$$\Rightarrow \lambda\vec{a} + \mu\vec{b} + \nu\vec{c} = 0$$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$  are L.D., which is a contradiction

Hence,  $\Delta$  can take any non-zero real values

164 (b)

We have,

$$(3\vec{a} - 2\vec{b}) = -8\hat{i} - 7\hat{j} + 3\hat{k} \text{ and } \vec{c} = \frac{1}{3}(2\hat{i} + 2\hat{j} - \hat{k})$$

$$\therefore \text{Required projection} = (3\vec{a} - 2\vec{b}) \cdot \vec{c}$$

$$= (-8\hat{i} - 7\hat{j} + 3\hat{k}) \cdot \frac{1}{3}(2\hat{i} + 2\hat{j} - \hat{k})$$

$$= \frac{1}{3}(-16 - 14 - 3) = -11$$

165 (a)

Angle between the faces  $OAB$  and  $ABC$  is same as angle between normals of faces  $OAB$  and  $ABC$ .

Vector along the normals of  $OAB$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 1 \\ 2 & 1 & 3 \end{vmatrix} = 5\hat{i} - \hat{j} - 3\hat{k} = \vec{a} \text{ (let)}$$

Vector along normals of  $ABC$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ -2 & -1 & 1 \end{vmatrix} = \hat{i} - 5\hat{j} - 3\hat{k} = \vec{b} \text{ (let)}$$

$$\therefore \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{5 + 5 + 9}{\sqrt{35} \sqrt{35}}$$

$$\Rightarrow \theta = \cos^{-1} \left( \frac{19}{35} \right)$$

167 (d)

$$\vec{a} \times [\vec{a} \times (\vec{a} \times \vec{b})] = \vec{a} \times \{ (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} \}$$

(Expanding by vector triple product)

$$= (\vec{a} \cdot \vec{b})(\vec{a} \times \vec{a}) - (\vec{a} \cdot \vec{a})(\vec{a} \times \vec{b})$$

$$= (\vec{a} \cdot \vec{a})(\vec{b} \times \vec{a}) \quad (\because (\vec{a} \times \vec{a}) = 0)$$

169 (b)

Taking  $A$  as the origin let the position vectors of  $B$

and  $C$  be  $\vec{b}$  and  $\vec{c}$  respectively

Equations of lines  $BF$  and  $AC$  are

$$\vec{r} = \vec{b} + \lambda \left( \frac{\vec{b} + \vec{c}}{4} - \vec{b} \right) \text{ and } \vec{r} = \vec{0} + \mu \vec{c} \text{ respectively}$$

For the point of intersection  $F$ , we have

$$\vec{b} + \lambda \left( \frac{\vec{c} - 3\vec{b}}{4} \right) = \mu \vec{c}$$

$$\Rightarrow 1 - \frac{3\lambda}{4} = 0 \text{ and } \frac{\lambda}{4} = \mu \Rightarrow \lambda = \frac{4}{3} \text{ and } \mu = \frac{1}{3}$$

So, the position vector of  $F$  is  $\vec{r} = \frac{1}{3}\vec{c}$

$$\text{Now, } \vec{AF} = \frac{1}{3}\vec{c} \Rightarrow \vec{AF} = \frac{1}{3}\vec{AC}$$

$$\text{Hence, } AF:AC = \frac{1}{3}:1 = \frac{1}{3}$$

170 (d)

$$\text{Given, } |\vec{a}| = 1, |\vec{b}| = 2$$

$$\therefore [(\vec{a} + 3\vec{b}) \times (3\vec{a} + \vec{b})]^2$$

$$= [0 + \vec{a} \times \vec{b} + 9\vec{b} \times \vec{a} + 0]^2$$

$$= [-8\vec{a} \times \vec{b}]^2$$

$$= 64[|\vec{a}|^2 |\vec{b}|^2 \sin^2 \theta]$$

$$= 64[1 \times 4 \times \sin^2 120^\circ]$$

$$= 64 \times 4 \times \frac{3}{4} = 192$$

171 (c)

$$(\vec{a} + \vec{b} + \vec{c}) \cdot [(\vec{a} + \vec{b}) \times (\vec{a} + \vec{c})]$$

$$= (\vec{a} + \vec{b} + \vec{c}) \cdot [\vec{a} \times \vec{c} + \vec{b} \times \vec{a} + \vec{b} \times \vec{c}]$$

$$0 + 0 + [\vec{a} \cdot \vec{b} \cdot \vec{c}] + [\vec{b} \cdot \vec{a} \cdot \vec{c}] + 0 + 0 + 0 + [\vec{c} \cdot \vec{b} \cdot \vec{a}] + 0$$

$$= -[\vec{a} \cdot \vec{b} \cdot \vec{c}]$$

172 (b)

$$\text{Clearly, } \vec{c} = \pm \frac{\vec{a} \times (\vec{a} \times \vec{b})}{|\vec{a} \times (\vec{a} \times \vec{b})|}$$

Now,

$$\vec{a} \times (\vec{a} \times \vec{b}) = (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}$$

$$\Rightarrow \vec{a} \times (\vec{a} \times \vec{b}) = -\hat{i} - \hat{j} + \hat{k} - 3(\hat{i} - \hat{j} + \hat{k})$$

$$= -4\hat{i} + 2\hat{j} - 2\hat{k}$$

$$\therefore \vec{c} = \pm \frac{1}{\sqrt{6}}(2\hat{i} - \hat{j} + \hat{k})$$

Since  $\vec{d}$  is a unit vector perpendicular to both  $\vec{a}$  and  $\vec{c}$

$$\therefore \vec{d} = \pm \frac{\vec{a} \times \vec{c}}{|\vec{a} \times \vec{c}|}$$

$$\text{Now, } \vec{a} \times \vec{c} = \pm \frac{1}{\sqrt{6}} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & -1 \\ 2 & -1 & 1 \end{vmatrix}$$

$$= \pm \frac{1}{\sqrt{6}}(-3\hat{j} - 3\hat{k})$$

$$\therefore \vec{d} = \pm \frac{1}{\sqrt{2}}(-\hat{j} - \hat{k}) = \pm \frac{1}{\sqrt{2}}(\hat{j} + \hat{k})$$

173 (d)

Since,  $G$  is the centroid of a triangle, then

$$\vec{GA} + \vec{GB} + \vec{GC} = \vec{0} \Rightarrow \vec{GA} + \vec{GC} = -\vec{GB} \dots (i)$$

$$\text{Now, } \vec{GA} + \vec{BG} + \vec{GC} = -\vec{GB} + \vec{BG} = 2\vec{BG}$$

[ from Eq. (i) ]

174 (c)

Let  $\vec{n}_1$  and  $\vec{n}_2$  be the vectors normal to the plane determined by  $\hat{i}, \hat{i} - \hat{j}$  and  $\hat{i} + \hat{j}, \hat{i} - \hat{k}$  respectively

$$\therefore \vec{n}_1 = \hat{i} \times (\hat{i} - \hat{j}) = -\hat{k}$$

$$\text{and } \vec{n}_2 = (\hat{i} + \hat{j}) \times (\hat{i} - \hat{k}) = -\hat{i} + \hat{j} - \hat{k}$$

Since,  $\vec{a}$  is parallel to the line of intersection of the planes determined by the given planes.

$$\therefore \vec{a} \parallel (\vec{n}_1 \times \vec{n}_2)$$

$$\Rightarrow \vec{a} = \lambda(\vec{n}_1 \times \vec{n}_2) = \lambda(\hat{i} + \hat{j})$$

Let  $\theta$  be the angle between  $\vec{a}$  and  $\hat{i} + 2\hat{j} - 2\hat{k}$

$$\therefore \cos \theta = \frac{\lambda((\hat{i} + \hat{j}) \cdot (\hat{i} + 2\hat{j} - 2\hat{k}))}{\sqrt{\lambda^2 + \lambda^2} \sqrt{1 + 4 + 4}}$$

$$= \frac{\lambda(1 + 2)}{\sqrt{2}\lambda \times 3} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \theta = \frac{\pi}{4}$$

175 (d)

$$|\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$$

$$\Rightarrow |\vec{a} \times \vec{b}|^2 = 25 \times 36 - (25)^2$$

$$= 25(36 - 25)$$

$$= 25 \times 11$$

$$\Rightarrow |\vec{a} \times \vec{b}| = 5\sqrt{11}$$


176 (c)

$$\vec{AB} + \vec{AE} + \vec{BC} + \vec{DC} + \vec{ED} + \vec{AC}$$

$$= (\vec{AB} + \vec{BC}) + (\vec{AE} + \vec{ED}) + \vec{DC} + \vec{AC}$$

$$= \vec{AC} + (\vec{AD} + \vec{DC}) + \vec{AC}$$

$$= \vec{AC} + \vec{AC} + \vec{AC} = 3\vec{AC}$$



177 (a)

We have,

$$(\vec{a} - \vec{d}) \times (\vec{b} - \vec{c})$$

$$= \vec{a} \times \vec{b} - \vec{a} \times \vec{c} - \vec{d} \times \vec{b} + \vec{d} \times \vec{c}$$

$$\Rightarrow (\vec{a} - \vec{d}) \times (\vec{b} - \vec{c})$$

$$= \vec{c} \times \vec{d} - \vec{b} \times \vec{d} - \vec{d} \times \vec{b} + \vec{d} \times \vec{c}$$

$$\Rightarrow (\vec{a} - \vec{d}) \times (\vec{b} - \vec{c})$$

$$= 0 \quad [\because \vec{a} \times \vec{b} = \vec{c} \times \vec{d}, \vec{a} \times \vec{c}$$

$$= \vec{b} \times \vec{d}]$$

$$\Rightarrow (\vec{a} - \vec{d}) \parallel (\vec{b} - \vec{c})$$

$$\Rightarrow \vec{a} - \vec{d} = \lambda(\vec{b} - \vec{c})$$

Similarly, we have

$$(\vec{a} + \vec{d}) \times (\vec{b} + \vec{c}) = \vec{0} \Rightarrow \vec{a} + \vec{d} \parallel \vec{b} + \vec{c} \Rightarrow \vec{a} + \vec{d}$$

$$= \lambda(\vec{b} + \vec{c})$$

178 (b)

We have,

$$\vec{a} \times \{\vec{a} \times (\vec{a} \times \vec{b})\} = \vec{a} \times \{(\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}\}$$

$$\Rightarrow \vec{a} \times \{\vec{a} \times (\vec{a} \times \vec{b})\} = -(\vec{a} \cdot \vec{a})(\vec{a} \times \vec{b})$$

$$= (\vec{a} \cdot \vec{a})(\vec{b} \times \vec{a})$$

179 (c)


We have,

$$\vec{AB} + \vec{DC} = \vec{AB} + \vec{BC} - \vec{BC} + \vec{DC}$$

$$\Rightarrow \vec{AB} + \vec{DC} = (\vec{AB} + \vec{BC}) - \vec{BC} + \vec{CD}$$

$$\Rightarrow \vec{AB} + \vec{DC} = (\vec{AB} + \vec{BC}) - (\vec{BC} + \vec{CD})$$

$$\Rightarrow \vec{AB} + \vec{DC} = \vec{AC} - \vec{BD} = \vec{AC} + \vec{DB}$$



182 (c)

Volume of parallelepiped,

$$f(a) = \begin{vmatrix} 1 & a & 1 \\ 0 & 1 & a \\ a & 0 & 1 \end{vmatrix} = 1 + a^3 - a$$

$$\text{Now, } f'(a) = 3a^2 - 1$$

$$\Rightarrow f''(a) = 6a$$

$$\text{Put } f'(a) = 0$$

$$\Rightarrow a \neq \pm \frac{1}{\sqrt{3}}$$

Which shows  $f(a)$  is maximum at

$$a = \frac{1}{\sqrt{3}} \text{ and maximum at}$$

$$a = -\frac{1}{\sqrt{3}}$$

183 (c)

$$\text{Let } \vec{a} = 4\hat{i} + 6\hat{j} - \hat{k}$$

$$\text{and } \vec{b} = 3\hat{i} + 8\hat{j} + \hat{k}$$

$$\therefore \vec{c} = \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 6 & -1 \\ 3 & 8 & 1 \end{vmatrix} = 14\hat{i} - 7\hat{j} + 14\hat{k}$$

$$\Rightarrow \hat{c} = \frac{14\hat{i} - 7\hat{j} + 14\hat{k}}{\sqrt{14^2 + 7^2 + 14^2}} = \frac{14\hat{i} - 7\hat{j} + 14\hat{k}}{21}$$

$\therefore$  Required vector

$$= 12 \cdot \frac{(14\hat{i} - 7\hat{j} + 14\hat{k})}{21} = 8\hat{i} - 4\hat{j} + 8\hat{k}$$

184 (b)

$$\text{Since, } \vec{a} \cdot \vec{b} = 0$$

$$\text{Also, } \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \cos \theta$$

$$\text{Now, } \vec{c} = \alpha\vec{a} + \beta\vec{b} + \gamma(\vec{a} \cdot \vec{b})$$

$$\vec{a} \cdot \vec{c} = \alpha\vec{a} \cdot \vec{a} + \beta\vec{a} \cdot \vec{b} + \gamma\vec{a} \cdot (\vec{a} \cdot \vec{b})$$

$$\Rightarrow |\vec{a}||\vec{c}| \cos \theta = \alpha + 0 + 0$$

$$\Rightarrow \cos \theta = \alpha \quad [\because \vec{a} \cdot \vec{b} = 0]$$

$$\text{and } \vec{b} \cdot \vec{c} = \alpha\vec{a} \cdot \vec{b} + \beta\vec{b} \cdot \vec{b} + \gamma(\vec{a} \cdot \vec{b}) \cdot \vec{b}$$

$$\Rightarrow |\vec{b}||\vec{c}| \cos \theta = \beta \Rightarrow \cos \theta = \beta$$

185 (a)

Given volume of parallelepiped

$$[\vec{a} \vec{b} \vec{c}] = 40$$

$\therefore$  Volume of parallelepiped

$$= [\vec{b} + \vec{c} \vec{c} + \vec{a} \vec{a} + \vec{b}] = 2[\vec{a} \vec{b} \vec{c}]$$

$$= 2 \times 40 = 80 \text{ cu units}$$

186 (a)

$$\text{Given, } \vec{OP} = \hat{a} \cos t + \hat{b} \sin t$$

$$\Rightarrow |\vec{OP}|$$

$$= \sqrt{(\hat{a} \cdot \hat{a} \cos^2 t + \hat{b} \cdot \hat{b} \sin^2 t + 2\hat{a} \cdot \hat{b} \sin t \cos t)}$$

$$\Rightarrow |\vec{OP}| = \sqrt{1 + \hat{a} \cdot \hat{b} \sin 2t}$$

$$\Rightarrow |\vec{OP}|_{\text{max}} = \sqrt{1 + \hat{a} \cdot \hat{b}}$$

$$[\text{Max}(\sin 2t) = 1 \Rightarrow t = \frac{\pi}{4}]$$

$$\Rightarrow \vec{OP} \left( \text{at } t = \frac{\pi}{4} \right) = \frac{1}{\sqrt{2}}(\hat{a} + \hat{b})$$

$$\therefore \text{Unit vector along } \overrightarrow{OP} \text{ at } \left(t = \frac{\pi}{4}\right) = \frac{\hat{a} + \hat{b}}{|\hat{a} + \hat{b}|}$$

187 (b)

The position vector of midpoint of line joining the points whose position vector are  $\hat{i} + \hat{j} - \hat{k}$  and  $\hat{i} - \hat{j} + \hat{k}$

$$= \frac{\hat{i} + \hat{j} - \hat{k} + \hat{i} - \hat{j} + \hat{k}}{2} = \hat{i}$$

188 (a)

The position vector of  $G$  is  $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$

$$\therefore \vec{GA} + \vec{GB} + \vec{GC}$$

$$= \left(\vec{a} - \frac{\vec{a} + \vec{b} + \vec{c}}{3}\right) + \left(\vec{b} - \frac{\vec{a} + \vec{b} + \vec{c}}{3}\right) + \left(\vec{c} - \frac{\vec{a} + \vec{b} + \vec{c}}{3}\right) = \vec{0}$$

189 (d)

A vector normal to first plane is  $\vec{n}_1 = \hat{i} \times (\hat{i} + \hat{j}) = \hat{k}$

A vector normal to second plane is  $\vec{n}_2 = (\hat{i} - \hat{j}) \times (\hat{i} + \hat{k}) = -\hat{j} + \hat{k} - \hat{i}$

Since,  $\vec{a}$  will be parallel to  $\vec{n}_1 \times \vec{n}_2 = \hat{i} - \hat{j}$

Let  $\theta$  be the angle between  $\vec{a}$  and  $\hat{i} - 2\hat{j} + 2\hat{k}$

$$\begin{aligned} \therefore \cos \theta &= \frac{(\hat{i} - \hat{j}) \cdot (\hat{i} - 2\hat{j} + 2\hat{k})}{\sqrt{1^2 + 1^2} \sqrt{1^2 + 2^2 + 2^2}} \\ &= \frac{1 + 2}{\sqrt{2} \cdot 3} = \frac{1}{\sqrt{2}} \\ \Rightarrow \theta &= \frac{\pi}{4} \end{aligned}$$

190 (a)

Since, given planes are perpendicular, it means its normal are perpendicular.

$$\therefore 2(\lambda) - \lambda(5) + 3(-1) = 0$$

$$\Rightarrow -3\lambda - 3 = 0$$

$$\Rightarrow \lambda = -1$$

$$\therefore \lambda^2 + \lambda = (-1)^2 - 1 = 0$$

191 (a)

$$2\overrightarrow{OA} + 3\overrightarrow{OB} = 2(\overrightarrow{OC} + \overrightarrow{CA}) + 3(\overrightarrow{OC} + \overrightarrow{CB})$$

$$= 5\overrightarrow{OC} + 2\overrightarrow{CA} + 3\overrightarrow{CB}$$

$$= 5\overrightarrow{OC} \quad [\because 2\overrightarrow{CA} = -3\overrightarrow{CB}]$$

192 (b)

If the vectors  $(\sec^2 A)\hat{i} + \hat{j} + \hat{k}$ ,  $\hat{i} + (\sec^2 B)\hat{j} + \hat{k}$  and  $\hat{i} + \hat{j} + (\sec^2 C)\hat{k}$  are coplanar, then

$$\begin{vmatrix} \sec^2 A & 1 & 1 \\ 1 & \sec^2 B & 1 \\ 1 & 1 & \sec^2 C \end{vmatrix} = 0$$

$$\Rightarrow \sec^2 A \sec^2 B \sec^2 C - \sec^2 A - \sec^2 B - \sec^2 C + 2 = 0$$

$$\Rightarrow (1 + \tan^2 A)(1 + \tan^2 B)(1 + \tan^2 C) - (1 + \tan^2 A)$$

$$- (1 + \tan^2 B) - (1 + \tan^2 C) + 2 = 0$$

$$\Rightarrow \tan^2 A \tan^2 B \tan^2 C + \tan^2 A \tan^2 B$$

$$+ \tan^2 B \tan^2 C + \tan^2 C \tan^2 A$$

$$= 0$$

$$\Rightarrow \cot^2 A + \cot^2 B + \cot^2 C + 1 = 0$$

$$\Rightarrow \operatorname{cosec}^2 A + \operatorname{cosec}^2 B + \operatorname{cosec}^2 C - 2 = 0$$

$$\Rightarrow \operatorname{cosec}^2 A + \operatorname{cosec}^2 B + \operatorname{cosec}^2 C = 2$$

193 (b)

It is given that the points  $P, Q$  and  $R$  with position vectors  $2\hat{i} + \hat{j} + \hat{k}$ ,  $6\hat{i} - \hat{j} + 2\hat{k}$  and  $14\hat{i} - 5\hat{j} + p\hat{k}$  respectively are collinear

$$\therefore \vec{PQ} = \lambda \vec{QR} \text{ for some scalar } \lambda$$

$$\Rightarrow 4\hat{i} - 2\hat{j} + \hat{k} = \lambda\{8\hat{i} - 4\hat{j} + (p-2)\hat{k}\}$$

$$\Rightarrow 4 = 8\lambda, -2 = -4\lambda \text{ and } \lambda(p-2) = 1 \Rightarrow p = 4$$

194 (c)

$$\text{Given, } \vec{\alpha} + \vec{\beta} + \vec{\gamma} = a\vec{\delta} \quad \dots(i)$$

$$\vec{\beta} + \vec{\gamma} + \vec{\delta} = b\vec{\alpha} \quad \dots(ii)$$

From Eq. (i)

$$\vec{\alpha} + \vec{\beta} + \vec{\gamma} + \vec{\delta} = (a+1)\vec{\delta} \quad \dots(iii)$$

From Eq. (ii)

$$\vec{\alpha} + \vec{\beta} + \vec{\gamma} + \vec{\delta} = (b+1)\vec{\alpha} \quad \dots(iv)$$

From Eq. (iii) and (iv),

$$(a+1)\vec{\delta} = (b+1)\vec{\alpha} \quad \dots(v)$$

Since,  $\vec{\alpha}$  is not parallel to  $\vec{\delta}$ .

$$\therefore \text{From Eq. (v),}$$

$$a+1 = 0 \text{ and } b+1 = 0$$

$$\therefore \text{From Eq. (iii),}$$

$$\vec{\alpha} + \vec{\beta} + \vec{\gamma} + \vec{\delta} = \vec{0}$$

196 (d)

We have,

$$\begin{aligned} [2\vec{a} + \vec{b} \quad 2\vec{b} + \vec{c} \quad 2\vec{c} + \vec{a}] &= \begin{vmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 1 & 0 & 2 \end{vmatrix} [\vec{a} \quad \vec{b} \quad \vec{c}] \\ &= 9 \times 3 = 27 \end{aligned}$$

Hence, required volume = 27 cubic units

197 (a)

In plane  $P_1$ , a vector is perpendicular to  $\vec{a}$  and  $\vec{b}$  is  $\vec{a} \times \vec{b}$ .

In plane  $P_2$ , a vector is perpendicular to  $\vec{c}$  and  $\vec{d}$  is  $\vec{c} \times \vec{d}$

$$\Rightarrow (\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = 0$$

$$\Rightarrow (\vec{a} \times \vec{b}) \parallel (\vec{c} \times \vec{d})$$

The angle between the planes is 0.

198 (a)

We have,

$$\frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{(\vec{c} \times \vec{a}) \cdot \vec{b}} + \frac{\vec{b} \cdot (\vec{a} \times \vec{c})}{\vec{c} \cdot (\vec{a} \times \vec{b})} = \frac{[\vec{a} \vec{b} \vec{c}]}{[\vec{b} \vec{c} \vec{a}]} + \frac{[\vec{b} \vec{a} \vec{c}]}{[\vec{c} \vec{a} \vec{b}]}$$

$$= \frac{[\vec{a} \vec{b} \vec{c}]}{[\vec{a} \vec{b} \vec{c}]} - \frac{[\vec{a} \vec{b} \vec{c}]}{[\vec{a} \vec{b} \vec{c}]} = 1 - 1 = 0$$

199 (a)

Given,  $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j}$ ,

Now,  $|\vec{a}| = \sqrt{4 + 1 + 4} = 3$

Since,  $|\vec{c} - \vec{a}| = 2\sqrt{2}$

$$\Rightarrow |\vec{c}|^2 + |\vec{a}|^2 - 2\vec{c} \cdot \vec{a} = 8$$

$$\Rightarrow |\vec{c}|^2 + 9 - 2|\vec{c}| = 8$$

$$\Rightarrow |\vec{c}|^2 - 2|\vec{c}| + 1 = 0$$

$$\Rightarrow |\vec{c}| = 1$$

Now,  $|(\vec{a} \times \vec{b}) \times \vec{c}| = |\vec{a} \times \vec{b}| |\vec{c}| \sin 30^\circ \dots (i)$

Now,  $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 1 & 1 & 0 \end{vmatrix}$

$$= \hat{i}(0 + 2) - \hat{j}(0 + 2) + \hat{k}(2 - 1)$$

$$= 2\hat{i} - 2\hat{j} + \hat{k}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = \sqrt{4 + 4 + 1} = 3$$

$\therefore$  From Eq. (i),

$$|(\vec{a} \times \vec{b}) \times \vec{c}| = 3 \cdot 1 \cdot \frac{1}{2} = \frac{3}{2}$$

201 (a)

Now,  $\vec{AB} + 2\vec{AD} + \vec{BC} - 2\vec{DC}$

$$= \vec{AC} + 2\vec{AD} - 2\vec{DC}$$

$$= \vec{AC} + 2(\vec{AC} + \vec{CD}) - 2\vec{DC}$$


$$= 3\vec{AC} - 4\vec{DC}$$

$$= 3(2\vec{QC}) - 4\left(\frac{3}{2}\vec{PC}\right)$$

$$= 6\vec{QC} - 6\vec{PC} = 6(\vec{QC} + \vec{CP})$$

$$\Rightarrow k\vec{PQ} = 6\vec{QP} = -6\vec{PQ} \text{ (given)}$$

$$\Rightarrow k = -6$$



202 (c)

Given,  $|\vec{a} \times \vec{b}| = 4 \Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 4 \dots (i)$

$$\Rightarrow \sin \theta = \frac{4}{|\vec{a}| |\vec{b}|}$$

Alos,  $|\vec{a} \cdot \vec{b}| = 2 \Rightarrow |\vec{a}| |\vec{b}| \cos \theta = 2$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 (1 - \sin^2 \theta) = 4$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 \left(1 - \frac{16}{|\vec{a}|^2 |\vec{b}|^2}\right) = 4 \text{ [From Eq. 1]}$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 = 20$$

203 (c)

We have,

$$\vec{p} = \frac{\vec{b} \times \vec{c}}{[\vec{a} \vec{b} \vec{c}]}, \vec{q} = \frac{\vec{c} \times \vec{a}}{[\vec{a} \vec{b} \vec{c}]}, \vec{r} = \frac{\vec{a} \times \vec{b}}{[\vec{a} \vec{b} \vec{c}]}$$

$$\therefore \vec{a} \times \vec{p} = \frac{1}{[\vec{a} \vec{b} \vec{c}]} \vec{a} \times (\vec{b} \times \vec{c}),$$

$$\vec{b} \times \vec{q} = \frac{1}{[\vec{a} \vec{b} \vec{c}]} \vec{b} \times (\vec{c} \times \vec{a})$$

$$\vec{c} \times \vec{r} = \frac{1}{[\vec{a} \vec{b} \vec{c}]} \vec{c} \times (\vec{a} \times \vec{b})$$

$$\therefore \vec{a} \times \vec{p} + \vec{b} \times \vec{q} + \vec{c} \times \vec{r}$$

$$= \frac{1}{[\vec{a} \vec{b} \vec{c}]} \{ \vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b}) \}$$

$$= \vec{0}$$

204 (b)

Since,  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

$$= \frac{c(\log_2 x)^2 - 12 + 6c \log_2 x}{\sqrt{(c \log_2 x)^2 + 36 + 9} \sqrt{(\log_2 x)^2 + 4 + 4(c \log_2 x)^2}}$$

For obtuse angle,

$$\cos \theta < 0$$

$$\Rightarrow c(\log_2 x)^2 - 12 + 6c \log_2 x < 0$$

$$\Rightarrow c < 0 \text{ and } D < 0$$

$$\Rightarrow c < 0 \text{ and } (6c)^2 + 48c < 0$$

$$\Rightarrow c < 0 \text{ and } c < -\frac{4}{3}$$

$$\therefore c \in \left(-\frac{4}{3}, 0\right)$$

206 (d)

Given lines can be rewritten as

$$\vec{r} = 2\hat{i} + \hat{j} + 2\hat{k} + t(-3\hat{i} + 2\hat{j} + 6\hat{k})$$

$$\text{and } \vec{r} = \hat{i} + 2\hat{j} - \hat{k} + s(4\hat{i} - \hat{j} + 8\hat{k})$$

here,  $a_1 = -3, b_1 = 2, c_1 = 6$

and  $a_2 = 4, b_2 = -1, c_2 = 8$

$$\therefore \cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

$$= \frac{-3 \times 4 + 2 \times (-1) + 6 \times 8}{\sqrt{9 + 4 + 36} \sqrt{16 + 1 + 64}} = \frac{34}{7 \times 9}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{34}{63}\right)$$

208 (a)

We have,

$$\vec{AB} = \hat{i} - 7\hat{j} + \hat{k} \text{ and } \vec{BC} = 3\hat{i} + \hat{j} + 2\hat{k}$$

$$\therefore \vec{AC} = \vec{AB} + \vec{BC} = 4\hat{i} - 6\hat{j} + 3\hat{k}$$

$$\Rightarrow |\vec{AC}| = \sqrt{16 + 36 + 9} = \sqrt{61}$$

210 (d)

$$(\hat{i} + \hat{j} + 2\hat{k}) \cdot \left( \frac{m\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{13 + m^2}} \right) = 2$$

$$\Rightarrow m + 2 + 6 = 2\sqrt{13 + m^2}$$

$$\Rightarrow (m + 8)^2 = 4(13 + m^2)$$

$$\Rightarrow m^2 + 16m + 64 = 4m^2 + 52$$

$$\Rightarrow 3m^2 - 16m - 12 = 0$$

$$\Rightarrow (3m + 2)(m - 6) = 0$$

$$\Rightarrow m = 6, -\frac{2}{3}$$

211 (c)

If  $\vec{a}$  and  $\vec{b}$  are non-zero and non-collinear vectors and there exists  $\alpha$  and  $\beta$  such that  $\alpha\vec{a} + \beta\vec{b} = \vec{0}$ , then  $\alpha = \beta = 0$

212 (d)

Given vectors are coplanar, if

$$\begin{vmatrix} 1 & 1 & m \\ 1 & 1 & m+1 \\ 1 & -1 & m \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 0 & 0 & -1 \\ 1 & 1 & m+1 \\ 1 & -1 & m \end{vmatrix} = 0 \quad [R_1 \rightarrow R_1 - R_2]$$

$$\Rightarrow -1(-1 - 1) = 0$$

$$\Rightarrow 2 \neq 0$$

$\therefore$  Now value of  $m$  for which vectors are coplanar.

213 (b)

Let the required unit vector  $\vec{c} = x\hat{i} + y\hat{k}$

We have,

$$|\vec{c}| = 1 \Rightarrow x^2 + y^2 = 1 \quad \dots(i)$$

Vectors  $\vec{a}$  and  $\vec{c}$  are inclined at an angle of  $45^\circ$

$$\therefore \cos 45^\circ = \frac{2x-y}{\sqrt{4+4+1}} \Rightarrow 2x - y = \frac{3}{\sqrt{2}} \quad \dots(ii)$$

Vectors  $\vec{b}$  and  $\vec{c}$  are inclined at an angle of  $60^\circ$

$$\therefore -\frac{y}{\sqrt{2}} = \cos 60^\circ \Rightarrow y = -\frac{1}{\sqrt{2}} \quad \dots(iii)$$

From (ii) and (iii), we get  $x = 1/\sqrt{2}$

Hence, the required unit vector is  $\frac{1}{\sqrt{2}}\hat{i} - \frac{1}{\sqrt{2}}\hat{k}$

214 (c)

Let  $\vec{A} = 2\hat{i} + 3\hat{j} - \hat{k}$ ,  $\vec{B} = \hat{i} - 2\hat{j} + 3\hat{k}$ ,

$\vec{C} = 3\hat{i} + 4\hat{j} - 2\hat{k}$  and  $\vec{D} = \hat{i} - 6\hat{j} + \lambda\hat{k}$

Now,  $\vec{AB} = -\hat{i} - 5\hat{j} + 4\hat{k}$ ,  $\vec{AC} = \hat{i} + \hat{j} - \hat{k}$

and  $\vec{AD} = -\hat{i} - 9\hat{j} + (\lambda + 1)\hat{k}$

These will be coplanar, if  $[\vec{AB} \ \vec{AC} \ \vec{AD}] = 0$

$$\therefore \begin{vmatrix} -1 & -5 & 4 \\ 1 & 1 & -1 \\ -1 & -9 & (\lambda + 1) \end{vmatrix} = 0$$

$$\Rightarrow -1(\lambda + 1 - 9) + 5(\lambda + 1 - 1) + 4(-9 + 1) = 0$$

$$\Rightarrow \lambda = 6$$

215 (b)

We have,

$$|\vec{a}| = |\vec{b}|$$

Now,

$$\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = |\vec{a}|^2 - |\vec{b}|^2$$

$$\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 0 \quad [\because |\vec{a}| = |\vec{b}|]$$

$$\Rightarrow (\vec{a} + \vec{b}) \perp (\vec{a} - \vec{b})$$

216 (a)

Adjacent sides of parallelogram are  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{b} = -3\hat{i} - 2\hat{j} + \hat{k}$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ -3 & -2 & 1 \end{vmatrix}$$

$$= \hat{i}(2 + 6) - \hat{j}(1 + 9) + \hat{k}(-2 + 6)$$

$$= 8\hat{i} - 10\hat{j} + 4\hat{k}$$

Therefore, area of parallelogram


$$= |\vec{a} \times \vec{b}|$$

$$= \sqrt{(8)^2 + (-10)^2 + (4)^2}$$

$$= \sqrt{64 + 100 + 16} = \sqrt{180} \text{ sq unit}$$

217 (d)

$$\therefore \vec{CP} + \vec{PA} + \vec{BA}$$



By triangle law,

$$\vec{CA} = \vec{CB} + \vec{BA}$$

$$\therefore \vec{CP} + \vec{PA} = \vec{CB} + \vec{BA}$$

218 (d)

We have,

$$\vec{c} = x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})$$

$$\Rightarrow \vec{c} \cdot \vec{a} = x \text{ and } \vec{c} \cdot \vec{b} = y \Rightarrow x = y = \cos \theta$$

Now,

$$\vec{c} \cdot \vec{c} = |\vec{c}|^2$$

$$\Rightarrow \{x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})\} \cdot \{x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})\} = |\vec{c}|^2$$

$$\Rightarrow 2x^2 + x^2|\vec{a} \times \vec{b}|^2 = 1$$

$$\Rightarrow 2x^2 + z^2\{|\vec{a}|^2|\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2\} = 1$$

$$\Rightarrow 2x^2 + z^2 = 1 \quad [\because |\vec{a}|^2 = 1, |\vec{b}|^2 = 1 \text{ and } \vec{a} \cdot \vec{b} = 0]$$

$$\Rightarrow z^2 = 1 - 2\cos^2 \theta = -\cos 2\theta$$

219 (a)

We have,

$$\vec{a} + \vec{b} = \vec{c}$$

$$\Rightarrow |\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow |\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2 \times 0 \quad [\because \vec{a} \cdot \vec{b} = 0]$$

$$\Rightarrow |\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2$$

221 (b)

All points  $A, B, C, D, E$  are in a plane.

$$\therefore \text{Resultant} = (\vec{AC} + \vec{AD} + \vec{AE}) + (\vec{CB} + \vec{DB} + \vec{EB})$$

$$= (\vec{AC} + \vec{CB}) + (\vec{AD} + \vec{DB}) + (\vec{AE} + \vec{EB}) \\ = \vec{AB} + \vec{AB} + \vec{AB} = 3\vec{AB}$$

222 (a)

Since,  $\vec{a}, \vec{b}, \vec{c}$  are coplanar.

$$\Rightarrow \begin{vmatrix} \alpha & 2 & \beta \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow \alpha(1-0) - 2(1-0) + \beta(1-0) = 0$$

$$\Rightarrow \alpha + \beta = 2 \text{ Which is possible for } \alpha = 1, \beta = 1$$

223 (c)

A unit perpendicular to the plane  $\vec{a}$  and  $\vec{b} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -6 & -3 \\ 4 & 3 & -1 \end{vmatrix}$$

$$= \hat{i}(6+9) - \hat{j}(-2+12) + \hat{k}(6+24)$$

$$= 15\hat{i} - 10\hat{j} + 30\hat{k}$$

$$\text{and } |\vec{a} \times \vec{b}| = \sqrt{15^2 + (-10)^2 + (30)^2}$$

$$= \sqrt{1225} = 35$$

$$\therefore \text{Required vector} = \frac{15\hat{i} - 10\hat{j} + 30\hat{k}}{35} = \frac{3\hat{i} - 2\hat{j} + 6\hat{k}}{7}$$

225 (d)

$$(\vec{a} \times \hat{j}) \cdot (2\hat{j} - 3\hat{k}) = \vec{a} \cdot \{\hat{j} \times (2\hat{j} - 3\hat{k})\}$$

$$= \vec{a} \cdot \{-3(\hat{j} \times \hat{k})\} = -3(\vec{a} \cdot \hat{i})$$

$$= -12 \quad [\because \vec{a} \cdot \hat{i} = 4, \text{ given}]$$

226 (b)

Volume of tetrahedron

$$= \frac{1}{6} [\vec{AB} \vec{AC} \vec{AD}]$$

$$= \frac{1}{6} \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 2 & -1 \end{vmatrix}$$

$$= \frac{1}{6} [-1 + 2 + 3] = \frac{2}{3} \text{ cu unit}$$

228 (c)

$$\text{Since, } \vec{a} \times (\vec{b} \times \vec{c}) = \frac{1}{2} \vec{b}$$

$$\Rightarrow (\vec{a} \cdot \vec{b})\vec{b} - (\vec{a} \cdot \vec{c})\vec{c} = \frac{1}{2} \vec{b}$$

On comparing both sides, we get

$$\vec{a} \cdot \vec{c} = \frac{1}{2} \text{ and } \vec{a} \cdot \vec{b} = 0$$

$$\text{Now, } \vec{a} \cdot \vec{b} = 0$$

$$\Rightarrow |\vec{a}| |\vec{c}| \cos \theta_2 = \frac{1}{2} \Rightarrow \cos \theta_2 = \frac{1}{2} \Rightarrow \theta_2 = \frac{\pi}{3}$$

$$\text{and } \vec{a} \cdot \vec{b} = 0$$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta_1 = 0$$

$$\Rightarrow \cos \theta_1 = \cos \frac{\pi}{2}$$

$$\Rightarrow \theta_1 = \frac{\pi}{2}$$

229 (b)

$$\vec{OA} + \vec{OB} + \vec{OC}$$

$$= \frac{1}{2} (2\vec{OA} + 2\vec{OB} + 2\vec{OC})$$

$$= \frac{1}{2} \{(\vec{OA} + \vec{OB}) + (\vec{OB} + \vec{OC}) + (\vec{OC} + \vec{OA})\}$$

$$= \frac{1}{2} \{2\vec{OP} + 2\vec{OQ} + 2\vec{OR}\}$$

$$= \vec{OP} + \vec{OQ} + \vec{OR}$$

230 (a)

$$\text{Given, } (\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = \vec{a}^2 \vec{b}^2 \sin^2 \theta + \vec{a}_2 \vec{b}_2 \cos^2 \theta = \vec{a}^2 \vec{b}^2$$

231 (a)

Since,  $\vec{a} = m\vec{b}$  for some scalar  $m$  ie,

$$\vec{a} = m \left( 6\hat{i} - 8\hat{j} - \frac{15}{2}\hat{k} \right)$$

$$\Rightarrow |\vec{a}| = |m| \sqrt{36 + 64 + \frac{225}{4}}$$

$$\Rightarrow 50 = \frac{25}{2} |m| \Rightarrow |m| = 4$$

$$\Rightarrow m = \pm 4$$

Since,  $\vec{a}$  makes an acute angle with the positive direction of  $z$ -axis, so its  $z$  component must be positive and hence,  $m$  must be  $-4$

$$\therefore \vec{a} = -4 \left( 6\hat{i} - 8\hat{j} - \frac{15}{2}\hat{k} \right)$$

$$= -24\hat{i} + 32\hat{j} + 30\hat{k}$$


232 (c)

In  $\triangle ABC$ , we have

$$\vec{AC} = \vec{a} + \vec{b}$$

In  $\triangle ACD$ , we have

$$\vec{AC} + \vec{CD} = \vec{AD} \Rightarrow \vec{CD} = \vec{b} - \vec{a} - \vec{b} = \vec{b} - \vec{a}$$



In  $\triangle CDE$ , we have

$$\vec{CD} + \vec{DE} = \vec{CE} \Rightarrow \vec{b} - \vec{a} - \vec{a} = \vec{CE} \Rightarrow \vec{CE} \\ = \vec{b} - 2\vec{a}$$

233 (b)

Given vectors will be coplanar, if  $\begin{vmatrix} 2 & -3 & 4 \\ 1 & 2 & -1 \\ m & -1 & 2 \end{vmatrix} =$

0

$$\Rightarrow 2(4 - 1) + 3(2 + m) + 4(-1 - 2m) = 0$$

$$\Rightarrow m = \frac{8}{5}$$

234 (d)

Given that,  $|\vec{a}| = 1$ ,  $|\vec{b}| = 3$  and  $|\vec{c}| = 5$

$$\begin{aligned} \therefore [\vec{a} - 2\vec{b} \vec{b} - 3\vec{c} \vec{c} - 4\vec{a}] \\ &= (\vec{a} - 2\vec{b}) \cdot \{ (\vec{b} - 3\vec{c}) \times (\vec{c} - 4\vec{a}) \} \\ &= (\vec{a} - 2\vec{b}) \cdot \{ \vec{b} \times \vec{c} - 4\vec{b} \times \vec{a} + 12\vec{c} \times \vec{a} \} \\ &= (\vec{a} - 2\vec{b}) \cdot (\vec{a} + 4\vec{c} + 12\vec{b}) \\ &= \vec{a} \cdot \vec{a} - 24\vec{b} \cdot \vec{b} = 1 - 24 \times 9 \\ &= 1 - 216 = -215 \end{aligned}$$

235 (a)

Now,  $\hat{i} \times (\hat{j} \times \hat{k}) = \hat{i} \times \hat{i} = \vec{0}$

$\hat{j} \times (\hat{k} \times \hat{i}) = \hat{j} \times \hat{j} = \vec{0}$

and  $\hat{k} \times (\hat{i} \times \hat{j}) \hat{k} \times \hat{k} = \vec{0}$

$$\therefore \hat{i} \times (\hat{j} \times \hat{k}) + \hat{j} \times (\hat{k} \times \hat{i}) + \hat{k} \times (\hat{i} \times \hat{j}) = \vec{0}$$

236 (c)

Given vectors will be coplanar, if

$$\begin{vmatrix} -\lambda^2 & 1 & 1 \\ 1 & -\lambda^2 & 1 \\ 1 & 1 & -\lambda^2 \end{vmatrix} = 0$$

$$\Rightarrow \lambda^6 - 3\lambda^2 - 2 = 0$$

$$\Rightarrow (1 + \lambda^2)^2(\lambda^2 - 2) = 0 \Rightarrow \lambda = \pm\sqrt{2}$$

237 (c)

$$\begin{aligned} \text{Here, force } \vec{F} &= 6 \times \frac{(9\hat{i} + 6\hat{j} + 2\hat{k})}{\sqrt{81 + 36 + 4}} \\ &= \frac{6(9\hat{i} + 6\hat{j} + 2\hat{k})}{11} \end{aligned}$$

Displacement vector  $\vec{d}$

$$= (7 - 3)\hat{i} + (-6 - 4)\hat{j} + (8 + 15)\hat{k}$$

$$= 4\hat{i} - 10\hat{j} + 23\hat{k}$$

$$\therefore \text{Work done} = \vec{F} \cdot \vec{d}$$

$$= \frac{6}{11} (9\hat{i} + 6\hat{j} + 2\hat{k}) \cdot (4\hat{i} - 10\hat{j} + 23\hat{k})$$

$$= \frac{6}{11} (36 - 60 + 46) = 12$$

238 (d)

Since,  $|2\hat{u} \times 3\hat{v}| = 1$

$$\Rightarrow 6|\hat{u}||\hat{v}|\sin\theta = 1$$

$$\Rightarrow \sin\theta = \frac{1}{6} \quad [\because |\hat{u}| = |\hat{v}| = 1]$$

Since,  $\theta$  is an acute angle, then there is exactly one value of  $\theta$  for which  $(2\hat{u} \times 3\hat{v})$  is a unit vector.

239 (d)

$$\therefore \text{Total force, } \vec{F} = \vec{F}_1 + \vec{F}_2$$

$$= 5\hat{i} + \hat{j} - \hat{k}$$

and displacement,  $\vec{d} = (5 - 3)\hat{i} + (5 - 2)\hat{j} + (3 - 1)\hat{k}$

$$= 2\hat{i} + 3\hat{j} + 2\hat{k}$$

$$\therefore W = \vec{F} \cdot \vec{d}$$

$$= (5\hat{i} + \hat{j} - \hat{k}) \cdot (2\hat{i} + 3\hat{j} + 2\hat{k})$$

$$= 11 \text{ units}$$

241 (a)

We have,

$$\vec{a} + t\vec{b} \perp \vec{c}$$

$$\Rightarrow (\vec{a} + t\vec{b}) \cdot \vec{c} = 0$$

$$\Rightarrow \vec{a} \cdot \vec{c} + t\vec{b} \cdot \vec{c} = 0 \Rightarrow t = -\frac{\vec{a} \cdot \vec{c}}{\vec{b} \cdot \vec{c}} = -\frac{6 + 2 + 0}{-3 + 2 + 0} = 8$$

242 (d)

$$\text{Given, } \vec{a} \cdot \vec{p} = \frac{\vec{a} \cdot \vec{b} \times \vec{c}}{[\vec{a} \vec{b} \vec{c}]} = 1$$

$$\text{and } \vec{a} \cdot \vec{q} = \vec{a} \cdot \frac{\vec{c} \times \vec{a}}{[\vec{a} \vec{b} \vec{c}]} = 0$$

Similarly,  $\vec{b} \cdot \vec{q} = \vec{c} \cdot \vec{r} = 1$ ,

and  $\vec{a} \cdot \vec{r} = \vec{b} \cdot \vec{p} = \vec{c} \cdot \vec{q} = \vec{c} \cdot \vec{p} = \vec{b} \cdot \vec{r} = 0$

$$\begin{aligned} \therefore (\vec{a} + \vec{b}) \cdot \vec{p} (\vec{b} + \vec{c}) \cdot \vec{q} + (\vec{c} + \vec{a}) \cdot \vec{r} \\ &= \vec{a} \cdot \vec{p} + \vec{b} \cdot \vec{p} + \vec{b} \cdot \vec{q} + \vec{c} \cdot \vec{q} + \vec{c} \cdot \vec{r} + \vec{a} \cdot \vec{r} \\ &= 1 + 1 + 1 = 3 \end{aligned}$$

243 (b)

$$\vec{a} \cdot \vec{b}_1 + \vec{a} \cdot \left( \vec{b} - \frac{\vec{b} \cdot \vec{a}}{|\vec{a}|^2} \vec{a} \right)$$

$$= \vec{a} \cdot \vec{b} - \frac{|\vec{a}|^2 (\vec{b} \cdot \vec{a})}{|\vec{a}|^2}$$

$$= \vec{a} \cdot \vec{b} - \vec{b} \cdot \vec{a} = 0$$

Similarly,  $\vec{a} \cdot \vec{c}_2 = \vec{b}_1 \cdot \vec{c}_2 = 0$

Hence,  $\{\vec{a}, \vec{b}_1, \vec{c}_2\}$  are mutually orthogonal vectors.

244 (c)

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} = \vec{0},$$

$$[\because \vec{a} \perp \vec{b} \text{ and } \vec{a} \perp \vec{c}]$$

245 (b)

Let  $\vec{a}, \vec{b}, \vec{c}$  be the position vectors of A, B and C respectively. Then, the position vector of G is

$$\frac{\vec{a} + \vec{b} + \vec{c}}{3}$$

Let the position vectors of A', B' and C' be

$\vec{a}, \vec{b}'$  and  $\vec{c}$  respectively. Then, the position vectors

$$\text{of } G' \text{ is } \frac{\vec{a} + \vec{b}' + \vec{c}}{3}$$

$$\begin{aligned} \therefore A\vec{A}' + B\vec{B}' + C\vec{C}' &= (\vec{a} - \vec{a}) + (\vec{b}' - \vec{b}) + (\vec{c} - \vec{c}) \\ \Rightarrow A\vec{A}' + B\vec{B}' + C\vec{C}' &= (\vec{a}' + \vec{b}' + \vec{c}') - (\vec{a} + \vec{b} + \vec{c}) \\ \Rightarrow A\vec{A}' + B\vec{B}' + C\vec{C}' &= 3 \left\{ \frac{\vec{a}' + \vec{b}' + \vec{c}'}{3} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right\} \\ &= 3G\vec{G}' \end{aligned}$$

246 (a)

We have,

$$\begin{aligned} \vec{u} &= \vec{a} - \vec{b}, \vec{v} = \vec{a} + \vec{b} \\ \Rightarrow \vec{u} \times \vec{v} &= (\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2(\vec{a} \times \vec{b}) \\ \Rightarrow |\vec{u} \times \vec{v}| &= 2|\vec{a} \times \vec{b}| \\ \Rightarrow |\vec{u} \times \vec{v}| &= 2\sqrt{|\vec{a}|^2 |\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2} \\ \Rightarrow |\vec{u} \times \vec{v}| &= 2\sqrt{16 - (\vec{a} \cdot \vec{b})^2} \end{aligned}$$

248 (b)

Let  $\vec{d} = d_1\hat{i} + d_2\hat{j} + d_3\hat{k}$

$$\vec{a} \cdot \vec{d} = d_1 - d_2 = 0 \Rightarrow d_1 = d_2 \quad \dots(i)$$

Also,  $\vec{d}$  is a unit vector.

$$\Rightarrow d_1^2 + d_2^2 + d_3^2 = 1 \quad \dots(ii)$$

$$\text{Also, } [\vec{b} \ \vec{c} \ \vec{d}] = 0 \Rightarrow \begin{vmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ d_1 & d_2 & d_3 \end{vmatrix} = 0$$

$$\Rightarrow -1(-d_3 - d_1) - 1(-d_2) = 0$$

$$\Rightarrow d_1 + d_2 + d_3 = 0 \Rightarrow 2d_1 + d_3 = 0 \quad [\text{from Eq. (i)}]$$

(i) ]

$$\Rightarrow d_3 = -2d_1 \quad \dots(iii)$$

Using Eqs. (iii) and (i) in Eq. (ii), we get

$$d_1^2 + d_1^2 + 4d_1^2 = 1 \Rightarrow d_1 = \pm \frac{1}{\sqrt{6}}$$

$$\therefore d_2 = \pm \frac{1}{\sqrt{6}}$$

$$\text{and } d_3 = \mp \frac{2}{\sqrt{6}}$$

Hence, required vector is

$$\pm \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} - 2\hat{k})$$

249 (b)

Since  $\vec{a}$  is collinear to vector  $\vec{b}$ . Therefore,

$$\vec{a} = m\vec{b} \text{ for some scalar } m$$

$$\Rightarrow \vec{a} = m \left( 6\hat{i} - 8\hat{j} - \frac{15}{2}\hat{k} \right)$$

$$\Rightarrow |\vec{a}| = \frac{25}{2}|m|$$

$$\begin{aligned} \Rightarrow 50 &= \frac{25}{2}|m| \Rightarrow |m| = 4 \Rightarrow m \\ &= \pm 4 \quad [ \because |\vec{a}| = 50 ] \end{aligned}$$

Since  $\vec{a}$  makes an acute angle with the positive direction of z-axis. So, its z-component must be positive, and hence 'm' must be -4

$$\therefore \vec{a} = -4 \left( 6\hat{i} - 8\hat{j} - \frac{15}{2}\hat{k} \right) = -24\hat{i} + 32\hat{j} + 30\hat{k}$$

251 (c)

Since  $\vec{a}$  and  $\vec{b}$  are coplanar. Therefore,  $\vec{a} \times \vec{b}$  is a vector perpendicular to the plane containing  $\vec{a}$  and  $\vec{b}$

Similarly,  $\vec{c} \times \vec{d}$  is a vector perpendicular to the plane containing  $\vec{c}$  and  $\vec{d}$

Two planes will be parallel if their normal i.e.  $\vec{a} \times \vec{b}$  and  $\vec{c} \times \vec{d}$  are parallel

$$\therefore (\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = 0$$

252 (c)

$$\text{Since, } \vec{a} \cdot (\vec{b} + \vec{c}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{c} \cdot \vec{a} = 0 \quad \dots(i)$$

$$\text{Similarly, } \vec{b} \cdot \vec{c} + \vec{a} \cdot \vec{b} = 0 \quad \dots(ii)$$

$$\text{and } \vec{c} \cdot \vec{a} + \vec{b} \cdot \vec{c} = 0 \quad \dots(iii)$$

On adding Eqs. (i), (ii) and (iii), we get

$$2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2$$

$$= 9 + 16 + 25 = 50$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = 5\sqrt{2}$$

253 (b)

We know that the diagonals of a parallelogram bisect each other. Therefore, M is the mid point of AC and BD both.

$$\therefore \vec{OA} + \vec{OC} = 2\vec{OM}$$

$$\text{and } \vec{OB} + \vec{OD} = 2\vec{OM}$$

$$\Rightarrow \vec{OA} + \vec{OB} + \vec{OC} + \vec{OD} = 4\vec{OM}$$

254 (b)

$$|\vec{OA}| = \sqrt{4 + 4 + 1} = 3$$

$$\text{and } |\vec{OB}| = \sqrt{4 + 16 + 16} = 6$$

$$\therefore \text{Required vector} = \lambda(\vec{OA} + \vec{OB})$$

$$= \lambda \left[ \frac{1}{3}(2\hat{i} + 2\hat{j} + \hat{k}) + \frac{1}{6}(2\hat{i} + 4\hat{j} + 4\hat{k}) \right]$$

$$= \frac{\lambda}{3}(3\hat{i} + 4\hat{j} + 3\hat{k})$$

$$\therefore \text{Length of vector} = \frac{\lambda}{3}\sqrt{9 + 16 + 9} = \frac{\lambda}{3}\sqrt{34}$$

Take  $\lambda = 2$

∴ Required length of a vector is  $\frac{\sqrt{136}}{3}$

255 (d)

Given that,  $\vec{A} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{B} = \hat{i}$ ,  $\vec{C} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$

Since,  $\vec{A}$ ,  $\vec{B}$ ,  $\vec{C}$  are coplanar.

$$\therefore [\vec{A} \vec{B} \vec{C}] = 0$$

$$\text{Now, } \vec{B} \times \vec{C} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & 0 \\ c_1 & c_2 & c_3 \end{vmatrix} = -c_3\hat{j} + c_2\hat{k}$$

$$\therefore \vec{A} \cdot (\vec{B} \times \vec{C}) = (\hat{i} + \hat{j} + \hat{k}) \cdot (-c_3\hat{j} + c_2\hat{k}) = 0$$


⇒ No value of  $c_1$  can be found.

256 (c)

We have,

$$\begin{aligned} & \vec{AB} \cdot \vec{AC} + \vec{BC} \cdot \vec{BA} + \vec{CA} \cdot \vec{CB} \\ &= (AB)(AC) \cos \theta + (BC)(BA) \sin \theta + 0 \\ &= AB(AC \cos \theta + BC \sin \theta) \\ &= AB \left\{ \frac{(AC)^2}{AB} + \frac{(BC)^2}{AB} \right\} \left[ \because \cos \theta = \frac{AC}{AB}, \sin \theta \right. \\ & \quad \left. = \frac{BC}{AB} \right] \end{aligned}$$

$$= AC^2 + BC^2 = AB^2 = p^2$$



257 (a)

The position vector of the centroid of the triangle is  $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$

Since the triangle is an equilateral. Therefore, the orthocenter coincides with the centroid and hence

$$\frac{\vec{a} + \vec{b} + \vec{c}}{3} = \vec{0} \Rightarrow \vec{a} + \vec{b} + \vec{c} = \vec{0}$$

258 (d)

$$\text{Given, } |\vec{a} \times \vec{b}| = 4$$

$$\Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 4$$

$$\Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 4 \quad [\because |\hat{n}| = 1] \dots (i)$$

$$\text{Also, } |\vec{a} \cdot \vec{b}| = 2$$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta = 2 \quad \dots (ii)$$

On squaring and then on adding Eqs. (i) and (ii), we get

$$|\vec{a}|^2 |\vec{b}|^2 \sin^2 \theta + |\vec{a}|^2 |\vec{b}|^2 \cos^2 \theta = 4^2 + 2^2$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 (\sin^2 \theta + \cos^2 \theta) = 16 + 4$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 = 20$$

260 (d)

Given that,  $\vec{a} = 2\hat{i} - 3\hat{j} + 5\hat{k}$ ,  $\vec{b} = 3\hat{i} - 4\hat{j} + 5\hat{k}$  and  $\vec{c} = 5\hat{i} - 3\hat{j} - 2\hat{k}$

∴ Volume of parallelepiped where sides are  $\vec{a} + \vec{b}$ ,  $\vec{b} + \vec{c}$  and  $\vec{c} + \vec{a}$ , is

$$\begin{aligned} [\vec{a} + \vec{b} \vec{b} + \vec{c} \vec{c} + \vec{a}] &= \begin{vmatrix} 2 & -3 & 5 \\ 3 & -4 & 5 \\ 5 & -3 & -2 \end{vmatrix} \\ &= [2(8 + 15) + 3(-6 - 25) + 5(-9 + 20)] \\ &= 46 - 93 + 55 = 8 \end{aligned}$$

261 (c)

Let  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$

$$\text{Given, } \vec{a} \cdot \hat{i} = \vec{a} \cdot (\hat{i} + \hat{j}) = \vec{a} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$$

$$\therefore a_1 = a_1 + a_2 = a_1 + a_2 + a_3 = 1$$

$$\Rightarrow a_1 = 1, a_2 = 0, a_3 = 0$$

$$\therefore \vec{a} = \hat{i}$$

263 (b)

$$\begin{aligned} & \vec{DA} + \vec{DB} + \vec{DC} + \vec{AE} + \vec{BE} + \vec{CE} \\ &= (\vec{DA} + \vec{AE}) + (\vec{DB} + \vec{BE}) + (\vec{DC} + \vec{CE}) \\ &= \vec{DE} + \vec{DE} + \vec{DE} \\ &= 3 \vec{DE} \end{aligned}$$

265 (d)

Given vertices are

$$A(3\hat{i} + \hat{j} + 2\hat{k}), B(\hat{i} - 2\hat{j} + 7\hat{k}) \text{ and } C(-2\hat{i} + 3\hat{j} + 5\hat{k}).$$

$$\text{Now, } \vec{AB} = (\hat{i} - 2\hat{j} + 7\hat{k}) - (3\hat{i} + \hat{j} + 2\hat{k})$$

$$= -2\hat{i} - 3\hat{j} + 5\hat{k}$$

$$\therefore |\vec{AB}| = \sqrt{4 + 9 + 25} = \sqrt{38}$$

$$\text{Similarly, } |\vec{BC}| = |\vec{CA}| = \sqrt{38}$$

$$\therefore |\vec{AB}| = |\vec{BC}| = |\vec{CA}| = \sqrt{38}$$

∴ Hence, triangle is an equilateral triangle.

267 (b)

We have,

$$|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = |\vec{a} - \vec{b}|^2$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = |\vec{a}|^2 + |\vec{b}|^2 - 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 4\vec{a} \cdot \vec{b} = 0 \Rightarrow \vec{a} \cdot \vec{b} = 0 \Rightarrow \vec{a} \perp \vec{b}$$

268 (d)

$$\because |\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = |\vec{a} - \vec{b}|^2$$

$$\Rightarrow a^2 + b^2 + 2\vec{a} \cdot \vec{b} = a^2 + b^2 - 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 4\vec{a} \cdot \vec{b} = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} = 0$$

$\therefore$  Angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{2}$ .

269 (c)

Given vectors are coplanar, if  $\begin{vmatrix} \alpha & 1 & 1 \\ 1 & \beta & 1 \\ 0 & c & \gamma \end{vmatrix} = 0$

Applying  $C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_2$

$$\Rightarrow \begin{vmatrix} \alpha & 1-\alpha & 0 \\ 1 & \beta-1 & 1-\beta \\ 1 & 0 & \gamma-1 \end{vmatrix} = 0$$

$$\Rightarrow (1-\alpha)(1-\beta)(1-\gamma) \begin{vmatrix} \alpha & 1 & 0 \\ 1-\alpha & -1 & 1 \\ 1-\gamma & 0 & -1 \end{vmatrix} = 0$$

$$\Rightarrow (1-\alpha)(1-\beta)(1-\gamma) \left[ \frac{\alpha}{1-\alpha} (1) - 1 \left( -\frac{1}{1-\beta} - \frac{1}{1-\gamma} \right) \right] = 0$$

But  $\alpha \neq 1, \beta \neq 1$  and  $\gamma \neq 1$

$$\therefore \frac{1}{(1-\alpha)} - 1 + \frac{1}{1-\beta} + \frac{1}{1-\gamma} = 0$$

$$\Rightarrow \frac{1}{1-\alpha} + \frac{1}{1-\beta} + \frac{1}{1-\gamma} = 1$$

270 (b)

Let the required vector be  $\vec{c} = x\hat{i} + z\hat{k}$

Since,  $|\vec{c}| = 1 \Rightarrow x^2 + z^2 = 1 \dots (i)$

$\vec{a}$  and  $\vec{c}$  are inclined at the angle  $45^\circ$

$$\therefore \cos 45^\circ = \frac{2x - z}{\sqrt{4 + 4 + 1}} \Rightarrow 2x - z = \frac{3}{\sqrt{2}} \dots (ii)$$

$\vec{b}$  and  $\vec{c}$  are inclined at an angle  $60^\circ$

$$\therefore -\frac{z}{\sqrt{2}} = \cos 60^\circ \Rightarrow z = -\frac{1}{\sqrt{2}} \dots (iii)$$

From Eqs. (ii) and (iii), we get  $x = \frac{1}{\sqrt{2}}$

Hence, the required vector is  $\frac{1}{\sqrt{2}}\hat{i} - \frac{1}{\sqrt{2}}\hat{k}$

271 (d)

Since  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar vectors. Therefore,

$$[\vec{a} \vec{b} \vec{c}] \neq 0$$

$$\Rightarrow \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} \neq 0 \Rightarrow \Delta \neq 0, \text{ where } \Delta = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix}$$

Now,

$$\begin{vmatrix} a & a^2 & 1+a^3 \\ b & b^2 & 1+b^3 \\ c & c^2 & 1+c^3 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} a & a^2 & 1 \\ b & b^2 & 1 \\ c & c^2 & 1 \end{vmatrix} + \begin{vmatrix} a & a^2 & a^3 \\ b & b^2 & b^3 \\ c & c^2 & c^3 \end{vmatrix} = 0$$

$$\Rightarrow \Delta(1 + abc) = 0 \Rightarrow abc = -1 \quad [\because \Delta \neq 0]$$

272 (c)

$$\hat{u} \cdot \hat{v} = 0$$

$$\Rightarrow |\hat{u}| |\hat{v}| \cos \theta = 0$$

$$\Rightarrow 1 \times 1 \times \cos \theta = 0 \quad (\because |\hat{u}| = |\hat{v}| = 1)$$

$$\Rightarrow \cos \theta = 0$$

$$\Rightarrow \theta = 90^\circ$$

Let  $\hat{n}$  be a unit vector perpendicular to the plane of vectors  $\hat{u}$  and  $\hat{v}$ .

$$\Rightarrow \hat{u} \times \hat{v} = |\hat{u}| |\hat{v}| \sin 90^\circ \cdot \hat{n} = \hat{n}$$

Since,  $\vec{r}$  is coplanar with  $\hat{u}$  and  $\hat{v}$

$\therefore \hat{n}$  is perpendicular to  $\vec{r}$

Let  $\Phi$  be the angle between  $\hat{n}$  and  $\vec{r}$

$$\Rightarrow \Phi = 90^\circ$$

$$\therefore |\vec{r} \times (\hat{u} \times \hat{v})| = |\vec{r} \times \hat{n}| = |\vec{r}| |\hat{n}| \sin \Phi$$

$$= |\vec{r}| \times 1 \times \sin 90^\circ$$

$$= |\vec{r}|$$

273 (b)

Let  $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}, \vec{b} = 4\hat{i} - 4\hat{j} + 7\hat{k}$ . Then,

$$\text{Projection of } \vec{a} \text{ on } \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{4+8+7}{\sqrt{16+16+49}} = \frac{19}{9}$$

274 (a)

$$\frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{\vec{b} \cdot (\vec{c} \times \vec{a})} + \frac{\vec{b} \cdot (\vec{a} \times \vec{c})}{\vec{a} \cdot (\vec{b} \times \vec{c})}$$

$$= \frac{[\vec{a} \vec{b} \vec{c}]}{[\vec{b} \vec{c} \vec{a}]} + \frac{[\vec{b} \vec{a} \vec{c}]}{[\vec{a} \vec{b} \vec{c}]} = 1 + 0 = 1$$

275 (b)

$$\left[ \frac{1}{2} |\vec{u}_2 - \vec{u}_1| \right]^{-2} = \frac{1}{4} [|\vec{u}_2|^2 + |\vec{u}_1|^2 - 2\vec{u}_2 \cdot \vec{u}_1]$$

$$= \frac{1}{4} [1 + 1 - 2|\vec{u}_2||\vec{u}_1| \cos \theta]$$

$$= \frac{1}{4} [2 - 2 \cos \theta] = \sin^2 \frac{\theta}{2}$$

$$\Rightarrow \frac{1}{2} |\vec{u}_2 - \vec{u}_1| = \sin \frac{\theta}{2}$$

276 (b)

Let  $\vec{c} = x\hat{i} + y\hat{j}$ . Then,

$$\vec{b} \perp \vec{c}$$

$$\Rightarrow \vec{b} \cdot \vec{c} = 0$$

$$\Rightarrow 4x + 3y = 0 \Rightarrow \frac{x}{3} = \frac{y}{-4} = \lambda \Rightarrow x = 3\lambda, y = -4\lambda$$

$$\therefore \vec{c} = \lambda(3\hat{i} - 4\hat{j})$$

Let the required vector be  $\alpha = p\hat{i} + q\hat{j}$ . Then the

projections of  $\vec{\alpha}$  on  $\vec{b}$  and  $\vec{c}$  are  $\frac{\vec{\alpha} \cdot \vec{b}}{|\vec{b}|}$  respectively

$$\therefore \frac{\vec{\alpha} \cdot \vec{b}}{|\vec{b}|} = 1 \text{ and } \frac{\vec{\alpha} \cdot \vec{c}}{|\vec{c}|} = 2$$

$$\Rightarrow 4p + 3q = 5 \text{ and } 3p - 4q = 10 \Rightarrow p = 2, q = -1$$

Hence, the required vector =  $2\hat{i} - \hat{j}$

277 (b)

Given equation of plane is

$$2x + 4y - 5z = 10$$

Here,  $a = 2, b = 4, c = -5$

Let  $OP$  be the perpendicular from  $O$  to the plane, then its equation is

$$\frac{x-0}{2} = \frac{y-0}{4} = \frac{z-0}{-5}$$

Here, direction ratio are  $(2, 4, -5)$ .

Now, equation of line in vector form is

$$\vec{r} = 0 + k(2, 4, -5)$$

$$= (2k, 4k, -5k), k \in R$$

[ $\therefore$  equation of line is  $\vec{r} = \vec{a} + \lambda \vec{b}$ ]

278 (a)

We have,

$$\vec{a} = \lambda \{ \vec{b} \times (\hat{i} \times \hat{j}) \} = \lambda \{ \vec{b} \cdot \hat{j} \} \hat{i} - (\vec{b} \cdot \hat{i}) \hat{j} \\ = \lambda(-3\hat{i} - 4\hat{j})$$

$$\text{Now, } |\vec{a}| = |\vec{b}| \Rightarrow 25\lambda^2 = 16 + 9 + 25 \Rightarrow \lambda = \pm\sqrt{2}$$

$$\text{Hence, } \vec{a} = \pm\sqrt{2}(3\hat{i} + 4\hat{j})$$

279 (d)

Given  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 3^2 + 4^2 + 5^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -25$$

280 (a)

We know that the position vector of the centroid of the triangle is

$$\frac{\vec{a} + \vec{b} + \vec{c}}{3}$$

Since, the triangle is an equilateral, therefore the orthocentre coincides

With the centroid and hence,

$$\frac{\vec{a} + \vec{b} + \vec{c}}{3} = \vec{0}$$

$$\Rightarrow \vec{a} + \vec{b} + \vec{c} = \vec{0}$$

281 (a)

$$\vec{AB} = 2\hat{i} + 3\hat{j} + 4\hat{k} - 4\hat{i} - 7\hat{j} - 8\hat{k} \\ = -2\hat{i} - 4\hat{j} - 4\hat{k}$$

$$\text{and } \vec{AC} = 2\hat{i} + 5\hat{j} + 7\hat{k} - 4\hat{i} - 7\hat{j} - 8\hat{k} = -2\hat{i} - 2\hat{j} - \hat{k}$$

$$\therefore |\vec{AB}| = 6 \text{ and } |\vec{AC}| = 3$$

$\therefore$  Position vector of required bisector

$$= \frac{6(2\hat{i} + 5\hat{j} + 7\hat{k}) + 3(2\hat{i} + 3\hat{j} + 4\hat{k})}{6 + 3}$$

$$= \frac{1}{3}(6\hat{i} + 13\hat{j} + 18\hat{k})$$

282 (a)

Since  $\vec{a}$  and  $\vec{b}$  are collinear vectors. Therefore,  $\vec{b} = \lambda \vec{a}$

$$\Rightarrow \vec{b} = \lambda(2\hat{i} + 3\hat{j} + 6\hat{k})$$

$$\Rightarrow |\vec{b}| = |\lambda|\sqrt{4 + 9 + 36} \Rightarrow 21 = 7|\lambda| \Rightarrow \lambda = \pm 3$$

$$\therefore \vec{b} = \pm 3\vec{a} = \pm(6\hat{i} + 9\hat{j} + 18\hat{k})$$

283 (a)

We have,

$$\vec{a} - \vec{b} + \vec{b} - \vec{c} + \vec{c} - \vec{a} = 0$$

$$\Rightarrow \vec{a} - \vec{b}, \vec{b} - \vec{c} \text{ and } \vec{c} - \vec{a} \text{ are coplanar}$$

$$\Rightarrow [\vec{a} - \vec{b} \vec{b} - \vec{c} \vec{c} - \vec{a}] = 0$$

284 (c)

$$\text{Here, } (\hat{i} - \hat{j} + 4\hat{k}) \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 0$$

It means line is parallel to the plane

General point on the line is  $(\lambda + 2, -\lambda - 2, 4\lambda + 3)$

For  $\lambda = 0$ , point on this line is  $(2, -2, 3)$  and

distance from

$$\vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 5 \text{ is}$$

$$d = \frac{|2 + 5(-2) + 3 - 5|}{\sqrt{(1)^2 + (5)^2 + (1)^2}} = \frac{10}{3\sqrt{3}}$$

286 (b)

$$\therefore \vec{a} + \vec{b} = \hat{i} + 2\hat{j} - 3\hat{k} + 3\hat{i} - \hat{j} + 2\hat{k}$$

$$= 4\hat{i} + \hat{j} - \hat{k}$$

$$\text{and } \vec{a} - \vec{b} = (\hat{i} + 2\hat{j} - 3\hat{k}) - (3\hat{i} - \hat{j} + 2\hat{k})$$

$$= -2\hat{i} + 3\hat{j} - 5\hat{k}$$

$$\therefore \cos \theta = \frac{(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b})}{|\vec{a} + \vec{b}| |\vec{a} - \vec{b}|}$$

$$= \frac{(4\hat{i} + \hat{j} - \hat{k}) \cdot (-2\hat{i} + 3\hat{j} - 5\hat{k})}{|4\hat{i} + \hat{j} - \hat{k}| |-2\hat{i} + 3\hat{j} - 5\hat{k}|}$$

$$= \frac{-8 + 3 + 5}{\sqrt{16 + 1 + 1} \sqrt{4 + 9 + 25}} = 0$$

$$\Rightarrow \theta = 90^\circ$$

288 (a)

$$\text{Given, } \vec{a} = \vec{b} + \vec{c}$$

$$\text{and } \vec{b} \perp \vec{c}$$

$$\text{then } |\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{b} \cdot \vec{c}$$

$$\Rightarrow a^2 = b^2 + c^2 (\because \vec{b} \cdot \vec{c} = 0)$$

289 (b)

$$\therefore \vec{AB} = \vec{OB} - \vec{OA}$$

$$\therefore \vec{OB} = \vec{AB} + \vec{OA}$$

$$= 3\hat{i} - \hat{j} + \hat{k} + 3\hat{i} - 2\hat{j} + 4\hat{k}$$

$$= 6\hat{i} - 3\hat{j} + 5\hat{k}$$

290 (a)

Given that,

$$\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 3\hat{j} + 5\hat{k}$$

$$\text{and } \vec{c} = 7\hat{i} + 9\hat{j} + 11\hat{k}$$

Let  $\vec{A} = \vec{a} + \vec{b} = (\hat{i} + \hat{j} + \hat{k}) + (\hat{i} + 3\hat{j} + 5\hat{k}) = 2\hat{i} + 4\hat{j} + 6\hat{k}$

And  $\vec{B} = \vec{b} + \vec{c} = (\hat{i} + 3\hat{j} + 5\hat{k}) + (7\hat{i} + 9\hat{j} + 11\hat{k}) = 8\hat{i} + 12\hat{j} + 16\hat{k}$

If  $\vec{A}$  and  $\vec{B}$  are diagonals, then area of parallelogram

$$\begin{aligned} &= \frac{1}{2} |\vec{A} \times \vec{B}| = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 4 & 6 \\ 8 & 12 & 16 \end{vmatrix} \\ &= \frac{1}{2} |\hat{i}(64 - 72) - \hat{j}(32 - 48) + \hat{k}(24 - 32)| \\ &= \frac{1}{2} |-8\hat{i} + 16\hat{j} - 8\hat{k}| \\ &= |-4\hat{i} + 8\hat{j} - 4\hat{k}| \\ &= \sqrt{(-4)^2 + (8)^2 + (-4)^2} \\ &= \sqrt{16 + 64 + 16} = \sqrt{96} = 4\sqrt{6} \end{aligned}$$

291 (a)

Given that,  $\vec{a} = (1, 1, 4) = \hat{i} + \hat{j} + 4\hat{k}$

and  $\vec{b} = (1, -1, 4) = \hat{i} - \hat{j} + 4\hat{k}$

$\therefore \vec{a} + \vec{b} = 2\hat{i} + 8\hat{k}$

$\Rightarrow \vec{a} - \vec{b} = 2\hat{j}$

Let  $\theta$  be the angle between  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$ , then

$$\begin{aligned} \cos \theta &= \frac{(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b})}{|\vec{a} + \vec{b}| |\vec{a} - \vec{b}|} \\ &= \frac{(2\hat{i} + 0\hat{j} + 8\hat{k}) \cdot (0\hat{i} + 2\hat{j} + 0\hat{k})}{\sqrt{2^2 + 0^2 + 8^2} \sqrt{0^2 + 2^2 + 0^2}} \\ &= \frac{0 + 0 + 0}{\sqrt{4 + 64} \sqrt{4}} = 0 \\ \Rightarrow \cos \theta &= \cos \theta^\circ \Rightarrow \theta = \frac{\pi}{2} = 90^\circ \end{aligned}$$

292 (c)

Area of rhombus  $= \frac{1}{2} |\vec{a} \times \vec{b}|$

$$\begin{aligned} &= \frac{1}{2} |(2\hat{i} - 3\hat{j} + 5\hat{k}) \times (-\hat{i} + \hat{j} + \hat{k})| \\ &= \frac{1}{2} |-8\hat{i} - 7\hat{j} - \hat{k}| = \frac{1}{2} \sqrt{144} \\ &= \sqrt{28.5} \end{aligned}$$

293 (a)

It is given that the vectors  $\hat{i} - 2x\hat{j} - 3y\hat{k}$  and  $\hat{i} + 3x\hat{j} + 2y\hat{k}$  are orthogonal

$\therefore (\hat{i} - 2x\hat{j} - 3y\hat{k}) \cdot (\hat{i} + 3x\hat{j} + 2y\hat{k}) = 0$

$\Rightarrow 1 - 6x^2 - 6y^2 = 0 \Rightarrow 6x^2 + 6y^2 = 1$

Clearly, it represents a circle

295 (a)

Given vectors are orthogonal.

$\therefore (3x\hat{i} + y\hat{j} - 3\hat{k}) \cdot (x\hat{i} - 4y\hat{j} + 4\hat{k}) = 0$

$\Rightarrow 3x^2 - 4y^2 - 12 = 0$

$$\Rightarrow \frac{x^2}{4} - \frac{y^2}{3} = 1$$

Hence, it represent a hyperbola.

296 (c)

We have,  $|\vec{a}| = 1$ ,  $|\vec{b}| = 1$  and  $|\vec{a} + \vec{b}| = 1$

Now,

$$|\vec{a} + \vec{b}|^2 + |\vec{a} - \vec{b}|^2 = 2\{|\vec{a}|^2 + |\vec{b}|^2\}$$

$$\Rightarrow 1 + |\vec{a} - \vec{b}|^2 = 4$$

$$\Rightarrow |\vec{a} - \vec{b}| = \sqrt{3}$$

298 (a)

Let unit vector is  $a\hat{i} + b\hat{j} + c\hat{k}$ .

$\therefore a\hat{i} + b\hat{j} + c\hat{k}$  is perpendicular to  $\hat{i} + \hat{j} + \hat{k}$ .

Then,  $a + b + c = 0$  ... (i)

and  $a\hat{i} + b\hat{j} + c\hat{k}$ ,  $(\hat{i} + \hat{j} + 2\hat{k})$  and  $(\hat{i} + 2\hat{j} + \hat{k})$  are coplanar.

$$\therefore \begin{vmatrix} a & b & c \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -3a + b + c = 0 \dots (ii)$$

From Eqs. (i) and (ii), we get

$$a = 0 \text{ and } c = -b$$

$\therefore a\hat{i} + b\hat{j} + c\hat{k}$  is a unit vector, then

$$a^2 + b^2 + c^2 = 1$$

$$\Rightarrow 0 + b^2 + b^2 = 1$$

$$\Rightarrow b = \frac{1}{\sqrt{2}}$$

$$\therefore a\hat{i} + b\hat{j} + c\hat{k} = \frac{1}{\sqrt{2}}\hat{j} - \frac{1}{\sqrt{2}}\hat{k} = \frac{\hat{j} - \hat{k}}{\sqrt{2}}$$

300 (b)

Given,  $\vec{r} = (1 + t)\hat{i} - (1 - t)\hat{j} + (1 - t)\hat{k}$

and  $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 5$

Since, they intersect, therefore

$$(1 + t) - (1 - t) + (1 - t) = 5$$

$$\Rightarrow t = 4$$

$$\therefore \vec{r} = (1 + 4)\hat{i} - (1 - 4)\hat{j} + (1 - 4)\hat{k}$$

$$= 5\hat{i} + 3\hat{j} - 3\hat{k}$$

301 (d)

We have,

$$|\vec{a}| = 3, |\vec{b}| = 5 \text{ and } |\vec{c}| = 7$$

Let  $\theta$  be the angle between  $\vec{a}$  and  $\vec{b}$

Now,  $\vec{a} + \vec{b} + \vec{c} = 0$

$$\Rightarrow |\vec{c}|^2 = |\vec{a} + \vec{b}|^2$$

$$\Rightarrow |\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow |\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos \theta$$

$$\Rightarrow 49 = 9 + 25 + 2 \times 3 \times 5 \cos \theta$$

$$\Rightarrow 15 = 30 \cos \theta \Rightarrow \cos \theta = 1/2 \Rightarrow \theta = \pi/3$$

302 (c)

$$\begin{aligned} \therefore [\vec{a} \vec{b} \vec{c}] &= \vec{a} \cdot \left( |\vec{b}| |\vec{c}| \sin \frac{2\pi}{3} \hat{n} \right) \\ &= |\vec{a}| |\vec{b}| |\vec{c}| \left( \sin \frac{2\pi}{3} \right) \\ [\because \vec{a} \cdot \hat{n} &= |\vec{a}| \hat{n} \cos 0^\circ = |\vec{a}|] \\ &= 2 \times 3 \times 4 \times \frac{\sqrt{3}}{2} = 12\sqrt{3} \end{aligned}$$

303 (a)

Given that,  $\vec{OA} = 2\hat{i} + \hat{j} - \hat{k}$ ,  $\vec{OB} = 3\hat{i} - 2\hat{j} + \hat{k}$  and

$$\vec{OC} = \hat{i} + 4\hat{j} - 3\hat{k}$$

$$\vec{AB} = \vec{OB} - \vec{OA}$$

$$\begin{aligned} &= (3 - 2)\hat{i} + (-2 - 1)\hat{j} + (1 + 1)\hat{k} \\ &= \hat{i} - 3\hat{j} + 2\hat{k} \end{aligned}$$

$$|\vec{AB}| = \sqrt{1^2 + (-3)^2 + 2^2}$$

$$= \sqrt{1 + 9 + 4} = \sqrt{14}$$

$$\vec{BC} = \vec{OC} - \vec{OB}$$

$$\begin{aligned} &= (1 - 3)\hat{i} + (4 + 2)\hat{j} + (-3 - 1)\hat{k} \\ &= -2\hat{i} + 6\hat{j} - 4\hat{k} \end{aligned}$$

$$|\vec{BC}| = \sqrt{(-2)^2 + 6^2 + (-4)^2}$$

$$= \sqrt{4 + 36 + 16} = \sqrt{56}$$

$$\vec{CA} = \vec{OA} - \vec{OC}$$

$$\begin{aligned} &= (2 - 1)\hat{i} + (1 - 4)\hat{j} + (-1 + 3)\hat{k} \\ &= \hat{i} - 3\hat{j} + 2\hat{k} \end{aligned}$$

$$|\vec{CA}| = \sqrt{1^2 + (-3)^2 + (2)^2}$$

$$= \sqrt{1 + 9 + 4} = \sqrt{14}$$

It is clear that two sides of a triangle are equal.

$\therefore$  Points  $A, B, C$  form an isosceles triangle.

304 (b)

The component of  $\vec{a}$  along  $\vec{b}$  is given by

$$\left\{ \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right\} = \frac{18}{25} (3\hat{j} + 4\hat{k})$$

305 (a)

It is given that  $\vec{c}$  and  $\vec{d}$  are collinear vectors

$$\therefore \vec{c} = \lambda \vec{d} \text{ for some scalar } \lambda$$

$$\Rightarrow (x - 2)\vec{a} + \vec{b} = \lambda \{ (2x + 1)\vec{a} - \vec{b} \}$$

$$\Rightarrow \{ x - 2 - \lambda(2x + 1) \} \vec{a} + (\lambda + 1)\vec{b} = \vec{0}$$

$$\Rightarrow \lambda + 1 = 0 \text{ and } x - 2 - \lambda(2x + 1) = 0 \quad [\because \vec{a}, \vec{b} \text{ are non-collinear}]$$

$$\Rightarrow \lambda = -1 \text{ and } x = \frac{1}{3}$$

306 (a)

Equation of plane is  $\vec{r} \cdot \hat{n} = d$ ,

where  $d$  is the perpendicular distance of the plane from origin


$$\therefore \text{Required plane is } (\alpha x + \beta y + \gamma z) = p$$

307 (c)

In  $\Delta ABC$ ,  $\vec{AB} + \vec{BC} + \vec{AC}$

$$\Rightarrow \vec{AC} = \vec{a} + \vec{b}$$

$AD$  is parallel to  $BC$  and  $AD = 2BC$



$$\therefore \vec{AD} = 2\vec{b}$$

In  $\Delta ACD$ ,  $\vec{AC} + \vec{CD} = \vec{AD}$


$$\Rightarrow \vec{CD} = 2\vec{b} - (\vec{a} + \vec{b}) = \vec{b} - \vec{a}$$

$$\text{Now, } \vec{CE} = \vec{CD} + \vec{DE} = \vec{b} - 2\vec{a}$$

309 (d)

Let  $\vec{R}_1 = 2\hat{i} + 4\hat{j} - 5\hat{k}$

and  $\vec{R}_2 = \hat{i} + 2\hat{j} + 3\hat{k}$



$$\therefore \vec{R} \text{ (along } \vec{AC}) = \vec{R}_1 + \vec{R}_2 = 3\hat{i} + 6\hat{j} - 2\hat{k}$$

$$\therefore \vec{a} \text{ (unit vector angle } \vec{AC}) = \frac{\vec{R}}{|\vec{R}|} = \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{9 + 36 + 4}}$$

$$= \frac{1}{7} (3\hat{i} + 6\hat{j} - 2\hat{k})$$

311 (b)

Since  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar vectors. Therefore,

$\vec{a}, \vec{b}, \vec{c}$  are linearly independent vectors

$$\therefore x\vec{a} + y\vec{b} + z\vec{c} = \vec{0} \Rightarrow x = y = z = 0$$

312 (a)

Suppose point  $\hat{i} + 2\hat{j} + 3\hat{k}$  divides the join of points  $-2\hat{i} + 3\hat{j} + 5\hat{k}$  and  $7\hat{i} - \hat{k}$  in the ratio  $\lambda : 1$ .

Then,

$$\hat{i} + 2\hat{j} + 3\hat{k} = \frac{\lambda(7\hat{i} - \hat{k}) + (-2\hat{i} + 3\hat{j} + 5\hat{k})}{\lambda + 1}$$

$$\Rightarrow (\lambda + 1)\hat{i} + 2(\lambda + 1)\hat{j} + 3(\lambda + 1)\hat{k}$$

$$= (7\lambda - 2)\hat{i} + 3\hat{j} + (-\lambda + 5)\hat{k}$$

$$\Rightarrow \lambda + 1 = 7\lambda - 2, 2(\lambda + 1) = 3 \text{ and } 3(\lambda + 1) = -\lambda + 5$$

$$\Rightarrow \lambda = \frac{1}{2}$$

Hence, required ratio is  $1 : 2$

313 (d)

Clearly,

$$\vec{a} - \vec{b} + \vec{b} - \vec{c} + \vec{c} - \vec{a} = \vec{0}$$

$$\therefore \vec{a} - \vec{b}, \vec{b} - \vec{c}, \vec{c} - \vec{a} \text{ are coplanar}$$

$$\Rightarrow (\vec{a} - \vec{b}) \cdot \{(\vec{b} - \vec{c}) \times (\vec{c} - \vec{a})\} = 0$$

314 (d)

Two given lines intersect, if

$$7\hat{i} + 10\hat{j} + 13\hat{k} + s(2\hat{i} + 3\hat{j} + 4\hat{k})$$

$$= 3\hat{i} + 5\hat{j} + 7\hat{k} + t(\hat{i} + 2\hat{j} + 3\hat{k})$$

$$\Rightarrow (7 + 2s)\hat{i} + (10 + 3s)\hat{j} + (13 + 4s)\hat{k}$$

$$= (3 + t)\hat{i} + (5 + 2t)\hat{j} + (7 + 3t)\hat{k}$$

$$\Rightarrow 7 + 2s = 3 + t$$

$$\Rightarrow 2s - t = -4 \dots(i)$$

$$10 + 3s = 5 + 2t$$

$$\Rightarrow 3s - 2t = -5 \dots(ii)$$

and  $13 + 4s = 7 + 3t$

$$\Rightarrow 4s - 3t = -6 \dots(iii)$$

On solving Eqs. (i) and (iii), we get

$$s = -3, t = -2$$

$\therefore$  Required line is

$$7\hat{i} + 10\hat{j} + 13\hat{k} + (-3)[2\hat{i} + 3\hat{j} + 4\hat{k}]$$

$$\Rightarrow \hat{i} + \hat{j} + \hat{k} \text{ is the required line.}$$

316 (c)

Given that,  $\vec{a} = \hat{i} + \hat{j}$  and  $\vec{b} = 2\hat{i} - \hat{k}$

Let  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ , then

$$\vec{r} \times \vec{a} = \vec{b} \times \vec{a} \Rightarrow (\vec{r} - \vec{b}) \times \vec{a} = \vec{0}$$

Now,  $\vec{r} - \vec{b} = (x\hat{i} + y\hat{j} + z\hat{k}) - (2\hat{i} - \hat{k})$

$$= (x - 2)\hat{i} + y\hat{j} + (z + 1)\hat{k}$$

$$\therefore (\vec{r} - \vec{b}) \times \vec{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x - 2 & y & z + 1 \\ 1 & 1 & 0 \end{vmatrix} = \vec{0}$$

$$\Rightarrow -(z + 1)\hat{i} + (z + 1)\hat{j} + (x - 2 - y)\hat{k} = \vec{0}$$

On equating the coefficient of  $\hat{i}, \hat{j}$  and  $\hat{k}$ , we get

$$z = -1, x - y = 2 \dots(i)$$

Now,  $\vec{r} \times \vec{b} = \vec{a} \times \vec{b} \Rightarrow (\vec{r} - \vec{a}) \times \vec{b} = \vec{0}$

And  $\vec{r} - \vec{a} = (x - 1)\hat{i} + (y - 1)\hat{j} + z\hat{k}$

$$\therefore (\vec{r} - \vec{a}) \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x - 1 & y - 1 & z \\ 2 & 0 & -1 \end{vmatrix} = \vec{0}$$

$$\Rightarrow (-y + 1)\hat{i} - \hat{j}(-x + 1 - 2z) + (-2y + 2)\hat{k} = \vec{0}$$

$$\Rightarrow y = 1, x + 2z = 1 \dots(ii)$$

From Eqs. (i) and (ii), we get

$$x = 3, y = 1, z = -1$$

$$\therefore \vec{r} = 3\hat{i} + \hat{j} - \hat{k}$$

317 (a)

Given,  $\vec{A} \times (\vec{B} \times \vec{C}) = \vec{B} \times (\vec{C} \times \vec{A}) \dots (i)$

Also,  $[\vec{A} \vec{B} \vec{C}] \neq 0$  i.e.  $\vec{A}, \vec{B}, \vec{C}$  are not coplanar.

$\therefore$  From Eq. (i)

$$(\vec{A} \cdot \vec{C}) - (\vec{A} \cdot \vec{B})\vec{C} = (\vec{B} \cdot \vec{A})\vec{C} - (\vec{B} \cdot \vec{C})\vec{A}$$

$$\Rightarrow (\vec{B} \cdot \vec{C})\vec{A} + (\vec{A} \cdot \vec{C})\vec{B} - [(\vec{A} \cdot \vec{B}) + (\vec{B} \cdot \vec{C})]\vec{C} = \vec{0}$$

$$\Rightarrow \vec{B} \cdot \vec{C} = \vec{A} \cdot \vec{C} = \vec{A} \cdot \vec{B} = 0$$

$$[\because [\vec{A} \vec{B} \vec{C}] \neq 0]$$

Now, consider

$$\vec{A} \times (\vec{B} \times \vec{C}) = (\vec{A} \cdot \vec{C})\vec{B} - (\vec{A} \cdot \vec{B})\vec{C}$$

$$= 0 \cdot \vec{B} - 0 \cdot \vec{C} = \vec{0}$$

319 (a)

$$[\vec{a} \vec{b} \vec{c}] = \begin{vmatrix} 1 & 0 & -1 \\ x & 1 & 1 - x \\ y & x & 1 + x - y \end{vmatrix}$$

Applying  $C_3 \rightarrow C_3 + C_1$

$$= \begin{vmatrix} 1 & 0 & 0 \\ x & 1 & 1 \\ y & x & 1 + x \end{vmatrix} = 1[1 + x - x] = 1$$

Hence,  $[\vec{a} \vec{b} \vec{c}]$  does not depend upon neither  $x$  nor  $y$ .

320 (b)

The required vector is given by

$$\hat{n} = \frac{\vec{A}\vec{B} \times \vec{A}\vec{C}}{|\vec{A}\vec{B} \times \vec{A}\vec{C}|} = \frac{\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}}{|\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|}$$

321 (d)

$$(\vec{a} - \vec{b}) \cdot (\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})$$

$$= (\vec{a} - \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}]$$

$$= \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) - \vec{b} \cdot (\vec{b} \times \vec{c}) - \vec{b} \cdot (\vec{b} \times \vec{a}) - \vec{b} \cdot (\vec{c} \times \vec{a})$$

$$= \vec{a} \cdot (\vec{b} \times \vec{c}) - \vec{a} \cdot (\vec{b} \times \vec{c})$$

$$= [\vec{a} \vec{b} \vec{c}] - [\vec{a} \vec{b} \vec{c}] = 0$$

322 (b)

$\therefore \vec{a}, \vec{b}$  and  $\vec{c}$  are coplanar vectors, so  $2\vec{a} - \vec{b} - 2\vec{b} - \vec{c}$  and  $2\vec{c} - \vec{a}$  are also coplanar. Thus

$$[2\vec{a} - \vec{b} - 2\vec{b} - \vec{c} \quad 2\vec{c} - \vec{a}] = 0$$

323 (b)

Clearly, angle between  $\vec{a}$  and  $\vec{b} = \frac{\pi}{2}$

$$\Rightarrow \vec{a} \cdot \vec{b} = 0$$

$$\therefore |\vec{a} + \vec{b}|^2 = a^2 + b^2 + 2\vec{a} \cdot \vec{b}$$

$$= 1 + 1 + 0 = 2$$

$$\Rightarrow |\vec{a} + \vec{b}| = \sqrt{2}$$

325 (d)

Given,  $(\vec{a} \times \vec{b}) \times \vec{c} = -\frac{1}{4}|\vec{b}||\vec{c}|\vec{a}$

$$\Rightarrow (\vec{c} \cdot \vec{a})\vec{b} - (\vec{c} \cdot \vec{b})\vec{a} = -\frac{1}{4}|\vec{b}||\vec{c}|\vec{a}$$

On comparing both sides, we get

$$(\vec{c} \cdot \vec{a})\vec{b} = 0$$

$$|\vec{c}||\vec{a}| \cos \theta = 0$$

$$\Rightarrow \theta = \frac{\pi}{2}$$

326 (c)

$$\text{Now, } (\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} + \hat{j}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix}$$

$$= \hat{i}(-1) + \hat{j}(1) + \hat{k}(0) = -\hat{i} + \hat{j}$$

$$\text{and } |(\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} + \hat{j})| = \sqrt{1^2 + 1^2} = \sqrt{2}$$

Vector perpendicular to both of the vectors  $\hat{i} + \hat{j} + \hat{k}$  and  $\hat{i} + \hat{j}$

$$= \frac{(\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} + \hat{j})}{|(\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} + \hat{j})|}$$

$$= \frac{-\hat{i} + \hat{j}}{\sqrt{2}} = \frac{-1}{\sqrt{2}}(\hat{i} - \hat{j})$$

$$= c(\hat{i} - \hat{j}), c \text{ is a scalar.}$$

327 (b)

It is given that  $(\vec{a} + \vec{b}) \parallel \vec{c}$  and  $(\vec{c} + \vec{a}) \parallel \vec{b}$

$$\therefore (\vec{a} + \vec{b}) \times \vec{c} = 0 \text{ and } (\vec{c} + \vec{a}) \times \vec{b} = 0$$

$$\Rightarrow \vec{a} \times \vec{c} + \vec{b} \times \vec{c} = 0 \text{ and } \vec{c} \times \vec{b} + \vec{a} \times \vec{b} = 0$$

$$\Rightarrow \vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$$

Hence,  $\vec{a}, \vec{b}, \vec{c}$  form the sides of a triangle

328 (a)

$$\therefore \text{Displacement, } \vec{AB} = (3 - 2)\hat{i} + (1 + 1)\hat{j} + (2 - 1)\hat{k}$$

$$= \hat{i} + 2\hat{j} + \hat{k}$$

$$\text{and force, } \vec{F} = \frac{\sqrt{6}(\hat{i} + 2\hat{j} + \hat{k})}{\sqrt{6}}$$

$$= (\hat{i} + 2\hat{j} + \hat{k})$$

$$\therefore \text{Work done} = \vec{F} \cdot \vec{AB} = (1 + 2\hat{j} + \hat{k}) \cdot (\hat{i} + 2\hat{j} + \hat{k}) = 6$$

329 (c)

let  $\vec{a} = l\hat{i} + m\hat{j} + n\hat{k}$  makes an angle  $\frac{\pi}{4}$  with z-axis

$$\text{Also, } l^2 + m^2 + n^2 = 1$$

$$\text{Here, } n = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}, \quad l^2 + m^2 = \frac{1}{2} \quad \dots \dots (i)$$

$$\therefore \vec{a} = l\hat{i} + m\hat{j} + \frac{\hat{k}}{\sqrt{2}}$$

$$\Rightarrow \vec{a} + \hat{i} + \hat{j} = (l + 1)\hat{i} + (m + 1)\hat{j} + \frac{\hat{k}}{\sqrt{2}}$$

$$\Rightarrow |\vec{a} + \hat{i} + \hat{j}| = \sqrt{(l + 1)^2 + (m + 1)^2 + \left(\frac{1}{\sqrt{2}}\right)^2}$$

$$\Rightarrow 1 = l^2 + m^2 + 2 + 2l + 2m + \frac{1}{2}$$

$$\Rightarrow l + m = -1 \text{ (From Eq. (i))}$$

$$\Rightarrow l^2 + m^2 + 2lm = 1$$

$$\Rightarrow 2lm = \frac{1}{2}$$

$$\Rightarrow l = m = -\frac{1}{2}$$

$$\left( \because l = m = \frac{1}{2}, \text{ is not satisfied the given equation} \right)$$

$$\therefore \vec{a} = -\frac{\hat{i}}{2} - \frac{\hat{j}}{2} + \frac{\hat{k}}{\sqrt{2}}$$

330 (b)

$$\text{Given, } |\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 144$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 (\sin^2\theta + \cos^2\theta) = 144$$

$$\Rightarrow 16|\vec{b}|^2 = 144$$

$$\Rightarrow |\vec{b}| = 3$$

331 (c)

Since,  $m\vec{a}$  is a unit vector, if and only, if

$$|m\vec{a}| = 1 \Rightarrow |m| |\vec{a}| = 1 \Rightarrow m|\vec{a}| = 1$$

$$\Rightarrow m = \frac{1}{|\vec{a}|}$$

332 (b)

Resultant force  $\vec{F}$  is given by

$$\vec{F} = (2\hat{i} - 5\hat{j} + 6\hat{k}) - (-\hat{i} + 2\hat{j} - \hat{k}) = \hat{i} - 3\hat{j} + 5\hat{k}$$

Let  $\vec{d}$  be the displacement vector. Then,

$$\vec{d} = A\vec{B}$$

$$\Rightarrow \vec{d} = (6\hat{i} + \hat{j} - 3\hat{k}) - (4\hat{i} - 3\hat{j} - 2\hat{k}) = 2\hat{i} + 4\hat{j} - \hat{k}$$

$$\therefore W = \text{Work done}$$

$$\Rightarrow W = \vec{F} \cdot \vec{d}$$

$$\Rightarrow W = (\hat{i} - 3\hat{j} + 5\hat{k}) \cdot (2\hat{i} + 4\hat{j} - \hat{k})$$

$$\Rightarrow W = 2 - 12 - 5 = -15 \text{ units}$$

333 (d)

Since,  $P, Q, R$  are collinear. Therefore,

$$\vec{PQ} = m \vec{QR} \text{ for same scalar } m$$

$$\Rightarrow -2\hat{j} = m[(a - 1)\hat{i} + (\vec{b} + 1)\hat{j} + c\hat{k}] \text{ for some non-zero scalar } m$$

$$\Rightarrow (a - 1)m = 0, (b + 1)m = -2, cm = 0$$

$$\Rightarrow a = 1, c = 0, b \in R$$

334 (b)

The direction cosines of a vector making equal angles with the coordinate axes are  $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$

Therefore, the unit vector along the vector making equal angles with the coordinate axes is

$$\vec{b} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k}$$

$$\therefore \text{Projection of } \vec{a} \text{ on } \vec{b} = \vec{a} \cdot \vec{b}$$

$$= (4\hat{i} - 3\hat{j} + 2\hat{k}) \cdot \left( \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k} \right) = \frac{4 - 3 + 2}{\sqrt{3}} = \sqrt{3}$$

335 (a)

$$[2\hat{i} 3\hat{j} - 5\hat{k}]$$

$$= -30 [\hat{i} \hat{j} \hat{k}]$$

$$= -30 (\because [\hat{i} \hat{j} \hat{k}] = 1)$$

336 (b)

We have,

$$\begin{aligned} & (\vec{a} \times \vec{b}) \times (\vec{a} \times \vec{c}) \cdot \vec{d} \\ &= \{((\vec{a} \times \vec{b}) \cdot \vec{c}) \vec{a} - ((\vec{a} \times \vec{b}) \cdot \vec{a}) \vec{c}\} \cdot \vec{d} \\ &= \{[\vec{a} \vec{b} \vec{c}] \vec{a} - 0\} \cdot \vec{d} = [\vec{a} \vec{b} \vec{c}] (\vec{a} \cdot \vec{d}) \end{aligned}$$

337 (d)

$$\text{Resultant force } \vec{F} = (2\hat{i} - 5\hat{j} + 6\hat{k}) + (-\hat{i} + 2\hat{j} - \hat{k})$$

$$= \hat{i} - 3\hat{j} + 5\hat{k}$$

$$\text{and displacement, } \vec{d} = (6\hat{i} + \hat{j} - 3\hat{k}) - (4\hat{i} - 3\hat{j} - 2\hat{k})$$

$$= 2\hat{i} + 4\hat{j} - \hat{k}$$

$$\therefore \text{work done } W = \vec{F} \cdot \vec{d}$$

$$= (\hat{i} - 3\hat{j} + 5\hat{k}) \cdot (2\hat{i} + 4\hat{j} - \hat{k})$$

$$= -15$$

$$= 15 \text{ units [neglecting - ve sign]}$$

338 (a)

The resultant force is given by

$$\begin{aligned} \vec{F} &= 6 \frac{(\hat{i} - 2\hat{j} + 2\hat{k})}{\sqrt{1+4+4}} + 7 \frac{(2\hat{i} - 3\hat{j} - 6\hat{k})}{\sqrt{4+9+36}} \\ &= 4\hat{i} - 7\hat{j} - 2\hat{k} \end{aligned}$$

$$\vec{d} = \text{Displacement} = \vec{PQ}$$

$$\vec{d} = (5\hat{i} - \hat{j} + \hat{k}) - (2\hat{i} - \hat{j} - 3\hat{k}) = 3\hat{i} + 4\hat{k}$$

$$\therefore \text{Work done} = \vec{F} \cdot \vec{d} = 12 + 0 - 8 = 4 \text{ units}$$

339 (c)

$$\text{We know, } [\vec{b} \times \vec{c} \vec{c} \times \vec{a} \vec{a} \times \vec{b}]$$

$$= (\vec{b} \times \vec{c}) \cdot [(\vec{c} \times \vec{a}) \cdot (\vec{a} \times \vec{b})]$$

$$= (\vec{b} \times \vec{c}) \cdot [((\vec{c} \times \vec{a}) \cdot \vec{b}) \vec{a} - ((\vec{c} \times \vec{a}) \cdot \vec{a}) \vec{b}]$$

$$= (\vec{b} \times \vec{c}) \cdot ([\vec{c} \vec{a} \vec{b}] \vec{a} - [\vec{c} \vec{a} \vec{a}] \vec{b})$$

$$= (\vec{b} \times \vec{c}) \cdot \vec{a} [\vec{a} \vec{b} \vec{c}] - 0$$

$$= [\vec{a} \vec{b} \vec{c}] [\vec{a} \vec{b} \vec{c}]$$

$$= [\vec{a} \vec{b} \vec{c}]^2$$

340 (d)

$$\therefore \vec{QP} \text{ is parallel to } \vec{AB} \text{ and } \vec{DC}.$$

$$\therefore \vec{AB} + \vec{DC} = \vec{QP} + \vec{QP} = 2\vec{QP}$$

341 (a)

Taking A as the origin, let the position vectors of B and C be  $\vec{b}$  and  $\vec{c}$  respectively

$$\begin{aligned} \therefore \vec{BE} + \vec{AF} &= \left(\frac{\vec{c}}{2} - \vec{b}\right) + \left(\frac{\vec{b} + \vec{c}}{2} - \vec{0}\right) = \vec{c} - \frac{\vec{b}}{2} \\ &= \vec{DC} \end{aligned}$$

342 (a)

Since,  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular unit vectors.

$$\Rightarrow |\vec{a}| = |\vec{b}| = |\vec{c}| = 1$$

$$\text{and } \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0 \quad \dots(i)$$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c})$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$= 1 + 1 + 1 + 0 = 3 \quad [\text{from Eq. (i)}]$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}$$

343 (c)

Any vector lying in the plane of  $\vec{a}$  and  $\vec{b}$  is of the form  $x\vec{a} + y\vec{b}$

It is given that  $\vec{c}$  is parallel to the plane of  $\vec{a}$  and  $\vec{b}$

$$\therefore \vec{c} = \lambda(x\vec{a} + y\vec{b}) \text{ for some scalar } \lambda$$

$$\Rightarrow d\hat{i} + \hat{j} + (2d - 1)\hat{k}$$

$$= \lambda\{x(\hat{i} - 2\hat{j} + 3\hat{k})$$

$$+ y(3\hat{i} + 3\hat{j} - \hat{k})\}$$

$$\Rightarrow d\hat{i} + \hat{j} + (2d - 1)\hat{k}$$

$$= \lambda\{(x + 3y)\hat{i} + (-2x + 3y)\hat{j}$$

$$+ (3x - y)\hat{k}\}$$

$$\Rightarrow \lambda(x + 3y) = d, \lambda(-2x + 3y) = 1 \text{ and } \lambda(3x - y) = (2d - 1)$$

[ $\therefore \hat{i}, \hat{j}, \hat{k}$  are non-coplanar]

Solving  $\lambda(x + 3y) = d$  and  $3x - y = 2d - 1$ , we get

$$x = \frac{7d-3}{10\lambda} \text{ and } y = \frac{d+1}{10\lambda}$$

Substituting these values in  $\lambda(x + 3y) = d$ , we get  $11d = -1$

ALTER clearly,  $\vec{c}$  is perpendicular to  $\vec{a} \times \vec{b}$

$$\therefore \vec{c} \cdot (\vec{a} \times \vec{b}) = 0$$

$$\Rightarrow [\vec{c} \vec{a} \vec{b}] = 0 \Rightarrow \begin{vmatrix} d & 1 & 2d - 1 \\ 1 & -2 & 3 \\ 3 & 3 & -1 \end{vmatrix} = 0 \Rightarrow 11d = -1$$

344 (c)

$\therefore \vec{p}, \vec{q}, \vec{r}$  are reciprocal vectors  $\vec{a}, \vec{b}, \vec{c}$  respectively.

$$\therefore \vec{p} \cdot \vec{a} = 1, \vec{p} \cdot \vec{b} = 0, \vec{p} \cdot \vec{c} \text{ etc.}$$

$$\therefore (l\vec{a} + m\vec{b} + n\vec{c}) \cdot (l\vec{p} + m\vec{q} + n\vec{r}) = l^2 + m^2 + n^2$$

345 (b)

$$\text{Given expression} = 2(1 + 1 + 1) - 2\sum(\vec{a} \cdot \vec{b})$$

$$= 6 - 2\sum(\vec{a} \cdot \vec{b}) \quad \dots(i)$$

$$\text{But } (\vec{a} + \vec{b} + \vec{c})^2 \geq 0$$

$$\therefore (1 + 1 + 1) + 2\sum \vec{a} \cdot \vec{b} \geq 0$$

$$\therefore 3 \geq -2\sum \vec{a} \cdot \vec{b} \quad \dots(ii)$$

From relations (i) and (ii), we get

$$\text{Given expression} \leq 6 + 3 = 9$$

346 (a)

Let  $\vec{OA} = \hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{OB} = 3\hat{i} + 4\hat{j} + 5\hat{k}$

$$\therefore \vec{AB} = 2\hat{i} + 2\hat{j} + 2\hat{k}$$

$\therefore$  work done,  $W = \vec{F} \cdot \vec{AB}$

$$= (2\hat{i} - 3\hat{j} + 2\hat{k}) \cdot (2\hat{i} + 2\hat{j} + 2\hat{k})$$

$$= 4 - 6 + 4 = 2$$

347 (d)

$$\vec{AC} = (a\hat{i} - 3\hat{j} + \hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = (a-2)\hat{i} - 2\hat{j}$$

$$\text{and } \vec{BC} = (a\hat{i} - 3\hat{j} + \hat{k}) - (\hat{i} - 3\hat{j} - 5\hat{k}) =$$

$$(a-1)\hat{i} + 6\hat{k}$$

Since, the  $\Delta ABC$  is right angled at  $C$ , then

$$\vec{AC} \cdot \vec{BC} = 0$$

$$\Rightarrow \{(a-2)\hat{i} - 2\hat{j}\} \cdot \{(a-1)\hat{i} + 6\hat{k}\} = 0$$

$$\Rightarrow (a-2)(a-1) = 0 \Rightarrow a = 1 \text{ and } 2$$

348 (a)

We have,

$$(\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$$

$$\Leftrightarrow -\vec{c} \times (\vec{a} \times \vec{b}) = \vec{a} \times (\vec{b} \times \vec{c})$$

$$\Leftrightarrow -\{(\vec{c} \cdot \vec{b})\vec{a} - (\vec{c} \cdot \vec{a})\vec{b}\} = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$$

$$\Leftrightarrow (\vec{a} \cdot \vec{b})\vec{c} - (\vec{c} \cdot \vec{b})\vec{a} = 0$$

$$\Leftrightarrow (\vec{b} \cdot \vec{a})\vec{c} - (\vec{b} \cdot \vec{c})\vec{a} = 0$$

$$\Leftrightarrow \vec{b} \times (\vec{c} \times \vec{a}) = 0$$

349 (b)

Clearly,

$$(\vec{a} + \vec{b}) \times \{\vec{c} - (\vec{a} + \vec{b})\}$$

$$= (\vec{a} + \vec{b}) \times \vec{c} - (\vec{a} + \vec{b}) \times (\vec{a} + \vec{b}) = (\vec{a} + \vec{b}) \times \vec{c}$$

350 (a)

$$\vec{PQ} = (2\hat{i} - \hat{j} + 3\hat{k}) - (\hat{i} - \hat{j} + 2\hat{k})$$

$$= \hat{i} + \hat{k}$$

$$\text{and } \vec{F} = 3\hat{i} + 2\hat{j} - 4\hat{k}$$

$$\therefore \text{Moment} = |\vec{PQ} \times \vec{F}|$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & 1 \\ 3 & 2 & -4 \end{vmatrix}$$

$$= -2\hat{i} + 7\hat{j} + 2\hat{k}$$

$$\therefore \text{Magnitude of moment} = \sqrt{4 + 49 + 4} = \sqrt{57}$$

351 (b)

$$\text{Since, } |\vec{a} + \vec{b}| = \sqrt{3}$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = 3$$

$$\Rightarrow \vec{a} \cdot \vec{b} = \frac{1}{2} \dots (i)$$

$$\because [|\vec{a}| = |\vec{b}| = 1, \text{ given}]$$

$$\therefore (3\vec{a} - 4\vec{b}) \cdot (2\vec{a} + 5\vec{b}) = 6 + 7\vec{a} \cdot \vec{b} - 20$$

$$= 6 + \frac{7}{2} - 20$$

$$= -\frac{21}{2} \quad [\text{from Eq. (i)}]$$

352 (c)

We have,

$$\hat{a} \times (\hat{b} \times \hat{c}) = \frac{1}{2}\hat{b}$$

$$\Rightarrow (\hat{a} \cdot \hat{c})\hat{b} - (\hat{a} \cdot \hat{b})\hat{c} = \frac{1}{2}\hat{b}$$

$$\Rightarrow \left\{(\hat{a} \cdot \hat{c}) - \frac{1}{2}\right\}\hat{b} - (\hat{a} \cdot \hat{b})\hat{c} = 0$$

$$\Rightarrow \hat{a} \cdot \hat{c} - \frac{1}{2} = 0 \text{ and } \hat{a} \cdot \hat{b} = 0$$

$$= 0 \quad \left[ \begin{array}{l} \because \hat{b}, \hat{c} \\ \text{are non-collinear vectors} \end{array} \right]$$

$$\Rightarrow \cos \theta = \frac{1}{2}, \text{ where } \theta \text{ is the angle between } \hat{a} \text{ and } \hat{c}$$

$$\Rightarrow \theta = \pi/3$$

354 (b)

The given line is parallel to the vector  $\vec{n}$

$$= \hat{i} - \hat{j}$$

$$+ 2\hat{k}. \text{ The required plane passing}$$

through the point  $(2, 3, 1)$  i.e.,  $2\hat{i} + 3\hat{j}$

+  $\hat{k}$  and is perpendicular to the vector

$$\vec{n} = \hat{i} - \hat{j} + 2\hat{k}$$

$\therefore$  Its equation is

$$[(\vec{r} - (2\hat{i} + 3\hat{j} + \hat{k})) \cdot (\hat{i} - \hat{j} + 2\hat{k})] = 0$$

$$\Rightarrow \vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 1$$

355 (c)

$$(\vec{a} - \vec{b}) \cdot (\vec{b} \times \vec{c} - \vec{b} \times \vec{a} + \vec{c} \times \vec{a})$$

$$= \vec{a} \cdot (\vec{b} \times \vec{c}) - \vec{b} \cdot (\vec{c} \times \vec{a})$$

$$= [\vec{a} \vec{b} \vec{c}] - [\vec{b} \vec{c} \vec{a}] = 0$$

356 (a)

We have,

$$|\hat{n}_1 + \hat{n}_2|^2 = |\hat{n}_1|^2 + |\hat{n}_2|^2 + 2\hat{n}_1 \cdot \hat{n}_2$$


$$\Rightarrow |\hat{n}_1 + \hat{n}_2|^2 = |\hat{n}_1|^2 + |\hat{n}_2|^2 + 2|\hat{n}_1||\hat{n}_2|\cos \theta$$

$$\Rightarrow |\hat{n}_1 + \hat{n}_2|^2 = 1 + 1 + 2\cos \theta = 4\cos^2 \frac{\theta}{2}$$

$$\therefore \cos \frac{\theta}{2} = \frac{1}{2} |\hat{n}_1 + \hat{n}_2|$$

357 (d)

$$\text{Let } \vec{R}_1 = 2\hat{i} + 4\hat{j} - 5\hat{k}$$



$$\text{and } \vec{R}_2 = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\therefore \vec{R} \text{ (along } \vec{AC}) = \vec{R}_1 + \vec{R}_2$$

$$= 3\hat{i} + 6\hat{j} - 2\hat{k}$$

$$\therefore \vec{a} \text{ (unit vector along } AC) = \frac{\vec{R}}{|\vec{R}|}$$

$$= \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{9 + 36 + 4}}$$

$$= \frac{1}{7}(3\hat{i} + 6\hat{j} - 2\hat{k})$$

358 (a)

Let  $P(60\hat{i} + 3\hat{j})$ ,  $Q(40\hat{i} - 8\hat{j})$  and  $R(a\hat{i} - 52\hat{j})$  be the collinear points. Then  $\overrightarrow{PQ} = \lambda\overrightarrow{QR}$  for some scalar  $\lambda$

$$\Rightarrow (-20\hat{i} - 11\hat{j}) = \lambda[(a - 40)\hat{i} - 44\hat{j}]$$

$$\Rightarrow \lambda(a - 40) = -20, -44\lambda = -11$$

$$\Rightarrow \lambda(a - 40) = -20, \lambda = \frac{1}{4}$$

$$\therefore a - 40 = -20 \times 4 \Rightarrow a = -40$$

359 (a)

We have,

$$\vec{a} + \vec{b} + \vec{c} = \alpha\vec{d} \text{ and } \vec{b} + \vec{c} + \vec{d} = \beta\vec{a}$$

$$\Rightarrow \vec{a} + \vec{b} + \vec{c} + \vec{d} = (\alpha + 1)\vec{d} \text{ and } \vec{a} + \vec{b} + \vec{c} + \vec{d} = (\beta + 1)\vec{a}$$

$$\Rightarrow (\alpha + 1)\vec{d} = (\beta + 1)\vec{a}$$

$$\Rightarrow (\alpha + 1)\vec{d} = (\beta + 1)\vec{a}$$

If  $\alpha \neq -1$ , then

$$(\alpha + 1)\vec{d} = (\beta + 1)\vec{a} \Rightarrow \vec{d} = \frac{\beta + 1}{\alpha + 1}\vec{a}$$

$$\therefore \vec{a} + \vec{b} + \vec{c} = \alpha\vec{d}$$

$$\Rightarrow \vec{a} + \vec{b} + \vec{c} = \alpha \left( \frac{\beta + 1}{\alpha + 1} \right) \vec{a}$$

$$\Rightarrow \left\{ 1 - \frac{\alpha(\beta + 1)}{\alpha + 1} \right\} \vec{a} + \vec{b} + \vec{c} = 0$$

$$\Rightarrow \vec{a}, \vec{b}, \vec{c} \text{ are coplanar}$$

It is a contradiction to the given condition

$$\therefore \alpha = -1 \Rightarrow \vec{a} + \vec{b} + \vec{c} = 0$$

360 (c)

Let the unit vector  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$  is perpendicular to  $\hat{i} - \hat{j}$ , then we get

$$(\hat{i} + \hat{j}) \cdot (\hat{i} - \hat{j}) = 0$$

$$\frac{(\hat{i} + \hat{j}) \cdot (\hat{i} - \hat{j})}{\sqrt{2}} = \frac{1 - 1}{\sqrt{2}} = 0$$

$$\therefore \frac{\hat{i} + \hat{j}}{\sqrt{2}} \text{ is the unit vector}$$

361 (c)

We have,

$$\left. \begin{aligned} \vec{r} \cdot \vec{a} = 0 &\Rightarrow \vec{r} \perp \vec{a} \\ \vec{r} \cdot \vec{b} = 0 &\Rightarrow \vec{r} \perp \vec{b} \\ \vec{r} \cdot \vec{c} = 0 &\Rightarrow \vec{r} \perp \vec{c} \end{aligned} \right\} \Rightarrow \vec{a}, \vec{b}, \vec{c} \text{ are coplanar}$$

$$\text{Hence, } [\vec{a}\vec{b}\vec{c}] = 0$$

362 (b)

$$\cos \frac{\pi}{3} = \frac{(\hat{i} + \hat{k}) \cdot (\hat{i} + \hat{j} + a\hat{k})}{\sqrt{2}\sqrt{1 + 1 + a^2}}$$

$$\Rightarrow \frac{1}{2} = \frac{1 + a}{\sqrt{2}\sqrt{2 + a^2}}$$

$$\Rightarrow \frac{1}{4} = \frac{(1 + a)^2}{2(2 + a^2)}$$

$$\Rightarrow 2 + a^2 = 2(1 + a^2 + 2a)$$

$$\Rightarrow a^2 + 4a = 0$$

$$\Rightarrow a = 0, -4$$

363 (b)

Let the required vector be  $\vec{d} = x\hat{i} + y\hat{j} + z\hat{k}$

It makes equal angles with the unit vectors

$$\vec{b} = \frac{1}{3}(\hat{i} - 2\hat{j} + 2\hat{k}), \vec{c} = \frac{1}{5}(-4\hat{i} - 3\hat{k}) \text{ and } \vec{d} = \hat{j}$$

$$\therefore \vec{d} \cdot \vec{b} = \vec{d} \cdot \vec{c} = \vec{d} \cdot \vec{d} \quad [\because \vec{b}, \vec{c}, \vec{d} \text{ are unit vectors}]$$

$$\Rightarrow \frac{1}{3}(x - 2y + 2z) = \frac{1}{5}(-4x - 3z) = y$$

$$\Rightarrow x - 2y + 2z = 3y \text{ and } -4x - 5y - 3z = 0$$

$$\Rightarrow x - 5y + 2z = 0 \text{ and } 4x + 5y + 3z = 0$$

$$\Rightarrow \frac{x}{-5} = \frac{y}{1} = \frac{z}{5} = \lambda \text{ (say)}$$

$$\Rightarrow x = -5\lambda, y = \lambda, z = 5\lambda \text{ for some scalar } \lambda$$

$$\Rightarrow \vec{d} = \lambda(-5\hat{i} + \hat{j} + 5\hat{k})$$

Clearly, option (b) is true for  $\lambda = 1$

364 (d)

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 2 & 1 \\ 1 & -2 & 2 \end{vmatrix}$$

$$= \hat{i}(4 + 2) - \hat{j}(4 - 1) + \hat{k}(-4 - 2)$$

$$= 6\hat{i} - 3\hat{j} - 6\hat{k}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = \sqrt{36 + 9 + 36} = \sqrt{81} = 9$$

$\therefore$  Required vectors are

$$\pm 6 \frac{|\vec{a} \times \vec{b}|}{|\vec{a} \times \vec{b}|}$$

$$= \pm \frac{6}{9}(6\hat{i} - 3\hat{j} - 6\hat{k})$$

$$= \pm 2(2\hat{i} - \hat{j} - 2\hat{k})$$

366 (d)

(a) Let  $\vec{p} = x\hat{i} + y\hat{j} + z\hat{k}$  where at least one of  $x, y, z$  is non-zero. Let

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

$$\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$$

$\therefore$  By given conditions

$$a_1x + a_2y + a_3z = 0$$

$$b_1x + b_2y + b_3z = 0$$

$$c_1x + c_2y + c_3z = 0$$

$$\Rightarrow \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = 0$$

$$\Rightarrow [\vec{a} \vec{b} \vec{c}] = 0$$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$  are coplanar.

(b) Vectors are coplanar, if

$$\begin{vmatrix} 1 & 3 & 0 \\ 2 & 0 & 1 \\ 0 & 1 & 1 \end{vmatrix} = 0$$

$$\text{ie, } -7 = 0$$

Which is not possible.

$$(c) \vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$$

$\Rightarrow \vec{a} \times (\vec{b} \times \vec{c})$  is coplanar with  $\vec{b}$  and  $\vec{c}$ .

$$(d) |\vec{a}| = |\vec{b}| = 1$$

$$\therefore |\vec{a} + \vec{b}|^2 = (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b})$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$= 1 + 1 + 2 \cdot 1 \cdot 1 \cos \frac{\pi}{3} = 3$$

$$\Rightarrow |\vec{a} + \vec{b}| = \sqrt{3} > 1$$

367 (d)

$$\text{Here, } \vec{a}_1 = 3\hat{i} + 6\hat{j}, \quad \vec{a}_2 = -2\hat{i} + 7\hat{k}$$

$$\vec{b}_1 = -4\hat{i} + 3\hat{j} + 2\hat{k} \text{ and } \vec{b}_2 = -4\hat{i} + \hat{j} + \hat{k}$$

$$\text{Now, } \vec{a}_2 - \vec{a}_1 = \hat{i} - 6\hat{j} + 7\hat{k}$$

and

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -4 & 3 & 2 \\ -4 & 1 & 1 \end{vmatrix} = \hat{i} - 4\hat{j} + 8\hat{k}$$

$$\Rightarrow |\vec{b}_1 \times \vec{b}_2| = \sqrt{1 + 16 + 64} = 9$$

Now,

$$\begin{aligned} (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) \\ = (\hat{i} - 6\hat{j} + 7\hat{k}) \cdot (\hat{i} - 4\hat{j} + 8\hat{k}) \end{aligned}$$

$$= 1 + 24 + 56 = 81$$

$\therefore$  Shortest distance,

$$d = \frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

$$= \frac{81}{9} = 9 \text{ unit}$$

368 (b)

We know that a vector perpendicular to the plane containing the points  $\vec{A}, \vec{B}, \vec{C}$  is given by  $\vec{A} \times \vec{B} + \vec{B} \times \vec{C} + \vec{C} \times \vec{A}$ .

$$\text{Given, } \vec{A} = \hat{i} - \hat{j} + 2\hat{k}, \vec{B} = 2\hat{i} + 0\hat{j} - \hat{k}$$

$$\text{and } \vec{C} = 0\hat{i} + 2\hat{j} + \hat{k}$$

Now,

$$\vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 2 & 0 & -1 \end{vmatrix} = \hat{i} + 5\hat{j} + 2\hat{k}$$

$$\vec{B} \times \vec{C} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & -1 \\ 0 & 2 & 1 \end{vmatrix} = 2\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\vec{C} \times \vec{A} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 2 & 1 \\ 1 & -1 & 2 \end{vmatrix} = 5\hat{i} + \hat{j} - 2\hat{k}$$

Thus,

$$\begin{aligned} \vec{A} \times \vec{B} + \vec{B} \times \vec{C} + \vec{C} \times \vec{A} \\ = (\hat{i} + 5\hat{j} + 2\hat{k}) + (2\hat{i} - 2\hat{j} + 4\hat{k}) + (5\hat{i} + \hat{j} - 2\hat{k}) \\ = 8\hat{i} + 4\hat{j} + 4\hat{k} \end{aligned}$$

369 (c)

Given,

$$(\vec{a} \times \vec{b}) \cdot (\vec{a} \times \vec{b}) = \frac{1}{4}$$

$$\Rightarrow (|\vec{a}||\vec{b}|\sin\theta)^2 = \frac{1}{4}$$

$$\Rightarrow \sin^2\theta = \frac{1}{4}$$

$$\Rightarrow \theta = \frac{\pi}{6}$$

370 (b)

Given that,  $|\vec{a}| = 3, |\vec{b}| = 4$  and  $\vec{a} + \lambda\vec{b}$  is perpendicular to  $\vec{a} - \lambda\vec{b}$ .

$$\therefore (\vec{a} + \lambda\vec{b}) \cdot (\vec{a} - \lambda\vec{b}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{a} - \vec{a} \cdot \lambda\vec{b} + \lambda\vec{b} \cdot \vec{a} - \lambda^2\vec{b} \cdot \vec{b} = 0$$

$$\Rightarrow |\vec{a}|^2 - \lambda^2|\vec{b}|^2 = 0$$

$$\Rightarrow \lambda^2 = \frac{|\vec{a}|^2}{|\vec{b}|^2} \Rightarrow \lambda = \frac{|\vec{a}|}{|\vec{b}|} = \frac{3}{4}$$

371 (a)

$$\begin{aligned} (\vec{x} - \vec{y}) \times (\vec{x} + \vec{y}) \\ = \vec{x} \times \vec{x} + \vec{x} \times \vec{y} - \vec{y} \times \vec{x} - \vec{y} \times \vec{y} \\ = \vec{0} + \vec{x} \times \vec{y} + \vec{x} \times \vec{y} - \vec{0} \\ = 2(\vec{x} \times \vec{y}) \end{aligned}$$

372 (a)

$$\vec{a} \cdot \vec{c} = 0$$

$$\Rightarrow \lambda - 1 + 2\mu = 0$$

$$\Rightarrow \lambda + 2\mu = 1 \dots(i)$$

$$\vec{b} \cdot \vec{c} = 0$$

$$\Rightarrow 2\lambda + 4 + \mu = 0$$

$$\Rightarrow 2\lambda + \mu = -4 \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$\lambda - 3, \mu = 2$$

375 (b)

The projection  $\vec{x} \times \vec{y}$  on  $\vec{z}$  is given by

$$\frac{(\vec{x} \times \vec{y}) \cdot \vec{z}}{|\vec{z}|} = \frac{1}{|\vec{z}|} [\vec{x} \vec{y} \vec{z}] = \frac{1}{13} \begin{vmatrix} 3 & -6 & -1 \\ 1 & 4 & -3 \\ 3 & -4 & -12 \end{vmatrix} = -14$$

376 (c)

We have,

$$\vec{a} \times \{\vec{a} \times \{\vec{a} \times (\vec{a} \times \vec{b})\}\}$$

$$\begin{aligned}
&= \vec{a} \times \left\{ \vec{a} \times \left\{ (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} \right\} \right\} \\
&= \vec{a} \times \left\{ \vec{0} - |\vec{a}|^2(\vec{a} \times \vec{b}) \right\} \\
&= -|\vec{a}|^2 \{ \vec{a} \times (\vec{a} \times \vec{b}) \} = -|\vec{a}|^2 \{ (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} \} \\
&= -|\vec{a}|^2 \{ 0 - |\vec{a}|^2 \vec{b} \} = |\vec{a}|^4 \vec{b}
\end{aligned}$$

379 (c)

For an obtuse angle

$$(cx\hat{i} - 6\hat{j} + 3\hat{k}) \cdot (x\hat{i} + 2\hat{j} + 2cx\hat{k}) < 0$$

$$\Rightarrow cx^2 - 12 + 6cx < 0$$

$$\Rightarrow cx^2 + 6cx - 12 < 0$$

$$\therefore (6c)^2 - 4c(-12) < 0 \quad [\because f(x) < 0 \Rightarrow D < 0]$$

$$\Rightarrow 36c \left( c + \frac{4}{3} \right) < 0$$

$$\Rightarrow -\frac{4}{3} < c < 0$$

380 (a)

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

$$\begin{aligned}
&= \frac{(2\hat{i} + 2\hat{j} - \hat{k}) \cdot (6\hat{i} - 3\hat{j} + 2\hat{k})}{\sqrt{2^2 + 2^2 + (-1)^2} \sqrt{6^2 + (-3)^2 + 2^2}} \\
&= \frac{12 - 6 - 2}{\sqrt{4 + 4 + 1} \sqrt{36 + 9 + 4}} = \frac{4}{21}
\end{aligned}$$

381 (b)

Given vectors  $2\hat{i} + 3\hat{j} - 4\hat{k}$  and  $a\hat{i} + b\hat{j} + c\hat{k}$  will be perpendicular, if

$$(2\hat{i} + 3\hat{j} - 4\hat{k}) \cdot (a\hat{i} + b\hat{j} + c\hat{k}) = 0 \Rightarrow 2a + 3b - c = 0$$

Clearly,  $a = 4, b = 4, c = 5$  satisfy the above equation

382 (a)

We have,  $\vec{\alpha} = x(\vec{a} \times \vec{b}) + y(\vec{b} \times \vec{c}) + z(\vec{c} \times \vec{a})$

Taking dot product with  $\vec{a}, \vec{b}, \vec{c}$  respectively, we get

$$\vec{\alpha} \cdot \vec{a} = y[\vec{a} \cdot \vec{b} \times \vec{c}] \Rightarrow y = 8(\vec{\alpha} \cdot \vec{a})$$

$$\vec{\alpha} \cdot \vec{b} = z[(\vec{c} \times \vec{a}) \cdot \vec{b}]$$

$$\Rightarrow \vec{\alpha} \cdot \vec{b} = z[\vec{a} \cdot \vec{b} \times \vec{c}] \Rightarrow z = 8(\vec{\alpha} \cdot \vec{b})$$

$$\text{and } \vec{\alpha} \cdot \vec{c} = x(\vec{a} \times \vec{b} \cdot \vec{c})$$

$$\vec{\alpha} \cdot \vec{c} = x[\vec{a} \cdot \vec{b} \times \vec{c}] \Rightarrow x = 8(\vec{\alpha} \cdot \vec{c})$$

$$\therefore x + y + z = 8\vec{\alpha} \cdot (\vec{a} + \vec{b} + \vec{c})$$

383 (d)

Let  $\vec{c} = 3\hat{i} + \hat{j} - 5\hat{k}$  and  $\vec{d} = a\hat{i} + b\hat{j} - 15\hat{k}$

$$\text{For collinears, } \vec{c} \times \vec{d} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & -5 \\ a & b & -15 \end{vmatrix} = \vec{0}$$

$$\Rightarrow \hat{i}(-15 + 5b) - \hat{j}(-45 + 5a) + \hat{k}(3b - a) = \vec{0}$$

$$\Rightarrow -15 + 5b = 0, \quad -45 + 5a = 0, \\ 3b - a = 0$$

$$\Rightarrow b = 3, a = 9$$

384 (d)

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|\cos \theta$$

$$= 1^2 + 1^2 - 2 \cdot 1 \cdot 1 \cdot \cos 60^\circ \quad [\because |\vec{a}| = |\vec{b}| = 1]$$

$$= 2 - 2 \cdot \frac{1}{2} = 1$$

385 (c)

Let  $\vec{a} = \hat{i} - 2\hat{j} - 3\hat{k}$ ,  $\vec{b} = 2\hat{i} + 0\hat{j} + 0\hat{k}$

Now take option (c).

Let  $\vec{c} = 0\hat{i} - 4\hat{j} - 6\hat{k}$

$$\text{Now, } \vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} 1 & -2 & -3 \\ 2 & 0 & 0 \\ 0 & -4 & -6 \end{vmatrix}$$

$$= 1(0) + 2(-12) - 3(-8) = 0$$

386 (a)

$$(\vec{a} + \vec{b}) \times (\vec{a} \times \vec{b})$$

$$= \vec{a} \times (\vec{a} \times \vec{b}) + \vec{b} \times (\vec{a} \times \vec{b})$$

$$= (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} + (\vec{b} \cdot \vec{b})\vec{a} - (\vec{b} \cdot \vec{a})\vec{b}$$

$$= (\vec{a} \cdot \vec{b})\vec{a} - \vec{b} + \vec{a} - (\vec{b} \cdot \vec{a})\vec{b}$$

$$= (\vec{a} - \vec{b})(\vec{a} \cdot \vec{b} - 1)$$

$\therefore$  Given vector is parallel to  $(\vec{a} - \vec{b})$ .

387 (a)

$$\vec{AB} = (2 - 1)\hat{i} + (0 - 2)\hat{j} + (3 + 1)\hat{k}$$

$$= \hat{i} - 2\hat{j} + 4\hat{k}$$

and

$$\vec{AC} = (3 - 1)\hat{i} + (-1 - 2)\hat{j} + (2 + 1)\hat{k}$$

$$= 2\hat{i} - 3\hat{j} + 3\hat{k}$$

$$\cos \theta = \frac{(\hat{i} - 2\hat{j} + 4\hat{k}) \cdot (2\hat{i} - 3\hat{j} + 3\hat{k})}{\sqrt{1 + 4 + 16} \sqrt{4 + 9 + 9}}$$

$$= \frac{2 + 6 + 12}{\sqrt{21}\sqrt{22}} = \frac{20}{\sqrt{462}}$$

$$\Rightarrow \sqrt{462} \cos \theta = 20$$

388 (c)

$$[\vec{u} \vec{v} \vec{w}] = |\vec{u} \cdot (\vec{v} \times \vec{w})|$$

$$= |\vec{u} \cdot (3\hat{i} - 7\hat{j} - \hat{k})|$$

$$= |\vec{u}| \sqrt{59} \cos \theta$$

$\therefore$  Maximum value of  $[\vec{u} \vec{v} \vec{w}] = \sqrt{59}$  ( $\because |\vec{u}| = 1, \cos \theta \leq 1$ )

390 (b)

$$\text{Given, force} = 5 \left( \frac{2\hat{i} - 2\hat{j} + \hat{k}}{|2\hat{i} - 2\hat{j} + \hat{k}|} \right) = \frac{5}{3} (2\hat{i} - 2\hat{j} + \hat{k})$$

$$\text{Displacement} = (5\hat{i} + 3\hat{j} + 7\hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k})$$

$$= (4\hat{i} + \hat{j} + 4\hat{k})$$

$\therefore$  Required work done = Force  $\cdot$  Displacement

$$= \frac{5}{3} [(2\hat{i} - 2\hat{j} + \hat{k}) \cdot (4\hat{i} + \hat{j} + 4\hat{k})]$$

$$= \frac{5}{3} [8 - 2 + 4] = \frac{50}{3} \text{ unit}$$

391 (b)

We know that the equation of the plane passing through three non-collinear points  $\vec{a}, \vec{b}, \vec{c}$  is

$$\vec{r} \cdot (\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}) = [\vec{a} \vec{b} \vec{c}]$$

392 (a)

We have,

Required vector  $\vec{r} = \lambda(\hat{a} + \hat{b})$ ,  $\lambda$  is a scalar

$$\begin{aligned} \Rightarrow \vec{r} &= \lambda \left\{ \frac{1}{9}(7\hat{i} - 4\hat{j} - 4\hat{k}) + \frac{1}{3}(-2\hat{i} - \hat{j} + 2\hat{k}) \right\} \\ &= \frac{\lambda}{9}(\hat{i} - 7\hat{j} + 2\hat{k}) \end{aligned}$$

Now,

$$\begin{aligned} |\vec{r}| &= 3\sqrt{6} \Rightarrow |\vec{r}|^2 = 54 \Rightarrow \frac{\lambda^2}{81}(1 + 49 + 4) = 54 \\ &\Rightarrow \lambda = \pm 9 \end{aligned}$$

Hence, required vector  $\vec{r} = \pm(\hat{i} - 7\hat{j} + 2\hat{k})$

Clearly, option (a) is true for  $\lambda = 1$

393 (b)

Given vectors are collinear, if  $\begin{vmatrix} 2 & 1 & 1 \\ 6 & -1 & 2 \\ 14 & -5 & p \end{vmatrix} = 0$

$$\Rightarrow 2[-p + 10] - 1[6p - 28] + 1[-30 + 14] = 0$$

$$\Rightarrow -8p + 32 = 0$$

$$\Rightarrow p = 4$$

394 (d)

Given,

$$\frac{1}{3}|\vec{b}||\vec{c}||\vec{a}| = (\vec{a} \times \vec{b}) \cdot \vec{c}$$

$$\therefore \frac{1}{3}|\vec{b}||\vec{c}||\vec{a}| = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a}$$

On comparing the coefficient of  $\vec{a}$  and  $\vec{b}$ , we get

$$\frac{1}{2}|\vec{b}||\vec{c}| = -\vec{b} \cdot \vec{c} \text{ and } \vec{a} \cdot \vec{c} = 0$$

$$\Rightarrow \frac{1}{3}|\vec{b}||\vec{c}| = -|\vec{b}||\vec{c}|\cos\theta \Rightarrow \cos\theta = -\frac{1}{3}$$

$$\Rightarrow 1 - \sin^2\theta = \frac{1}{9} \Rightarrow \sin\theta = \frac{2\sqrt{2}}{3}$$

395 (c)

Let  $\vec{A} = 7\hat{j} + 10\hat{k}$ ,  $\vec{B} = -\hat{i} + 6\hat{j} + 6\hat{k}$  and  $\vec{C} = -4\hat{i} + 9\hat{j} + 6\hat{k}$

Now,  $\vec{AB} = -\hat{i} - \hat{j} - 4\hat{k}$ ,  $\vec{BC} = -3\hat{i} + 3\hat{j}$

and  $\vec{CA} = 4\hat{i} - 2\hat{j} + 4\hat{k}$

Here,  $|\vec{AB}| = |\vec{BC}| = 3\sqrt{2}$  and  $|\vec{CA}| = 6$

Now,  $|\vec{AB}|^2 + |\vec{BC}|^2 = |\vec{AC}|^2$

Hence, the triangle is right angled isosceles triangle.

396 (b)

We know that if  $A$  and  $B$  are two points and  $P$  is any point on  $AB$ . Then,

$m\vec{PA} + n\vec{PB} = (m+n)\vec{PC}$ , where  $C$  divides  $AB$  in the ratio  $n:m$

Here,  $m = n = 1$

$$\therefore \vec{PA} + \vec{PB} = 2\vec{PC}$$

397 (a)

$$\begin{aligned} &(2\vec{a} + 3\vec{b}) \times (5\vec{a} + 7\vec{b}) + \vec{a} \times \vec{b} \\ &= \vec{0} + 14(\vec{a} \times \vec{b}) - 15(\vec{a} \times \vec{b}) + \vec{0} + \vec{a} \times \vec{b} \\ &= \vec{0} \end{aligned}$$

399 (c)

Let  $\vec{OA} = 2\hat{i} - \hat{j} + \hat{k}$ ,  $\vec{OB} = \hat{i} - 3\hat{j} - 5\hat{k}$

and  $\vec{OC} = 3\hat{i} - 4\hat{j} - 4\hat{k}$

$$\therefore a = |\vec{OA}| = \sqrt{6}, b = |\vec{OB}| = \sqrt{35}$$

and  $c = |\vec{OC}| = \sqrt{41}$

$$\therefore \cos A = \frac{b^2 + c^2 + a^2}{2bc}$$

$$= \frac{(\sqrt{35})^2 + (\sqrt{41})^2 - (\sqrt{6})^2}{2\sqrt{35}\sqrt{41}}$$

$$\Rightarrow \cos A = \frac{\sqrt{35}}{\sqrt{41}}$$

$$\Rightarrow \sin^2 A = \frac{35}{41}$$

400 (d)

Let  $\vec{p} \neq \vec{0}$ . Then,

$$\vec{r} \cdot \vec{a} = \vec{r} \cdot \vec{b} = \vec{r} \cdot \vec{c} = 0$$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$  are coplanar, which is a contradiction

Hence,  $\vec{r} = \vec{0}$

401 (c)

Let  $\vec{\alpha} = \lambda\vec{a} + \mu\vec{b} + t\vec{c}$  ... (i)

Now,  $\vec{a} \cdot \vec{p} = \vec{b} \cdot \vec{q} = \vec{c} \cdot \vec{r} = 1$

$$\Rightarrow \vec{\alpha} \cdot \vec{p} = \lambda(\vec{a} \cdot \vec{p}) + 0 + 0$$

$$\Rightarrow \lambda = \vec{\alpha} \cdot \vec{p}$$

Similarly,  $\mu = \vec{\alpha} \cdot \vec{q}$

and  $t = \vec{\alpha} \cdot \vec{r}$

From Eq. (i), we get

$$\vec{\alpha} = (\vec{\alpha} \cdot \vec{p})\vec{a} + (\vec{\alpha} \cdot \vec{q})\vec{b} + (\vec{\alpha} \cdot \vec{r})\vec{c}$$

402 (a)

Since,  $\vec{b} \times \vec{c}$  is a vector perpendicular to  $\vec{b}, \vec{c}$ .

Therefore  $\vec{a} \times (\vec{b} \times \vec{c})$  is a vector again in plane of  $\vec{b}, \vec{c}$ .

403 (c)

$$(\vec{a} \cdot \vec{b})\vec{b} + \vec{b} \times (\vec{a} \times \vec{b})$$

$$= (\vec{a} \cdot \vec{b})\vec{b} + (\vec{b} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{b})\vec{b}$$

$$= \vec{a} \quad [\because |\vec{b}| = 1]$$

404 (d)

$$\begin{aligned} \therefore \sum_{i=1}^n \vec{a}_i &= \vec{0} \\ \therefore \left( \sum_{i=1}^n \vec{a}_i \right) \cdot \left( \sum_{i=1}^n \vec{a}_i \right) &= \sum_{i=1}^n |\vec{a}_i|^2 + 2 \sum_{1 \leq i < j \leq n} \vec{a}_i \cdot \vec{a}_j \\ \Rightarrow 0 &= n + 2 \sum_{1 \leq i < j \leq n} \vec{a}_i \cdot \vec{a}_j \\ \therefore \sum_{1 \leq i < j \leq n} \vec{a}_i \cdot \vec{a}_j &= -\frac{n}{2} \end{aligned}$$

405 (b)

Since, given vectors are perpendicular.

$$\begin{aligned} \therefore (3\hat{i} - 2\hat{j} - 5\hat{k}) \cdot (6\hat{i} - \hat{j} + c\hat{k}) &= 0 \\ \Rightarrow 18 + 2 - 5c &= 0 \Rightarrow c = 4 \end{aligned}$$

406 (d)

Given,  $\vec{a} \times \vec{b} = \vec{0}$  and  $\vec{a} \cdot \vec{b} = 0$

$\Rightarrow \vec{a}$  is parallel to  $\vec{b}$  and  $\vec{a}$  is perpendicular to  $\vec{b}$  which is possible only if

$$\vec{a} = \vec{0} \text{ or } \vec{b} = \vec{0}$$

407 (a)

Let  $\vec{a} = 2\hat{i} + 4\hat{j} - 5\hat{k}$ ,  $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$

First diagonal,  $\vec{a} + \vec{b} = 3\hat{i} + 6\hat{j} - 2\hat{k}$

$$\Rightarrow |\vec{a} + \vec{b}| = 7$$

Second diagonal,  $\vec{a} - \vec{b} = \hat{i} + 2\hat{j} - 8\hat{k}$

$$\Rightarrow |\vec{a} - \vec{b}| = \sqrt{69}$$

408 (b)

Given  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\Rightarrow \vec{a} \times \vec{a} + \vec{a} \times \vec{b} + \vec{a} \times \vec{c} = 0$$

$$\Rightarrow \vec{a} \times \vec{b} = \vec{c} \times \vec{a}$$

Similarly,  $\vec{b} \times \vec{c} = \vec{c} \times \vec{a}$

$$\therefore \vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$$

**Alternate:** Since,  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors and  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ ,

so  $\vec{a}, \vec{b}, \vec{c}$  represent an equilateral triangle.

$$\therefore \vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$$

409 (c)

We have,

$$\begin{aligned} \vec{AB} + \vec{AC} + \vec{AD} + \vec{AE} + \vec{AF} &= \vec{ED} + \vec{AC} + \vec{AD} + \vec{AE} \\ &\quad + \vec{CD} \quad [\because \vec{AB} = \vec{ED} \text{ and } \vec{AF} = \vec{CD}] \\ &= (\vec{AC} + \vec{CD}) + (\vec{AE} + \vec{ED}) + \vec{AD} \\ &= 3\vec{AD} = 6\vec{AG} \quad [\because \vec{AD} = 2\vec{AG}] \end{aligned}$$

410 (c)

I. It is true that non-zero, non-collinear vectors are linearly independent.

II. It is also true that any three coplanar vectors are linearly dependent.

$\therefore$  Both I and II are true.

411 (a)

Let  $\vec{\alpha} = 2\vec{a} - 3\vec{b}, \vec{\beta} = 7\vec{b} - 9\vec{c}$  and  $\vec{\gamma} = 12\vec{c} - 23\vec{a}$

Then,

$$[\vec{\alpha} \vec{\beta} \vec{\gamma}] = \begin{vmatrix} 2 & -3 & 0 \\ 0 & 7 & -9 \\ -23 & 0 & 12 \end{vmatrix} [\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow [\vec{\alpha} \vec{\beta} \vec{\gamma}] = (168 + 3 \times -207) [\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow [\vec{\alpha} \vec{\beta} \vec{\gamma}] = 0 \quad [\because [\vec{a} \vec{b} \vec{c}] = 0]$$

$\Rightarrow \vec{\alpha}, \vec{\beta}, \vec{\gamma}$  are coplanar vectors

412 (b)

Given,  $[\vec{a} + \vec{b} \vec{b} + \vec{c} \vec{c} + \vec{a}] = [\vec{a} \vec{b} \vec{c}]$

$$\Rightarrow 2[\vec{a} \vec{b} \vec{c}] = [\vec{a} \vec{b} \vec{c}]$$

$$= [\vec{a} \vec{b} \vec{c}] = 0$$

Hence,  $\vec{a}, \vec{b}$  and  $\vec{c}$  are coplanar.

413 (c)

Given,  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = \sqrt{37}, |\vec{b}| = 3$ , and  $|\vec{c}| = 4$

Now,  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\Rightarrow \vec{a} = -(\vec{b} + \vec{c})$$

$$\Rightarrow |\vec{a}|^2 = |-(\vec{b} + \vec{c})|^2$$

$$\Rightarrow |\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2 + 2|\vec{b}||\vec{c}|\cos\theta$$

$$= 9 + 16 + 24\cos\theta$$

$$\Rightarrow 37 = 25 + 24\cos\theta$$

$$\Rightarrow 24\cos\theta = 12 \Rightarrow \theta = 60^\circ$$

414 (a)

Let unit vector be  $a\hat{i} + b\hat{j} + c\hat{k}$

$\therefore a\hat{i} + b\hat{j} + c\hat{k}$  is perpendicular to  $\hat{i} + \hat{j} + \hat{k}$ ,

Then  $a + b + c = 0$  .....(i)

Since,  $a\hat{i} + b\hat{j} + c\hat{k}, (\hat{i} + \hat{j} + 2\hat{k}), (\hat{i} + 2\hat{j} + \hat{k})$  are coplanar

$$\therefore \begin{vmatrix} a & b & c \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -3a + b + c = 0 \text{ ....(ii)}$$

From Eqs. (i) and (ii), we get

$$a = 0 \text{ and } c = -b$$

Also,  $a^2 + b^2 + c^2 = 1$

$$\Rightarrow 0 + b^2 + b^2 = 1$$

$$\Rightarrow b = \frac{1}{\sqrt{2}}$$

$$\therefore a\hat{i} + b\hat{j} + c\hat{k} = \frac{1}{\sqrt{2}}\hat{j} - \frac{1}{\sqrt{2}}\hat{k}$$

416 (b)

$$\text{Given, } \vec{OA} = 2\hat{i} - 2\hat{j} + \hat{k}$$

$$\vec{OB} = 5\hat{i} - 4\hat{j} + 4\hat{k}$$

$$\text{and } \vec{OC} = \hat{i} - 2\hat{j} + 4\hat{k}$$

volume of parallelepiped

$$= |\vec{OA} \vec{OB} \vec{OC}|$$

$$= \begin{vmatrix} 2 & -2 & 1 \\ 5 & -4 & 4 \\ 1 & -2 & 4 \end{vmatrix}$$

$$= 2(-16 + 8) + 2(20 - 4) + 1(-10 + 4)$$

$$= 10 \text{ cu units}$$

418 (a)

We have,

$$\vec{a} = \lambda(\vec{b} \times \vec{c}) = \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ -2 & 4 & 1 \end{vmatrix}$$

$$= \lambda(-10\hat{i} - 7\hat{k} + 8\hat{k})$$

Now,

$$\vec{a} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = -6$$

$$\Rightarrow \lambda(-10 + 14 + 8) = -6 \Rightarrow \lambda = -\frac{1}{2}$$

$$\text{Hence, } \vec{a} = -\frac{1}{2}(-10\hat{i} - 7\hat{k} + 8\hat{k}) = 5\hat{i} + \frac{7}{2}\hat{j} - 4\hat{k}$$

419 (c)

The projection of

$$\vec{a} \text{ on } \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$

$$= \frac{(3\hat{i} - \hat{j} + 5\hat{k}) \cdot (2\hat{i} + 3\hat{j} + \hat{k})}{\sqrt{2^2 + 3^2 + 1^2}} = \frac{8}{\sqrt{14}}$$

421 (d)

$$\begin{vmatrix} 7 & -11 & 1 \\ 5 & 3 & -2 \\ 12 & -8 & -1 \end{vmatrix}$$

$$= 7(-3 - 16) + 11(-5 + 24) + 1(-40 - 36)$$

$$= -133 + 209 - 76 = 0$$

$\therefore$  Vectors are collinear.

422 (c)

Let the position vectors of the points A, B, C are

$$\vec{0}, \vec{a} + \vec{b}, \vec{a} - \vec{b} \text{ respectively and } \theta = 90^\circ$$

$$\therefore \text{Area of triangle} = \frac{1}{2} |\vec{AB} \times \vec{AC}|$$

$$= \frac{1}{2} |(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b})|$$

$$= \frac{1}{2} |2\vec{b} \times \vec{a}|$$

$$= |\vec{b}| |\vec{a}| \sin \theta = 3 \times 2 \sin 90^\circ = 6$$

423 (a)

$$\text{We have, } |[\vec{a} \vec{b} \vec{c}]| = V$$

Let  $V_1$  be the volume of the parallelepiped formed by the vectors  $\vec{a}, \vec{\beta}$  and  $\vec{\gamma}$ . Then,

$$V_1 = |[\vec{a} \vec{\beta} \vec{\gamma}]|$$

Now,

$$[\vec{a} \vec{\beta} \vec{\gamma}] = \begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{a} \cdot \vec{b} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{a} \cdot \vec{c} & \vec{b} \cdot \vec{c} & \vec{c} \cdot \vec{c} \end{vmatrix} [\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow [\vec{a} \vec{\beta} \vec{\gamma}] = [\vec{a} \vec{b} \vec{c}]^2 [\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow [\vec{a} \vec{\beta} \vec{\gamma}] = [\vec{a} \vec{b} \vec{c}]^3$$

$$\therefore V_1 = |[\vec{a} \vec{\beta} \vec{\gamma}]| = |[\vec{a} \vec{b} \vec{c}]^3| = V^3$$

424 (a)

Let  $l, m, n$  be the direction cosines of the required vector. As it makes equal angles with X and Y axes

$$\therefore l = m$$

$$\therefore \text{Required vector } \vec{r} = l\hat{i} + m\hat{j} + n\hat{k} = l\hat{i} + l\hat{j} + n\hat{k}$$

$$\text{Now, } l^2 + m^2 + n^2 = 1 \Rightarrow 2l^2 + n^2 = 1 \quad \dots(i)$$

Since,  $\vec{r}$  is perpendicular to  $-\hat{i} + 2\hat{j} + 2\hat{k}$

$$\therefore \vec{r} \cdot (-\hat{i} + 2\hat{j} + 2\hat{k}) = 0 \Rightarrow -l + 2l + 2n = 0 \Rightarrow$$

$$l + 2n = 0 \quad \dots(ii)$$

$$\text{From (i) and (ii), we get } n = \mp \frac{1}{3}, l = \mp \frac{2}{3}$$

$$\text{Hence, } \vec{r} = \frac{1}{3}(\pm 2\hat{i} \pm 2\hat{j} \mp \hat{k}) = \pm \frac{1}{3}(2\hat{i} + 2\hat{j} - \hat{k})$$

425 (a)

Let the required vector be  $\vec{a}$ . Then,  $\hat{i} - \hat{j}, \hat{i} + \hat{j}$  and  $\vec{a}$  form a right handed system

$$\therefore \vec{a} = (\hat{i} - \hat{j}) \times (\hat{i} + \hat{j}) = \hat{k} + \hat{k} = 2\hat{k}$$

$$\text{Hence, the required unit vector } \hat{a} = \frac{\vec{a}}{|\vec{a}|} = \hat{k}$$

426 (b)

$$\vec{p} = x\vec{a} + y\vec{b} + z\vec{c}$$

$$\Rightarrow 3\hat{i} + 2\hat{j} + 4\hat{k} = x(\hat{i} + \hat{j}) + y(\hat{j} + \hat{k}) + z(\hat{i} + \hat{k})$$

$$\Rightarrow 3\hat{i} + 2\hat{j} + 4\hat{k} = (x + z)\hat{i} + (x + y)\hat{j} + (y + z)\hat{k}$$

On comparing both sides the coefficients of  $\hat{i}, \hat{j}, \hat{k}$ , we get

$$x + z = 3 \quad \dots (i)$$

$$x + y = 2 \quad \dots (ii)$$

$$\text{and } y + z = 4 \quad \dots (iii)$$

on solving Eqs. (i), (ii) and (iii), we get


$$x = \frac{1}{2}, y = \frac{3}{2}, z = \frac{5}{2}$$

427 (a)

From geometry

$$\vec{AO} = 2\vec{SD}$$

Where D is the mid point of BC



$$\begin{aligned} &\therefore \vec{SA} + \vec{SB} + \vec{SC} \\ &= \vec{SA} + 2\vec{SD} \quad (\because \vec{SB} + \vec{SC} = 2\vec{SD}) \\ &= \vec{SA} + \vec{AO} \\ &= \vec{SO} \end{aligned}$$

428 (c)

We have,

$$\vec{a} \cdot \vec{b} = 0 \text{ and } \vec{a} \times \vec{b} = \vec{0}$$

$$\Rightarrow |\vec{a}||\vec{b}| \cos \theta = 0 \text{ and } |\vec{a}||\vec{b}| \sin \theta = 0$$

$$\Rightarrow (|\vec{a}| = 0 \text{ or } |\vec{b}| = 0 \text{ or } \cos \theta = 0)$$

And,

$$(|\vec{a}| = 0 \text{ or } |\vec{b}| = 0 \text{ or } \sin \theta = 0)$$

$$\Rightarrow |\vec{a}| = 0 \text{ or } |\vec{b}| = 0$$

$$0 \left[ \begin{array}{l} \because \cos \theta \text{ and } \sin \theta \\ \text{are not zero simultaneously} \end{array} \right]$$

430 (c)

$$\text{Given } |\vec{a} + \vec{b}|^2 = |\vec{a} - \vec{b}|^2$$

$$\Rightarrow 4\vec{a} \cdot \vec{b} = 0 \Rightarrow \vec{a} \cdot \vec{b} = 0$$

So, angle between them is  $90^\circ$

431 (c)

We have,

$$\vec{r} \times \vec{a} = \vec{b} \times \vec{a}$$

$$\Rightarrow (\vec{r} - \vec{b}) \times \vec{a} = \vec{0}$$

$$\Rightarrow \vec{r} - \vec{b} \text{ is parallel to } \vec{a}$$

$$\Rightarrow \vec{r} - \vec{b} = \lambda \vec{a} \text{ for some scalar } \lambda$$

$$\Rightarrow \vec{r} - \vec{b} + \lambda \vec{a} \dots (i)$$

Now,

$$\vec{r} \perp \vec{c}$$

$$\Rightarrow \vec{r} \cdot \vec{c} = 0$$

$$\Rightarrow \vec{b} \cdot \vec{c} + \lambda(\vec{a} \cdot \vec{c}) = 0 \Rightarrow \lambda = -\frac{\vec{b} \cdot \vec{c}}{\vec{a} \cdot \vec{c}}$$

Putting the value of  $\lambda$  in (i), we get

$$\vec{r} = \vec{b} - \left( \frac{\vec{b} \cdot \vec{c}}{\vec{a} \cdot \vec{c}} \right) \vec{a}$$

432 (d)

$$\text{We have, } |\vec{\alpha}| = 1 = |\vec{\beta}| \text{ and } \vec{\alpha} \cdot \vec{\beta} = 0$$

Now,

$$\vec{\gamma} = x\vec{\alpha} + y\vec{\beta} + z(\vec{\alpha} \times \vec{\beta})$$

$$\Rightarrow \vec{\alpha} \cdot \vec{\gamma} = x(\vec{\alpha} \cdot \vec{\alpha}) + y(\vec{\alpha} \cdot \vec{\beta}) + z\{\vec{\alpha} \cdot (\vec{\alpha} \times \vec{\beta})\}$$

$$\vec{\beta} \cdot \vec{\gamma} = x(\vec{\beta} \cdot \vec{\alpha}) + y(\vec{\beta} \cdot \vec{\beta}) + z\{\vec{\beta} \cdot (\vec{\alpha} \times \vec{\beta})\}$$

And,

$$\begin{aligned} (\vec{\alpha} \times \vec{\beta}) \cdot \vec{\gamma} &= x\{\vec{\alpha} \cdot (\vec{\alpha} \times \vec{\beta})\} + y\{\vec{\beta} \cdot (\vec{\alpha} \times \vec{\beta})\} \\ &\quad + z\{(\vec{\alpha} \times \vec{\beta}) \cdot (\vec{\alpha} \times \vec{\beta})\} \end{aligned}$$

$$\Rightarrow \cos \theta = x, \cos \theta = y \text{ and } [\vec{\alpha} \vec{\beta} \vec{\gamma}] = z|\vec{\alpha} \times \vec{\beta}|^2$$

$$\Rightarrow x = \cos \theta, y = \cos \theta \text{ and } [\vec{\alpha} \vec{\beta} \vec{\gamma}] = z$$

$$[\because |\vec{\alpha} \times \vec{\beta}| = |\vec{\alpha}| |\vec{\beta}| \sin 90^\circ = 1]$$

Now,

$$[\vec{\alpha} \vec{\beta} \vec{\gamma}]^2 = \begin{vmatrix} \vec{\alpha} \cdot \vec{\alpha} & \vec{\alpha} \cdot \vec{\beta} & \vec{\alpha} \cdot \vec{\gamma} \\ \vec{\beta} \cdot \vec{\alpha} & \vec{\beta} \cdot \vec{\beta} & \vec{\beta} \cdot \vec{\gamma} \\ \vec{\gamma} \cdot \vec{\alpha} & \vec{\gamma} \cdot \vec{\beta} & \vec{\gamma} \cdot \vec{\gamma} \end{vmatrix}$$

$$\Rightarrow [\vec{\alpha} \vec{\beta} \vec{\gamma}]^2 = \begin{vmatrix} 1 & 0 & \cos \theta \\ 0 & 1 & \cos \theta \\ \cos \theta & \cos \theta & 1 \end{vmatrix} = 1 - 2 \cos^2 \theta$$

$$\Rightarrow z^2 = 1 - 2 \cos^2 \theta$$

$$\text{Also, } z^2 = 1 - 2y^2 \text{ and } z^2 = 1 - x^2 - y^2$$

433 (a)

$$(\vec{a} - \vec{d}) \times (\vec{b} - \vec{c})$$

$$= \vec{a} \times \vec{b} - \vec{a} \times \vec{c} - \vec{d} \times \vec{b} + \vec{d} \times \vec{c}$$

$$= \vec{c} \times \vec{d} - \vec{b} \times \vec{d} - \vec{d} \times \vec{b} + \vec{d} \times \vec{c} = \vec{0}$$

$$[\because \vec{a} \times \vec{b} = \vec{c} \times \vec{d}, \vec{a} \times \vec{c} = \vec{b} \times \vec{d}, \text{ given}]$$

$$\Rightarrow (\vec{a} - \vec{d}) \parallel (\vec{b} - \vec{c})$$

$$\Rightarrow \vec{a} - \vec{d} = \lambda(\vec{b} - \vec{c})$$

434 (a)

Since  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar unit vectors

$\therefore [\vec{a} \vec{b} \vec{c}] = \text{Volume of a parallelepiped whose each edge is of one unit length}$

$$\Rightarrow [\vec{a} \vec{b} \vec{c}] = \pm 1$$

436 (d)

Let  $D$  be the mid-point of  $BC$ . Then,

$$\vec{AB} + \vec{AC} = 2\vec{AD}$$

$$\Rightarrow 2\vec{AD} = 8\hat{i} + 2\hat{j} + 8\hat{k}$$

$$\Rightarrow \vec{AD} = 4\hat{i} + \hat{j} + 4\hat{k}$$

$$\Rightarrow |\vec{AD}| = \sqrt{16 + 1 + 16} = \sqrt{33}$$

437 (c)

$$\therefore \text{Median vector through } \vec{A} = \frac{1}{2}(\vec{AB} + \vec{AC})$$

$$= \frac{1}{2}[(3\hat{i} + 5\hat{j} + 4\hat{k}) + (5\hat{i} - 5\hat{j} + 2\hat{k})]$$

$$= 4\hat{i} + 3\hat{k}$$

$$\therefore \text{Length of the median} = \sqrt{4^2 + 3^2} = 5 \text{ units}$$

438 (d)

$$\text{Given, } (\vec{a} - \lambda \vec{b}) \cdot (\vec{b} - 2\vec{c}) \times (\vec{c} + 2\vec{a}) = 0$$

$$\Rightarrow (\vec{a} - \lambda \vec{b}) \cdot \{\vec{b} \times \vec{c} + \vec{b} \times 2\vec{a} - 4(\vec{c} \times \vec{a})\} = 0$$

$$\Rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times 2\vec{a}) - \vec{a} \cdot 4(\vec{c} \times \vec{a})$$

$$- \lambda \vec{b} \cdot (\vec{b} \times \vec{c}) - \lambda \vec{b} \cdot (\vec{b} \times 2\vec{a}) + 4\lambda \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$$

$$\Rightarrow \vec{a}(\vec{b} \times \vec{c}) + 4\lambda \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$$

$$\Rightarrow \{\vec{a} \cdot (\vec{b} \times \vec{c})\}(1 + 4\lambda) = 0$$

$$\Rightarrow \lambda = -\frac{1}{4} [\because \vec{a} \cdot (\vec{b} \times \vec{c}) \neq 0, \text{ given}]$$

440 (d)

$$\therefore \text{Total force } \vec{P} = \vec{P}_1 + \vec{P}_2 + \vec{P}_3$$

$$= \hat{i} - \hat{j} + \hat{k} - \hat{i} + 2\hat{j} - \hat{k} + \hat{j} - \hat{k} = 2\hat{j}$$

$$\begin{aligned} \text{and displacement } \overrightarrow{AB} &= 6\hat{i} + \hat{j} - 3\hat{k} - (4\hat{i} + 3\hat{j} - 2\hat{k}) \\ &= 2\hat{i} + 4\hat{j} - \hat{k} \end{aligned}$$

$$\begin{aligned} \therefore \text{Work done} &= \vec{P} \cdot \overrightarrow{AB} \\ &= 2\hat{j} \cdot (2\hat{i} + 4\hat{j} - \hat{k}) = 8 \end{aligned}$$

441 (a)

$$\begin{aligned} \text{The point of intersection of } \vec{r} \times \vec{a} = \vec{b} \times \vec{a} \text{ and } \vec{r} \times \\ \vec{b} = \vec{a} \times \vec{b} \text{ is } \vec{r} = \vec{a} + \vec{b} \\ \therefore \vec{r} = (\hat{i} + \hat{j}) + (2\hat{i} - \hat{k}) = 3\hat{i} + \hat{j} - \hat{k} \end{aligned}$$

442 (a)

$$\begin{aligned} \text{Since } \vec{a}, \vec{b} \text{ and } \vec{a} \times \vec{b} \text{ are non-coplanar vectors} \\ \therefore \vec{r} = x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b}) \text{ for some scalars } x, y, z \\ \dots(i) \end{aligned}$$

Now,

$$\begin{aligned} \vec{b} &= \vec{r} \times \vec{a} \\ \Rightarrow \vec{b} &= \{x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})\} \times \vec{a} \\ \Rightarrow \vec{b} &= y(\vec{b} \times \vec{a}) + z((\vec{a} \times \vec{b}) \times \vec{a}) \\ \Rightarrow \vec{b} &= y(\vec{b} \times \vec{a}) - z(\vec{a} \times (\vec{a} \times \vec{b})) \\ \Rightarrow \vec{b} &= y(\vec{b} \times \vec{a}) - z\{(\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}\} \\ \Rightarrow \vec{b} &= y(\vec{b} \times \vec{a}) + z(\vec{a} \cdot \vec{a})\vec{b} \quad [\because \vec{a} \cdot \vec{b} = 0] \end{aligned}$$

Comparing the coefficients, we get

$$y = 0, z = \frac{1}{\vec{a} \cdot \vec{a}} = \frac{1}{|\vec{a}|^2}$$

Putting the values of  $y$  and  $z$  in (i), we get

$$\vec{r} = x\vec{a} + \frac{1}{|\vec{a}|^2}(\vec{a} \times \vec{b})$$

444 (b)

$$\begin{aligned} (\vec{u} + \vec{v} - \vec{w}) \cdot [(\vec{u} - \vec{v}) \times (\vec{v} - \vec{w})] \\ = (\vec{u} + \vec{v} - \vec{w}) \cdot [\vec{u} \times \vec{v} - \vec{u} \times \vec{w} + \vec{v} \times \vec{w}] \\ = \vec{u} \cdot \vec{v} \times \vec{w} - \vec{v} \cdot \vec{u} \times \vec{w} - \vec{w} \cdot \vec{u} \times \vec{v} \\ = \vec{u} \cdot \vec{v} \times \vec{w} + \vec{w} \cdot \vec{u} \times \vec{v} - \vec{w} \cdot \vec{u} \times \vec{v} \\ = \vec{u} \cdot \vec{v} \times \vec{w} \end{aligned}$$

445 (d)

$$\begin{aligned} \therefore \vec{p} - 2\vec{q} &= 7\hat{i} - 2\hat{j} + 3\hat{k} - 2(3\hat{i} + \hat{j} + 5\hat{k}) \\ &= \hat{i} - 4\hat{j} - 7\hat{k} \\ \Rightarrow |\vec{p} - 2\vec{q}| &= \sqrt{1^2 + (-4)^2 + (-7)^2} = \sqrt{66} \end{aligned}$$

447 (a)

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix}$$

$$= -\hat{i} + \hat{j}$$

$$\Rightarrow (\vec{a} \times \vec{b}) \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & 0 \\ 1 & 0 & 0 \end{vmatrix} = -\hat{k}$$

$$\begin{aligned} \text{Now, } \lambda \vec{a} + \mu \vec{b} &= \lambda(\hat{i} + \hat{j} + \hat{k}) + \mu(\hat{i} + \hat{j}) \\ &= (\lambda + \mu)\hat{i} + (\lambda + \mu)\hat{j} + \lambda\hat{k} \end{aligned}$$

$$\therefore \lambda \vec{a} + \mu \vec{b} = (\vec{a} \times \vec{b}) \times \vec{c}$$

$$\Rightarrow (\lambda + \mu)\hat{i} + (\lambda + \mu)\hat{j} + \lambda\hat{k} = -\hat{k}$$

On equating the coefficient of  $\hat{i}$  we get  $\lambda + \mu = 0$

453 (d)

We have,

$$\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$$

$$\Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0$$

$$\Rightarrow \vec{a} \perp (\vec{b} - \vec{c}) \text{ or, } \vec{b} - \vec{c} = 0 \Rightarrow \vec{a} \perp (\vec{b} - \vec{c}) \text{ or,}$$

$$\vec{b} = \vec{c}$$

454 (c)

$$\text{Given that, } |\vec{a}| = 2\sqrt{2}, |\vec{b}| = 3$$

The longer vectors is  $5\vec{a} + 2\vec{b} + \vec{a} - 3\vec{b} = 6\vec{a} - \vec{b}$

Length of one diagonal

$$= |6\vec{a} - \vec{b}|$$

$$= \sqrt{36\vec{a}^2 + \vec{b}^2 - 2 \times 6|\vec{a}||\vec{b}| \cos 45^\circ}$$

$$= \sqrt{36 \times 8 + 9 - 12 \times 2\sqrt{2} \times 3 \times \frac{1}{\sqrt{2}}}$$

$$= \sqrt{288 + 9 - 12 \times 6} = \sqrt{225} = 15$$

Other diagonal is  $4\vec{a} + 5\vec{b}$ .

$$\text{Its length} = \sqrt{16 \times 8 + 25 \times 9 + 40 \times 6} = \sqrt{593}$$

455 (a)

Given projection of  $\vec{a}$  on  $\vec{b} = |\vec{a} \times \vec{b}|$

$$\Rightarrow \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = |\vec{a} \times \vec{b}|$$

$$\Rightarrow \frac{|\vec{a}||\vec{b}| \cos \theta}{|\vec{b}|} = |\vec{a}||\vec{b}| \sin \theta$$

$$\Rightarrow \tan \theta = \frac{1}{|\vec{b}|}$$

$$\Rightarrow \tan \theta = \frac{1}{\frac{1}{3}\sqrt{1^2 + 1^2 + 1^2}}$$

$$\Rightarrow \tan \theta = \sqrt{3}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

457 (c)

Since,  $\vec{a} + 2\vec{b} = k\vec{c}$

$$\begin{aligned} \therefore \vec{a} + 2\vec{b} + 6\vec{c} &= k\vec{c} + 6\vec{c} \\ &= (k + 6)\vec{c} = \lambda\vec{c} \quad (\because \lambda \neq 0) \end{aligned}$$

458 (d)

$$\vec{u} \times \vec{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 0 \\ 1 & -1 & 0 \end{vmatrix} = -2\hat{k}$$


$$\therefore |\vec{w} \cdot \hat{n}| = \frac{|\vec{w} \cdot \vec{u} \times \vec{v}|}{|\vec{u} \times \vec{v}|}$$

$$\Rightarrow |\vec{w} \cdot \hat{n}| = \frac{|-6\hat{k}|}{|-2\hat{k}|} = 3$$

459 (c)

Let the position of B is  $\vec{r}$ .

$$\therefore \vec{a} = \frac{2\vec{r} + 3(\vec{a} + 2\vec{b})}{2 + 3}$$



$$\Rightarrow 5\vec{a} = 2\vec{r} + 3\vec{a} + 6\vec{b}$$

$$\Rightarrow 2\vec{r} = 2\vec{a} - 6\vec{b}$$

$$\therefore \vec{r} = \vec{a} - 3\vec{b}$$

460 (a)

Since,  $(\vec{A} + t\vec{B}) \cdot \vec{C} = 0$  [given]

$$\Rightarrow [(1-t)\hat{i} + (2+2t)\hat{j} + (3+t)\hat{k}] \cdot (3\hat{i} + \hat{j}) = 0$$

$$\Rightarrow 3(1-t) + (2+2t) = 0 \Rightarrow t = 5$$

461 (a)

We have,

$$|\vec{a}| = 1, |\vec{b}| = 1 \text{ and } \vec{a} \cdot \vec{b} = \cos \theta$$

$$\text{Now, } |\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow |\vec{a} - \vec{b}|^2 = 1 + 1 - 2|\vec{a}| |\vec{b}| \cos \theta$$

$$\Rightarrow |\vec{a} - \vec{b}|^2 = 4 \sin^2 \frac{\theta}{2}$$

$$\Rightarrow \left| \frac{\vec{a} - \vec{b}}{2} \right|^2 = \sin^2 \frac{\theta}{2} \Rightarrow \left| \frac{\vec{a} - \vec{b}}{2} \right| = \sin \frac{\theta}{2}$$

462 (c)


If  $\vec{a}, \vec{b}$  are two non-zero non-collinear vectors and  $x, y$  are two scalars such that  $x\vec{a} + y\vec{b} = \vec{0}$ , then  $x = 0, y = 0$ .

Because otherwise one will be a scalar multiple of the other and hence collinear, which is a contradiction

463 (b)

$$\vec{AB} = 2\hat{i} + 10\hat{j} + 11\hat{k}$$

$$\vec{AD} = -\hat{i} + 2\hat{j} + 2\hat{k}$$



$$\vec{AB} \cdot \vec{AD} = -2 + 20 + 22 = 40$$

$$|\vec{AB}| = \sqrt{4 + 100 + 120} = \sqrt{225} = 15$$

$$|\vec{AD}| = \sqrt{1 + 4 + 4} = \sqrt{9} = 3$$

$$\therefore \cos \theta = \frac{40}{45} = \frac{8}{9}$$

$$\therefore \theta + \alpha = 90^\circ$$

$$\Rightarrow \alpha = 90^\circ - \theta$$

$$\Rightarrow \cos \alpha = \sin \theta = \sqrt{1 - \frac{64}{81}} = \frac{\sqrt{17}}{9}$$

464 (a)

$$\text{Let } \vec{a} = x\hat{i} + \hat{j} + \hat{k} \text{ and } \vec{b} = 2\hat{i} - \hat{j} + 5\hat{k}$$

$$\text{Since, } \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{1}{\sqrt{30}}$$

$$\Rightarrow \frac{(x\hat{i} + \hat{j} + \hat{k}) \cdot (2\hat{i} - \hat{j} + 5\hat{k})}{|\sqrt{4 + 1 + 25}|} = \frac{1}{\sqrt{30}}$$

$$\Rightarrow 2x - 1 + 5 = 1$$

$$\Rightarrow x = -\frac{3}{2}$$

465 (b)

$$\text{Now, } 2\vec{a} - \vec{c} = 2(-\hat{i} + \hat{j} + 2\hat{k}) - (2\hat{i} + \hat{j} + 3\hat{k})$$

$$= \hat{j} + 3\hat{k}$$

$$\text{and } \vec{a} + \vec{b} = -\hat{i} + \hat{j} + 2\hat{k} + 2\hat{i} - \hat{j} - \hat{k}$$

$$= \hat{i} + \hat{k}$$

let  $\theta$  be the angle between  $2\vec{a} - \vec{c}$  and  $\vec{a} + \vec{b}$ .

$$\therefore \cos \theta = \frac{(\hat{j} + \hat{k}) \cdot (\hat{i} + \hat{j})}{\sqrt{1^2 + 1^2} \sqrt{1^2 + 1^2}}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}\sqrt{2}} = \frac{1}{2}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

466 (d)

Since  $\vec{a} + \vec{b}$  and  $\vec{b} + \vec{c}$  are collinear with  $\vec{c}$  and  $\vec{a}$  respectively. Therefore, there exist scalars  $x, y$

such that  $\vec{a} + \vec{b} = x\vec{c}$  and  $\vec{b} + \vec{c} = y\vec{a}$ . Now,

$$\vec{a} + \vec{b} = x\vec{c} \Rightarrow \vec{a} + \vec{b} + \vec{c} = (x+1)\vec{c} \dots (i)$$

and,

$$\vec{b} + \vec{c} = y\vec{a} \Rightarrow \vec{a} + \vec{b} + \vec{c} = (y+1)\vec{a} \dots (ii)$$

From (i) and (ii), we get

$$(x+1)\vec{c} = (y+1)\vec{a}$$

If  $x \neq -1$ , then

$$(x+1)\vec{c} = (y+1)\vec{a} \Rightarrow \vec{c} = \frac{y+1}{x+1}\vec{a}$$

$\Rightarrow \vec{c}$  and  $\vec{a}$  are collinear

This is a contradiction to the given condition.

Therefore,  $x = -1$

Putting  $x = -1$  in  $\vec{a} + \vec{b} = x\vec{c}$ , we get

$$\vec{a} + \vec{b} + \vec{c} = (-1+1)\vec{c} = \vec{0}$$

467 (b)

$$\text{We have, } [\vec{a} \vec{b} + \vec{c} \vec{a} + \vec{b} + \vec{c}]$$

$$= \vec{a} \cdot [(\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})]$$

$$\begin{aligned}
&= \vec{a} \cdot (\vec{b} \times \vec{a} + \vec{b} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b} + \vec{c} \times \vec{c}) \\
&= \vec{a} \cdot (\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} - \vec{b} \times \vec{c}) \\
&= \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) \\
&= [\vec{a} \vec{b} \vec{c}] + [\vec{a} \vec{c} \vec{a}] = 0
\end{aligned}$$

468 (a)

It is given that points  $P, Q$  and  $R$  with position vectors  $60\hat{i} + 3\hat{j}, 40\hat{i} - 8\hat{j}$  and  $a\hat{i} - 52\hat{j}$  respectively are collinear

$$\begin{aligned}
\therefore \vec{PQ} &= \lambda \vec{QR} \text{ for some scalar } \lambda \\
\Rightarrow -20\hat{i} - 11\hat{j} &= \lambda\{(a - 40)\hat{i} - 44\hat{j}\} \\
\Rightarrow \lambda(a - 40) &= -20, -11 = -44\lambda \\
\Rightarrow \lambda &= \frac{1}{4} \text{ and } a = -40
\end{aligned}$$

469 (a)

Required unit vector

$$\vec{c} = \frac{\vec{a} \times (\vec{a} \times \vec{b})}{|\vec{a} \times (\vec{a} \times \vec{b})|}$$

Now,

$$\begin{aligned}
\vec{a} \times (\vec{a} \times \vec{b}) &= (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} \\
&= 3(2\hat{i} + \hat{j} + \hat{k}) - 6(\hat{i} + 2\hat{j} - \hat{k}) \\
&= -9\hat{j} + 9\hat{k} \\
\therefore \vec{c} &= \frac{-9\hat{j} + 9\hat{k}}{\sqrt{9^2 + 9^2}} = \pm \frac{1}{\sqrt{2}}(-\hat{j} + \hat{k})
\end{aligned}$$

470 (b)

$$\begin{aligned}
\begin{vmatrix} 2 & 1 & 4 \\ 4 & -2 & 3 \\ 2 & -3 & -\lambda \end{vmatrix} &= 0 \\
\Rightarrow 2(2\lambda + 9) - 1(-4\lambda - 6) + 4(-12 + 4) &= 0 \\
\Rightarrow 4\lambda + 18 + 4\lambda + 6 - 48 + 16 &= 0 \\
\Rightarrow 8\lambda &= 8 \\
\Rightarrow \lambda &= 1
\end{aligned}$$

471 (b)

We have,

$$\begin{aligned}
[\vec{u} \vec{v} \vec{w}] &= \begin{vmatrix} al + a_1l_1 & am + a_1m_1 & an + a_1n_1 \\ bl + b_1l_1 & bm + b_1m_1 & bn + b_1n_1 \\ cl + c_1l_1 & cm + c_1m_1 & cn + a_1n_1 \end{vmatrix} \\
\Rightarrow [\vec{u} \vec{v} \vec{w}] &= \begin{vmatrix} a & a_1 & 0 \\ b & b_1 & 0 \\ c & c_1 & 0 \end{vmatrix} \begin{vmatrix} l & l_1 & 0 \\ m & m_1 & 0 \\ n & n_1 & 0 \end{vmatrix} = 0
\end{aligned}$$

Hence, the given vectors are coplanar

473 (a)

Given that  $\vec{a}, \vec{b}, \vec{c}$  are coplanar

$$\therefore \vec{a} \perp \vec{b} \times \vec{c} \Rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) = 0 \Rightarrow [\vec{a} \vec{b} \vec{c}] = 0$$

474 (c)

$$\begin{aligned}
&(\vec{d} + \vec{a}) \cdot [\vec{a} \times \{\vec{b} \times (\vec{c} \times \vec{d})\}] \\
&= (\vec{d} + \vec{a}) \cdot [\vec{a} \times \{\vec{b} \cdot \vec{d}\vec{c} - (\vec{b} \cdot \vec{c})\vec{d}\}] \\
&= (\vec{b} \cdot \vec{d})[\vec{d} \cdot (\vec{a} \times \vec{c})] - (\vec{b} \cdot \vec{c})[\vec{d} \cdot (\vec{a} \times \vec{d})]
\end{aligned}$$

$$\begin{aligned}
&+ (\vec{b} \cdot \vec{d})[\vec{a} \cdot (\vec{a} \times \vec{c})] - (\vec{b} \cdot \vec{c})[\vec{a} \cdot (\vec{a} \times \vec{d})] \\
&= (\vec{b} \cdot \vec{d})[\vec{d} \cdot \vec{a} \vec{c}] = (\vec{b} \cdot \vec{d})[\vec{a} \vec{c} \vec{d}]
\end{aligned}$$

476 (a)

$$\begin{aligned}
\text{Let } \vec{a} &= \hat{i} - 2\hat{j} + 3\hat{k}, \vec{b} = -2\hat{i} + 3\hat{j} - 4\hat{k} \\
\text{and } \vec{c} &= \lambda\hat{i} - \hat{j} + 2\hat{k}
\end{aligned}$$

$$\therefore [\vec{a}, \vec{b}, \vec{c}] = 0$$

$$\Rightarrow \begin{vmatrix} 1 & -2 & 3 \\ -2 & 3 & -4 \\ \lambda & -1 & 2 \end{vmatrix} = 0$$

$$\begin{aligned}
\Rightarrow 1(6 - 4) + 2(-4 + 4\lambda) + 3(2 - 3\lambda) &= 0 \\
\Rightarrow \lambda &= 0
\end{aligned}$$

477 (b)

$$\text{Let } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$|\vec{a}|^2 = a_1^2 + a_2^2 + a_3^2$$

$$\text{and } \vec{a} \times \hat{i} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \times \hat{i}$$

$$= -a_2\hat{k} + a_3\hat{j}$$

$$(\vec{a} \times \hat{i})^2 = a_2^2 + a_3^2$$

$$\text{Similarly, } (\vec{a} \times \hat{j})^2 = a_3^2 + a_1^2$$

$$\text{and } (\vec{a} \times \hat{k})^2 = a_1^2 + a_2^2$$

$$\text{Now, } (\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2$$

$$= a_2^2 + a_3^2 + a_3^2 + a_1^2 + a_1^2 + a_2^2$$

$$= 2(a_1^2 + a_2^2 + a_3^2) = 2(\vec{a})^2$$

478 (d)

Since,  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 2\hat{i} - 4\hat{k}, \vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$  are coplanar.

$$\therefore [\vec{a} \vec{b} \vec{c}] = 0 \Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 2 & 0 & -4 \\ 1 & \lambda & 3 \end{vmatrix} = 0$$

$$\Rightarrow 4\lambda - 1(6 + 4) + 2\lambda = 0$$

$$\Rightarrow 6\lambda = 10 \Rightarrow \lambda = \frac{5}{3}$$

480 (c)

$\vec{A}, \vec{B}$  and  $\vec{C}$  are three vectors, then volume of parallelepiped

$$V = [\vec{A} \vec{B} \vec{C}]$$

$$= \begin{vmatrix} 1 & a & 1 \\ 0 & 1 & a \\ a & 0 & 1 \end{vmatrix} = 1 + a^3 - a$$

$$\Rightarrow V = 1 + a^3 - a$$

On differentiating with respect to  $a$ , we get

$$\frac{dV}{da} = 3a^2 - 1 = 0$$

For maximum or minimum, put  $\frac{dV}{da} = 0$

$$\Rightarrow a = \pm \frac{1}{\sqrt{3}}$$

$$\frac{d^2V}{da^2} = 6a, \text{ positive at } a = \frac{1}{\sqrt{3}}$$

$\therefore V$  is minimum at  $a = \frac{1}{\sqrt{3}}$

481 (c)

By the properties of midpoint theorem,

$$\overrightarrow{PA} + \overrightarrow{PB} = 2\overrightarrow{PC}$$

482 (a)

The vector equation of line passing through points (3, 2, 1) and (-2, 1, 3)

$$\begin{aligned} \vec{r} &= 3\hat{i} + 2\hat{j} + \hat{k} + \lambda[(-2 - 3)\hat{i} + (1 - 2)\hat{j} \\ &\quad + (3 - 1)\hat{k}] \\ &= 3\hat{i} + 2\hat{j} + \hat{k} + \lambda(-5\hat{i} - \hat{j} + 2\hat{k}) \end{aligned}$$

483 (d)

$$\begin{aligned} \because \vec{a} \cdot \vec{b} &= |\vec{a}||\vec{b}| \cos \frac{5\pi}{6} \\ &= -\frac{|\vec{a}||\vec{b}|\sqrt{3}}{2} \end{aligned}$$

Since, the projection of  $\vec{a}$  in the direction of

$$\begin{aligned} \vec{b} &= -\frac{6}{\sqrt{3}} \\ \Rightarrow -\frac{|\vec{a}||\vec{b}|\sqrt{3}}{2|\vec{b}|} &= -\frac{6}{\sqrt{3}} \\ \Rightarrow |\vec{a}| &= \frac{6 \times 2}{3} = 4 \end{aligned}$$

484 (d)

Let  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  in  $OXYZ$  system

Also, let  $\vec{r} = X\hat{i} + Y\hat{j} + Z\hat{k}$  in the new coordinate system

Since the right handed rectangular system  $OXYZ$  is rotated about  $z$ -axis through  $\frac{\pi}{4}$  in anticlockwise direction. Therefore,

$$\begin{aligned} x &= X \cos \theta - Y \sin \theta \text{ and } y = X \sin \theta + Y \cos \theta \\ \Rightarrow x &= X \cos \frac{\pi}{4} - Y \sin \frac{\pi}{4}, y = X \sin \frac{\pi}{4} + Y \cos \frac{\pi}{4} \\ \text{and, } z &= Z \end{aligned}$$

It is given that  $X = 2\sqrt{2}, Y = 3\sqrt{2}$  and  $Z = 4$

$$\therefore x = 2 - 3 = -1, y = 5 \text{ and } z = 4$$

$$\text{Hence, } \vec{r} = -\hat{i} + 5\hat{j} + 4\hat{k}$$

ALITER Let  $l_1, m_1, n_1; l_2, m_2, n_2$  and  $l_3, m_3, n_3$  be the direction cosines of the new axes with respect to the old axes. Then,

$$\begin{aligned} l_1 &= \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}, m_1 = \cos \left(-\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}, n_1 \\ &= \cos \frac{\pi}{2} = 0 \end{aligned}$$

$$\begin{aligned} l_2 &= \cos \frac{3\pi}{4} = -\frac{1}{\sqrt{2}}, m_2 = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}, n_2 \\ &= \cos \frac{\pi}{2} = 0 \end{aligned}$$

$$l_3 = \cos \frac{\pi}{2} = 0, m_3 = \cos \frac{\pi}{2} = 0, n_3 = \cos 0 = 1$$

Let  $x', y', z'$  and  $x, y, z$  be the components of the given vector with respect to new and old axes.

Then,

$$\begin{aligned} \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \\ n_1 & n_2 & n_3 \end{bmatrix} \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} \\ \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2\sqrt{2} \\ 3\sqrt{2} \\ 4 \end{bmatrix} \\ \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} 2 & -3 & +0 \\ 2 & +3 & +0 \\ 0 & 0 & +4 \end{bmatrix} = \begin{bmatrix} -1 \\ 5 \\ 4 \end{bmatrix} \end{aligned}$$

Hence, the components of  $\vec{a}$  in the  $Oxyz$  coordinate system are  $-1, 5, 4$

485 (d)

$$\because \vec{x} \cdot \vec{a} = \vec{x} \cdot \vec{b} = \vec{x} \cdot \vec{c} = 0$$

For non-zero vector  $\vec{x}$

$$[\vec{a} \vec{b} \vec{c}] = 0 \quad (\text{three vectors } \vec{a}, \vec{b}, \vec{c} \text{ are coplanar})$$

$$\text{and } [\vec{a} \times \vec{b} \vec{b} \times \vec{c} \vec{c} \times \vec{a}]$$


$$= [\vec{a} \vec{b} \vec{c}]^2 = 0$$

486 (d)

$ABCDEF$  is a regular hexagon. We know from the hexagon that  $\overrightarrow{AD}$  is parallel to  $\overrightarrow{BC}$ .

$$\Rightarrow \overrightarrow{AD} = 2\overrightarrow{BC}$$

Similarly,  $\overrightarrow{EB}$  is a parallel to  $\overrightarrow{FA}$



$$\Rightarrow \overrightarrow{EB} = 2\overrightarrow{FA}$$

and  $\overrightarrow{FC}$  is parallel to  $\overrightarrow{AB}$ .

$$\Rightarrow \overrightarrow{FC} = 2\overrightarrow{AB}$$

$$\text{Thus, } \overrightarrow{AD} + \overrightarrow{EB} + \overrightarrow{FC} = 2\overrightarrow{BC} + 2\overrightarrow{FA} + 2\overrightarrow{AB}$$

$$= 2(\overrightarrow{FA} + \overrightarrow{AB} + \overrightarrow{BC})$$

$$= 2(\overrightarrow{FC}) = 2(2\overrightarrow{AB}) = 4\overrightarrow{AB}$$

487 (d)

$$\text{Here, } \vec{a}_1 = 6\hat{i} + 2\hat{j} + 2\hat{k}, \vec{a}_2 = -4\hat{i} + 0\hat{j} - \hat{k},$$

$$\vec{b}_1 = \hat{i} - 2\hat{j} + 2\hat{k} \text{ and } \vec{b}_2 = 3\hat{i} - 2\hat{j} - 2\hat{k}$$

∴ Shortest distance

$$= \frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

$$= \frac{|(-10\hat{i} - 2\hat{j} - 3\hat{k}) \cdot (8\hat{i} + 8\hat{j} + 4\hat{k})|}{\sqrt{64 + 64 + 16}}$$

$$= \left| -\frac{108}{12} \right| = 9$$

488 (c)


$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -6 & -3 \\ 4 & 3 & -1 \end{vmatrix} = 15\hat{i} - 10\hat{j} + 30\hat{k}$$

$$\text{and } |\vec{a} \times \vec{b}| = \sqrt{15^2 + (-10)^2 + (30)^2} = 35$$

$$\therefore \text{Required vector} = \frac{3\hat{i} - 2\hat{j} + 6\hat{k}}{7}$$

490 (a)

Let  $O$  be the origin



$$\therefore \vec{BE} + \vec{AF} = \vec{OE} - \vec{OB} + \vec{OF} - \vec{OA}$$

$$= \frac{\vec{OA} + \vec{OC}}{2} - \vec{OB} + \frac{\vec{OB} + \vec{OC}}{2} - \vec{OA}$$

$$= \frac{\vec{OC}}{2} + \frac{\vec{OC}}{2} + \frac{\vec{OA}}{2} - \vec{OA} + \frac{\vec{OB}}{2} - \vec{OB}$$

$$= \vec{OC} - \frac{\vec{OA} + \vec{OB}}{2} = \vec{OC} - \vec{OD} = \vec{DC}$$

491 (d)

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|\cos\theta$$

$$\Rightarrow |\vec{a} - \vec{b}|^2 = 1 + 1 - 2\cos 60^\circ = 2 - 1$$

$$\Rightarrow |\vec{a} - \vec{b}| = 1$$

492 (b)

$$\text{Given, } 2\vec{a} + 3\vec{b} + \vec{c} = \vec{0}$$

$$\Rightarrow 2\vec{a} + 3\vec{b} = -\vec{c}$$

Taking cross product with  $\vec{a}$  and  $\vec{b}$  respectively, we get

$$2(\vec{a} \times \vec{a}) + 3(\vec{a} \times \vec{b}) = -\vec{a} \times \vec{c}$$

$$\Rightarrow 3(\vec{a} \times \vec{b}) = -\vec{c} \times \vec{a} \dots (i)$$

$$\text{and } 2(\vec{b} \times \vec{a}) + 3(\vec{b} \times \vec{b}) = -\vec{b} \times \vec{c}$$

$$\Rightarrow 2(\vec{a} \times \vec{b}) = \vec{b} \times \vec{c} \dots (ii)$$

$$\text{Now, } \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$$

$$= \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + 3(\vec{a} \times \vec{b}) \quad [\text{using Eq. (i)}]$$

$$= 4(\vec{a} \times \vec{b}) + \vec{b} \times \vec{c}$$

$$= 2(\vec{b} \times \vec{c}) + \vec{b} \times \vec{c} \quad [\text{using Eq. (ii)}]$$

$$= 3(\vec{b} \times \vec{c})$$

493 (d)

$$[\vec{a} - 2\vec{b}, \vec{b} - 3\vec{c}, \vec{c} - 4\vec{a}]$$

$$= (\vec{a} - 2\vec{b}) \cdot \{(\vec{b} - 3\vec{c}) \times (\vec{c} - 4\vec{a})\}$$

$$= (\vec{a} - 2\vec{b}) \cdot \{\vec{b} \times \vec{c} - 4\vec{b} \times \vec{a} + 12\vec{c} \times \vec{a}\}$$

$$= (\vec{a} - 2\vec{b}) \cdot (\vec{a} + 4\vec{c} + 12\vec{b})$$

$$= \vec{a} \cdot \vec{a} - 24\vec{b} \cdot \vec{b}$$

$$= 1 - 24 \times 9 = 1 - 216 = -215$$

494 (b)

$$\text{Given, area} = |\vec{a} \times \vec{b}| = 15$$

If the sides are  $(3\vec{a} + 2\vec{b})$  and  $(\vec{a} + 3\vec{b})$ , then

Area of parallelogram

$$= |(3\vec{a} + 2\vec{b}) \times (\vec{a} + 3\vec{b})| = 7|\vec{a} \times \vec{b}|$$

$$= 7 \times 15 = 105 \text{ sq units}$$

498 (a)

$$\text{Given, } \vec{a} \cdot (\vec{b} + \vec{c}) = 0 \Rightarrow \vec{a} \cdot \vec{b} + \vec{c} \cdot \vec{a} = 0$$

$$\vec{b} \cdot (\vec{c} + \vec{a}) = 0$$

$$\Rightarrow \vec{b} \cdot \vec{c} + \vec{a} \cdot \vec{b} = 0$$

$$\text{and } \vec{c} \cdot (\vec{a} + \vec{b}) = 0$$

$$\Rightarrow \vec{c} \cdot \vec{a} + \vec{b} \cdot \vec{c} = 0$$

$$\therefore \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = 0$$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 9 + 16 + 25 + 0 = 50$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = 5\sqrt{2}$$

499 (b)

We have,

$$(\vec{b} \times \vec{c}) \times \vec{a} = -\{\vec{a} \times (\vec{b} \times \vec{c})\}$$

$$\Rightarrow (\vec{b} \times \vec{c}) \times \vec{a} = -\{(\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}\}$$

$$= (\vec{a} \cdot \vec{b})\vec{c} - (\vec{a} \cdot \vec{c})\vec{b}$$

501 (c)

$$\text{Since, } |\vec{u}| = 1, |\vec{v}| = 2, |\vec{w}| = 3$$

$$\text{The projection of } \vec{v} \text{ along } \vec{u} = \frac{\vec{v} \cdot \vec{u}}{|\vec{u}|}$$

$$\text{and the projection of } \vec{w} \text{ along } \vec{u} = \frac{\vec{w} \cdot \vec{u}}{|\vec{u}|}$$

according to given condition,

$$\frac{\vec{v} \cdot \vec{u}}{|\vec{u}|} = \frac{\vec{w} \cdot \vec{u}}{|\vec{u}|} \Rightarrow \vec{v} \cdot \vec{u} = \vec{w} \cdot \vec{u} \dots (i)$$

$$\text{Also, } \vec{v} \cdot \vec{w} = 0$$

$$\text{Now, } |\vec{u} - \vec{v} + \vec{w}|^2 = |\vec{u}|^2 + |\vec{v}|^2 + |\vec{w}|^2$$

$$-2\vec{u} \cdot \vec{v} - 2\vec{v} \cdot \vec{w} + 2\vec{u} \cdot \vec{w}$$

$$= 1 + 4 + 9 - 2\vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{u} \quad [\text{from Eq. (i)}]$$

$$\Rightarrow |\vec{u} - \vec{v} + \vec{w}|^2 = 14 + 0$$

$$\Rightarrow |\vec{u} - \vec{v} + \vec{w}| = \sqrt{14}$$

502 (b)

$$\text{Area of triangle} = \frac{1}{2} \{ \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} \}$$

503 (c)

$$\because (\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$$

$$\Rightarrow (\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a} = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$$

$$\Rightarrow (\vec{b} \cdot \vec{c})\vec{a} = (\vec{a} \cdot \vec{b})\vec{c}$$

$$\Rightarrow \vec{a} \text{ is parallel to } \vec{c}$$

504 (d)

Let  $\vec{r}$  be a unit vector such that

$$\vec{r} = x(\hat{i} + 2\hat{j} + \hat{k}) + y(\hat{i} + \hat{j} + 2\hat{k})$$

$$= (x + y)\hat{i} + (2x + y)\hat{j} + (x + 2y)\hat{k}$$

Since,  $\vec{r} \cdot (2\hat{i} + \hat{j} + \hat{k}) = 0$

$$\Rightarrow 2x + 2y + 2x + y + x + 2y = 0$$

$$\Rightarrow y = -x$$

$$\therefore \vec{r} = x\hat{i} - x\hat{k} \Rightarrow \vec{r} = \frac{\hat{i} - \hat{k}}{\sqrt{2}}$$

505 (a)

Since  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are unit vectors inclined at an angle  $\theta$ . Therefore,

$$|\vec{a}| = |\vec{b}| = |\vec{c}| = 1 \text{ and } \cos \theta = \vec{a} \cdot \vec{c} = \vec{b} \cdot \vec{c}$$

Now,

$$\vec{c} = \alpha\vec{a} + \beta\vec{b} + \gamma(\vec{a} \times \vec{b}) \quad \dots(i)$$

$$\Rightarrow \vec{a} \cdot \vec{c} = \alpha(\vec{a} \cdot \vec{a}) + \beta(\vec{a} \cdot \vec{b}) + \gamma\{\vec{a} \cdot (\vec{a} \times \vec{b})\}$$

$$\Rightarrow \cos \theta = \alpha|\vec{a}|^2 \quad [\because \vec{a} \cdot \vec{b} = 0, \vec{a} \cdot (\vec{a} \times \vec{b}) = 0]$$

$$\Rightarrow \cos \theta = \alpha$$

Similarly, by taking dot product on both sides of

$$(i) \text{ by } \vec{b}, \text{ we get, } \beta = \cos \theta$$

$$\therefore \alpha = \beta$$

Thus, option (a) is incorrect

Again,

$$\vec{c} = \alpha\vec{a} + \beta\vec{b} + \gamma(\vec{a} \times \vec{b})$$

$$\Rightarrow |\vec{c}|^2 = |\alpha\vec{a} + \beta\vec{b} + \gamma(\vec{a} \times \vec{b})|^2$$

$$\Rightarrow |\vec{c}|^2 = \alpha^2|\vec{a}|^2 + \beta^2|\vec{b}|^2 + \gamma^2|\vec{a} \times \vec{b}|^2$$

$$+ 2\alpha\beta(\vec{a} \cdot \vec{b}) + 2\alpha\gamma\{\vec{a} \cdot (\vec{a} \times \vec{b})\}$$

$$+ 2\beta\gamma\{\vec{b} \cdot (\vec{a} \times \vec{b})\}$$

$$\Rightarrow 1 = \alpha^2 + \beta^2 + \gamma^2|\vec{a} \times \vec{b}|^2$$

$$\Rightarrow 1 = 2\alpha^2 + \gamma^2\left\{|\vec{a}|^2|\vec{b}|^2 \sin^2 \frac{\pi}{2}\right\}$$

$$\Rightarrow 1 = 2\alpha^2 + \gamma^2$$

$$\Rightarrow \alpha^2 = \frac{1 - \gamma^2}{2}$$

$$\text{But, } \alpha = \beta = \cos \theta$$

$$\therefore 1 = 2\alpha^2 + \gamma^2 \Rightarrow \gamma^2 = 1 - 2\cos^2 \theta = -\cos 2\theta$$

$$\therefore \alpha^2 = \beta^2 = \frac{1 - \gamma^2}{2} = \frac{1 + \cos 2\theta}{2}$$

Thus, option (b), (c) and (d) are correct

506 (d)

Let  $\vec{a} = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $\vec{b} = -2\hat{i} - \hat{j} + 2\hat{k}$  be the position vectors of points  $A$  and  $B$  respectively.

Then the bisector of  $\angle AOB$  divides  $AB$  in the ratio  $OA : OB$  i.e.  $9 : 3$  or  $3 : 1$ . Therefore, the vector lying along the bisector is

$$\frac{3(-2\hat{i} - \hat{j} + 2\hat{k}) + (7\hat{i} - 4\hat{j} - 4\hat{k})}{3 + 1}$$

$$= \frac{1}{4}(i - 7j + 2k)$$

$$\therefore \text{Required vector} = \pm 5\sqrt{6} \left( \frac{i - 7j + 2k}{\sqrt{54}} \right) = \pm \frac{5}{3}(i - 7j + 2k)$$

507 (b)

Since,  $\vec{a}$  and  $\vec{b}$  are collinear.

$$\therefore \vec{b} = m\vec{a}$$

$$\Rightarrow |\vec{b}| = m|\vec{a}|$$

$$\Rightarrow |\vec{b}| = m\sqrt{4 + 9 + 36} = \pm 7m$$

$$\Rightarrow 21 = \pm 7m \Rightarrow m = \pm 3$$

$$\therefore \vec{b} = \pm 3\vec{a} = \pm(2\hat{i} + 3\hat{j} + 6\hat{k})$$

510 (a)

Position vectors of vertices  $A, B$  and  $C$  of the triangle  $ABC$  are  $\vec{a}, \vec{b}$  and  $\vec{c}$

$\therefore$  Centroid of triangle

$$(G) = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$$

$$\text{Now, } \vec{GA} + \vec{GB} + \vec{GC}$$

$$= \left( \vec{a} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right) + \left( \vec{b} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right)$$

$$+ \left( \vec{c} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} \right)$$

$$= \vec{0}$$

511 (d)

Since  $X$  and  $Y$  divide  $\vec{AB}$  internally and externally in the ratio  $2 : 1$ . Therefore, the position vectors of

$X$  and  $Y$  are given by  $\frac{2\vec{b} + \vec{a}}{3}$  and  $2\vec{b} - \vec{a}$  respectively

$$\text{Hence, } \vec{XY} = (2\vec{b} - \vec{a}) - \frac{1}{3}(2\vec{b} + \vec{a}) = \frac{4}{3}(\vec{b} - \vec{a})$$

512 (a)

Let  $\vec{a} = (2, 1, -1)$ ,  $\vec{b} = (1, -1, 0)$  and  $\vec{c} = (5, -1, 1)$

$$\therefore \vec{a} + \vec{b} - \vec{c} = (2 + 1 - 5)\hat{i} + (1 - 1 + 1)\hat{j} + (-1 + 0 - 1)\hat{k}$$

$$= -(2\hat{i} - \hat{j} + 2\hat{k})$$

∴ Unit vector of

$$(\vec{a} + \vec{b} - \vec{c}) = -\frac{(2\hat{i} - \hat{j} + 2\hat{k})}{3}$$

∴ Required unit vector of

$$(\vec{a} + \vec{b} - \vec{c}) = \frac{(2\hat{i} - \hat{j} + 2\hat{k})}{3}$$

513 (b)

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = \hat{i} - \hat{j} + \hat{k}$$

∴ Unit vector

$$= \pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \pm \frac{\hat{i} - \hat{j} + \hat{k}}{1^2 + 1^2 + 1^2}$$

$$= \pm \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$$

So, there are two perpendicular vectors of unit length.

514 (b)

$$\text{Let } \vec{r} = (3\hat{i} + 4\hat{j} + 5\hat{k}) + b(6\hat{i} - 7\hat{j} - 3\hat{k})$$

$$= (3 + 6b)\hat{i} + (4 - 7b)\hat{j} + (5 - 3b)\hat{k}$$

$$\text{Since, } \vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 0$$

$$\Rightarrow (3 + 6b)1 + (4 - 7b)1 - (5 - 3b)1 = 0$$

$$\Rightarrow b = -1$$

$$\therefore \vec{r} = -3\hat{i} + 11\hat{j} + 8\hat{k}$$

515 (d)

$$\text{Given } |\vec{x}| = |\vec{y}| = 1 \text{ and } \vec{x} \cdot \vec{y} = 0$$

$$|\vec{x} + \vec{y}|^2 = |\vec{x}|^2 + |\vec{y}|^2 + 2(\vec{x} \cdot \vec{y})$$

$$\Rightarrow |\vec{x} + \vec{y}|^2 = 1 + 1 + 0$$

$$\Rightarrow |\vec{x} + \vec{y}| = \sqrt{2}$$

516 (c)

$$\text{Let } \vec{A} = \vec{a} \times \vec{b}, \vec{B} = \vec{b} \times \vec{c}, \vec{C} = \vec{c} \times \vec{a}$$

$$\text{Given, } [\vec{A} \vec{B} \vec{C}] = 9 \text{ cu units}$$

$$\text{Using the relation } [\vec{A} \times \vec{B} \vec{B} \times \vec{C} \vec{C} \times \vec{A}] =$$

$$[\vec{A} \vec{B} \vec{C}]^2 = (9)^2 = 81 \text{ cu units}$$

517 (a)

$$\text{Since, } \vec{a} = 8\vec{b} \text{ and } \vec{c} = -7\vec{b}$$

$$\therefore \vec{a} \text{ is parallel to } \vec{b} \text{ and } \vec{c} \text{ is anti-parallel to } \vec{b}$$

$$\Rightarrow \vec{a} \text{ and } \vec{c} \text{ are anti-parallel}$$

$$\Rightarrow \text{Angle between } \vec{a} \text{ and } \vec{c} \text{ is } \pi$$

519 (a)

$$\vec{a} \cdot \vec{c} = (\hat{i} + \hat{j} + \hat{k}) \cdot \hat{i} = 1$$

$$\text{and } \vec{b} \cdot \vec{c} = (\hat{i} + \hat{j}) \cdot \hat{i} = 1$$

$$\text{Now, } (\vec{a} \times \vec{b})\vec{c} = (\vec{c} \cdot \vec{a})\vec{b} - (\vec{c} \cdot \vec{b})\vec{a} = \mu\vec{b} + \lambda\vec{a}$$

$$\Rightarrow \mu = \vec{c} \cdot \vec{a} \text{ and } \lambda = -\vec{c} \cdot \vec{b}$$

$$\Rightarrow \mu = 1 \text{ and } \lambda = -1$$

$$\therefore \mu + \lambda = 1 - 1 = 0$$

520 (b)

Let angle between  $\vec{b}$  and  $\vec{c}$  is  $\alpha$ .

$$\text{Given, } |\vec{b} \times \vec{c}| = \sqrt{15}$$

$$\Rightarrow |\vec{b}||\vec{c}| \sin \alpha = \sqrt{15}$$

$$\Rightarrow \sin \alpha = \frac{\sqrt{15}}{4}$$

$$\therefore \cos \alpha = \sqrt{1 - \sin^2 \alpha} = \sqrt{1 - \frac{15}{16}}$$

$$= \frac{1}{4}$$

$$\therefore \vec{b} - 2\vec{c} = \lambda \vec{a} \quad [\text{given}]$$

$$\Rightarrow (\vec{b} - 2\vec{c})^2 = \lambda^2 (\vec{a})^2$$

$$\Rightarrow \vec{b}^2 + 4\vec{c}^2 - 4\vec{b} \cdot \vec{c} = \lambda^2 \vec{a}^2$$

$$\Rightarrow 16 + 4 \times 1 - 4(|\vec{b}||\vec{c}| \cos \alpha) = \lambda^2 \cdot 1^2$$

$$\Rightarrow 20 - 4 = \lambda^2$$

$$\Rightarrow \lambda = \pm 4$$

521 (a)

The given condition mean that  $\vec{r}$  is perpendicular to all three vectors  $\vec{a} \cdot \vec{b}$  and  $\vec{c}$ . This is possible only if they are coplanar.

$$\therefore [\vec{a} \vec{b} \vec{c}] = 0$$

523 (d)

$$\text{Let } \vec{a} = \hat{i} + \hat{j} \text{ and } \vec{b} = \hat{j} + \hat{k}$$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix}$$

$$= \hat{i} - \hat{j} + \hat{k}$$

$$\text{and } |\vec{a} \times \vec{b}| = \sqrt{1^2 + (-1)^2 + 1^2} = \sqrt{3}$$

∴ Required unit vector

$$= \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$$

Alternate Let  $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$

$$\text{Since, } \vec{a} \cdot (\hat{i} + \hat{j}) = 0 \text{ and } \vec{a} \cdot (\hat{j} + \hat{k}) = 0$$

$$\Rightarrow x + y = 0 \text{ and } y + z = 0$$

$$\text{Also } x^2 + y^2 + z^2 = 1$$

$$\Rightarrow x = 1, y = -1 \text{ and } z = 1$$

$$\therefore \vec{a} = \frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$$

524 (a)

$$\text{Let } \vec{r} = \vec{a} + t\vec{b}$$

$$\Rightarrow \vec{r} = \hat{i}(1 + t) + \hat{j}(2 - t) + \hat{k}(1 + t)$$

Since, The projection of  $\vec{r}$  on  $\vec{c}$ ,

$$\frac{\vec{r} \cdot \vec{c}}{|\vec{c}|} = \frac{|1|}{|\sqrt{3}|} \quad [\text{given}]$$

$$\Rightarrow \frac{1 \cdot (1 + t) + 1 \cdot (2 - t) - 1 \cdot (1 + t)}{\sqrt{3}} = \pm \frac{1}{\sqrt{3}}$$

$$\Rightarrow 2 - t = \pm 1$$

$$\Rightarrow t = 1 \text{ or } 3$$

$$\text{When, } t = 1, \vec{r} = 2\hat{i} + \hat{j} + 2\hat{k}$$

$$\text{When, } t = 3, \vec{r} = 4\hat{i} - \hat{j} + 4\hat{k}$$

525 (a)

$$\text{Given, } \vec{u} \times \vec{v} + \vec{u} = \vec{w} \text{ and } \vec{w} \times \vec{u} = \vec{v}$$

$$\Rightarrow (\vec{u} \times \vec{v} + \vec{u}) \times \vec{u} = \vec{v}$$

$$\Rightarrow (\vec{u} \times \vec{v}) \times \vec{u} = \vec{v}$$

$$\Rightarrow \vec{v} - (\vec{u} \cdot \vec{v})\vec{u} = \vec{v}$$

$$\Rightarrow (\vec{u} \cdot \vec{v})\vec{u} = 0$$

$$\Rightarrow (\vec{u} \cdot \vec{v}) = 0$$

$$\text{Now, } [\vec{u} \vec{v} \vec{w}] = \vec{u} \cdot (\vec{v} \times \vec{w})$$

$$= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v} + \vec{u}))$$

$$= \vec{u} \cdot (\vec{v}(\vec{u} \times \vec{v}) + \vec{v} + \vec{u})$$

$$= \vec{u} \cdot (\vec{v}^2 \times \vec{u} - (\vec{u} \cdot \vec{v}) \cdot \vec{v} + \vec{v} + \vec{u})$$

$$= \vec{v}^2 \vec{u}^2 = 1$$

527 (b)

$$\text{Given, } \frac{(\vec{b} \cdot \vec{a}) \cdot \vec{a}}{|\vec{a}|^2} = \frac{4}{3}(\hat{i} - \hat{j} - \hat{k})$$

$$\Rightarrow \frac{\{(\lambda\hat{i} - 3\hat{j} + \hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k})\}(\hat{i} - \hat{j} - \hat{k})}{(1 + 1 + 1)}$$

$$= \frac{4}{3}(\hat{i} - \hat{j} - \hat{k})$$

$$\Rightarrow (\lambda + 3 - 1)(\hat{i} - \hat{j} - \hat{k}) = 4(\hat{i} - \hat{j} - \hat{k})$$

$$\Rightarrow (\lambda + 2)(\hat{i} - \hat{j} - \hat{k}) = 4(\hat{i} - \hat{j} - \hat{k})$$

On equating the coefficient of  $\hat{i}$ , we get

$$\lambda + 2 = 4 \Rightarrow \lambda = 2$$

528 (a)

$$\text{Given that, } \vec{OA} = \hat{i} + x\hat{j} + 3\hat{k}$$

$$\vec{OB} = 3\hat{i} + 4\hat{j} + 7\hat{k}$$

$$\text{and } \vec{OC} = y\hat{i} - 2\hat{j} - 5\hat{k}$$

Since  $A, B, C$  are collinear. Then  $\vec{A} = \lambda \vec{BC}$

$$\Rightarrow 2\hat{i} + (4 - x)\hat{j} + 4\hat{k} = \lambda[(y - 3)\hat{i} - 6\hat{j} - 12\hat{k}]$$

On comparing the coefficient of  $\hat{i}, \hat{j}$  and  $\hat{k}$ , we get

$$2 = (y - 3)\lambda \quad \dots(i)$$

$$4 - x = -6\lambda \quad \dots(ii)$$

$$\text{and } 4 = -12\lambda \Rightarrow \lambda = -\frac{1}{3} \quad \dots(iii)$$

On putting the value of  $\lambda$  is Eqs. (i) and (ii), we get

$$y = -3 \text{ and } x = 2$$

529 (b)

Given have magnitude of  $\vec{OA}$  and  $\vec{OB}$  are 5 and 6 respectively

$$\text{and } \angle BOA = 60^\circ$$

$$\therefore \vec{OA} \cdot \vec{OB} = |\vec{OA}| |\vec{OB}| \cdot \cos 60^\circ$$

$$\Rightarrow \vec{OA} \cdot \vec{OB} = 5 \cdot 6 \cos 60^\circ$$

$$\Rightarrow \vec{OA} \cdot \vec{OB} = 5 \times 6 \times \frac{1}{2} = 15$$

530 (d)

$$\text{It is given that } |\vec{a}| = |\vec{b}| = |\vec{a} + \vec{b}| = 1$$

We have,

$$|\vec{a} + \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 1 = 1 + 1 + 2|\vec{a}||\vec{b}| \cos \theta$$

$$\Rightarrow \cos \theta = -\frac{1}{2} \Rightarrow \theta = \frac{2\pi}{3}$$

531 (a)

$$\text{Area of } \Delta ABC = \frac{1}{2} |\vec{AB} \times \vec{AC}|$$

$$= \frac{1}{2} \sqrt{4 + 16 + 16} = 3 \text{ sq units}$$

532 (a)

Since,  $\vec{a}, \vec{b}, \vec{c}$  from a right handed system

$$\therefore \vec{c} = \vec{b} \times \vec{a}$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & 0 \\ x & y & z \end{vmatrix} = z\hat{i} - x\hat{k}$$

533 (b)

$$\text{Given that, } |\vec{a}| = |\vec{c}| = 1, |\vec{b}| = 4$$

Let angle between  $\vec{b}$  and  $\vec{c}$  is  $\alpha$ , then

$$|\vec{b} \times \vec{c}| = \sqrt{15} \quad (\text{given})$$

$$\Rightarrow |\vec{b}||\vec{c}| \sin \alpha = \sqrt{15}$$

$$\Rightarrow \sin \alpha = \frac{\sqrt{15}}{4 \times 1} = \frac{\sqrt{15}}{4}$$

$$\therefore \cos \alpha = \sqrt{1 - \sin^2 \alpha} = \frac{1}{4}$$

$$\text{We have, } \vec{b} = 2\vec{c} = \lambda \vec{a}$$

On squaring both sides, we get

$$(\vec{b} - 2\vec{c})^2 = \lambda^2 (\vec{a})^2$$

$$\Rightarrow \vec{b}^2 + 4\vec{c}^2 - 4\vec{b} \cdot \vec{c} = \lambda^2 \vec{a}^2$$

$$\Rightarrow 16 + 4 - 4|\vec{b}||\vec{c}| \cos \alpha = \lambda^2$$

$$\Rightarrow 16 + 4 - 4 \times 4 \times 1 \times \frac{1}{4} = \lambda^2$$

$$\Rightarrow \lambda^2 = 16 + 4 - 4 = 16$$

$$\Rightarrow \lambda = \pm 4$$

534 (a)

We have,

$$\overbrace{P(\vec{a}) \quad Q(\vec{b})} \quad R$$

$$PR = 5 PQ \Rightarrow PQ + QR = 5 PQ \Rightarrow 4 PQ = QR$$

$$\therefore PR : QR = 5 : 4$$

$$\Rightarrow R \text{ divides } PQ \text{ externally in the ratio } 5 : 4$$

$$\Rightarrow \text{Position vector of } R \text{ is } 5\vec{b} - 4\vec{a}$$

536 (a)

We have,

$$\vec{BA} + \vec{BC} + \vec{CD} + \vec{DA}$$

$$= \vec{BA} + (\vec{BC} + \vec{CD}) + \vec{DA} = \vec{BA} + (\vec{BD} + \vec{DA})$$

$$= \vec{BA} + \vec{BA} = 2\vec{BA}$$

537 (a)

Given centre of sphere = (1, 0, 1) and radius = 4

∴ Vector equation of sphere is

$|\vec{r} - \vec{a}| = R$  Where  $\vec{a}$  centre of sphere and  $R$  radius of sphere.

Hence, the vector equation of sphere is


$$|\vec{r} - (\hat{i} + \hat{k})| = 4$$

538 (b)

We have,  $||[\vec{a} \vec{b} \vec{c}]|| = V$

Volume  $V_1$  of the parallelepiped having diagonals of the given parallelepiped as three concurrent edges is given by

$$V_1 = ||[\vec{a} + \vec{b} \vec{b} + \vec{c} \vec{c} + \vec{a}]|| = |2[\vec{a} \vec{b} \vec{c}]| = 2V$$



540 (d)

The given equation is

$$\vec{r}^2 - 2\vec{r} \cdot \vec{c} + h = 0, |\vec{c}| > \sqrt{h}$$

This is the equation of sphere in diameter form.

$$\text{i.e., } (\vec{r} - \vec{a}) \cdot (\vec{r} - \vec{b}) = 0$$

541 (c)

Let the given points be  $A, B, C$  respectively.

If  $A, B, C$  are collinear, then

$$A\vec{B} = \lambda B\vec{C} \text{ for some scalar } \lambda$$

$$\Rightarrow 2\hat{i} - 8\hat{j} = \lambda \{(a-12)\hat{i} + 16\hat{j}\}$$

$$\Rightarrow \lambda(a-12) = 2 \text{ and } 16\lambda = -8$$

$$\Rightarrow a-12 = -4 \Rightarrow a = 8$$

542 (a)

We have,

$$\vec{a} \times (\vec{a} \times \vec{b}) = \vec{b} \times (\vec{b} \times \vec{c})$$

$$\Rightarrow (\vec{a} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b} = (\vec{b} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{b})\vec{c}$$

Taking dot product on both sides by  $\vec{b} \times \vec{c}$ , we get

$$\Rightarrow (\vec{a} \cdot \vec{b})\{\vec{a} \cdot (\vec{b} \times \vec{c})\} - (\vec{a} \cdot \vec{a})\{\vec{b} \cdot (\vec{b} \times \vec{c})\}$$

$$= (\vec{b} \cdot \vec{c})\{\vec{b} \cdot (\vec{b} \times \vec{c})\}$$

$$- (\vec{b} \cdot \vec{b})\{\vec{c} \cdot (\vec{b} \times \vec{c})\}$$

$$\Rightarrow (\vec{a} \cdot \vec{b})[\vec{a} \vec{b} \vec{c}] = 0$$

$$\Rightarrow [\vec{a} \vec{b} \vec{c}] = 0 \quad [\because \vec{a} \cdot \vec{b} \neq 0]$$

543 (a)

We have,

$$[\vec{a} \vec{b} \vec{a} \times \vec{b}] + (\vec{a} \cdot \vec{b})^2$$

$$= (\vec{a} \times \vec{b}) \cdot (\vec{a} \times \vec{b}) + (\vec{a} \cdot \vec{b})^2$$

$$\Rightarrow [\vec{a} \vec{b} \vec{a} \times \vec{b}] + (\vec{a} \cdot \vec{b})^2 = |\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2$$

$$= |\vec{a}|^2 |\vec{b}|^2$$

544 (d)

Since,

$$[3\vec{v} p \vec{v} p \vec{w}] - [p \vec{v} \vec{w} q \vec{u}] - [2 \vec{w} q \vec{v} q \vec{u}] = 0$$

$$\therefore 3p^2[\vec{u} \cdot (\vec{v} \times \vec{w})] - pq[\vec{v} \cdot (\vec{w} \times \vec{u})]$$

$$- 2q^2[\vec{w} \cdot (\vec{v} \times \vec{u})] = 0$$

$$\Rightarrow (3p^2 - pq + 2q^2)[\vec{u} \cdot (\vec{v} \times \vec{w})] = 0$$

But  $[\vec{u} \vec{v} \vec{w}] \neq 0$

$$\Rightarrow 3p^2 - pq + 2q^2 = 0$$

$$\Rightarrow p = q = 0$$

545 (a)

$$\vec{a} \cdot [(\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})]$$

$$= \vec{a} \cdot [\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b}]$$

$$= \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{c} \times \vec{b})$$

$$= [\vec{a} \vec{b} \vec{c}] + [\vec{a} \vec{c} \vec{b}] = 0 \quad [\because [\vec{a} \vec{c} \vec{b}] = -[\vec{a} \vec{b} \vec{c}]]$$

546 (b)

$$\text{Let, } \vec{a} = \hat{i} - 3\hat{j} + 2\hat{k}, \vec{b} = -\hat{i} + 2\hat{j}$$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ -1 & 2 & 0 \end{vmatrix} = -4\hat{i} - 2\hat{j} - \hat{k}$$

$$\therefore \text{Area of parallelogram} = \frac{1}{2} |\vec{a} \times \vec{b}|$$

$$= \frac{1}{2} \sqrt{16 + 4 + 1} = \frac{\sqrt{21}}{2}$$

547 (c)

Since,  $\vec{a} \cdot \vec{b} = 0 \dots (i)$

$$\text{Also, } (\vec{a} + 3\vec{b}) \cdot (2\vec{a} - \vec{b}) = -10$$

$$\Rightarrow 2|\vec{a}|^2 - \vec{a} \cdot \vec{b} + 6\vec{b} \cdot \vec{a} - 3|\vec{b}|^2 = -10$$

$$\Rightarrow 2 - 3|\vec{b}|^2 = -10 \Rightarrow |\vec{b}| = 2 \text{ [from Eq. (i)]}$$

548 (a)

We have,  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + \hat{j}, \vec{c} = \hat{i}$

$$\therefore (\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$$

$$\Rightarrow (\vec{c} \cdot \vec{a})\vec{b} - (\vec{c} \cdot \vec{b})\vec{a} = \lambda \vec{a} + \mu \vec{b}$$

$$\Rightarrow \vec{b} - \vec{a} = \lambda \vec{a} + \mu \vec{b}$$

$$\Rightarrow (\lambda + 1)\vec{a} + (\mu - 1)\vec{b} = \vec{0}$$

$$\Rightarrow \lambda + 1 = 0 \text{ and } \mu - 1 = 0 \quad [\because \vec{a}, \vec{b}, \text{ are non-collinear}]$$

$$\Rightarrow \lambda + \mu = 0$$

550 (c)

Let angle between  $\vec{a}$  and  $\vec{b}$  be  $\theta_1$ .  $\vec{c}$  and  $\vec{d}$  be  $\theta_2$  and  $\vec{a} \times \vec{b}$  and  $\vec{c} \times \vec{d}$  be  $\theta$

$$\text{Since, } (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) = 1$$

$$\Rightarrow \sin \theta_1 \cdot \sin \theta_2 \cdot \cos \theta = 1 \quad (\because |\vec{a}| = |\vec{b}| = |\vec{c}| = |\vec{d}| = 1)$$

$$\Rightarrow \theta_1 = 90^\circ, \theta_2 = 90^\circ, \theta = 0^\circ$$

$$\Rightarrow \vec{a} \perp \vec{b}, \vec{c} \perp \vec{d}, (\vec{a} \times \vec{b}) \parallel (\vec{c} \times \vec{d})$$

$$\text{So, } \vec{a} \times \vec{b} = k(\vec{c} \times \vec{d}) \text{ and } \vec{a} \times \vec{b} = k(\vec{c} \times \vec{d})$$

$$\Rightarrow (\vec{a} \times \vec{b}) \cdot \vec{c} = k(\vec{c} \times \vec{d}) \cdot \vec{c}$$

$$\text{and } (\vec{a} \times \vec{b}) \cdot \vec{d} = k(\vec{c} \times \vec{d}) \cdot \vec{d}$$

$$\Rightarrow [\vec{a} \vec{b} \vec{c}] = 0 \text{ and } [\vec{a} \vec{b} \vec{d}] = 0$$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$  and  $\vec{a}, \vec{b}, \vec{d}$  are coplanar vector so option (A) and (B) are incorrect.

$$\text{Let } \vec{b} \parallel \vec{d} \Rightarrow \vec{b} = \pm \vec{d}$$

$$\text{As } (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) = 1 \Rightarrow (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{b}) = \pm 1$$

$$\Rightarrow [\vec{a} \times \vec{b} \vec{c} \vec{b}] = \pm 1$$

$$\Rightarrow [\vec{c} \vec{b} \vec{a} \times \vec{b}] = \pm 1$$


$$\Rightarrow \vec{c} \cdot [\vec{b} \times (\vec{a} \times \vec{b})] = \pm 1$$

$$\Rightarrow \vec{c} \cdot [\vec{a} - (\vec{b} \cdot \vec{a})\vec{b}] = \pm 1$$

$$\Rightarrow \vec{c} \cdot \vec{a} = \pm 1 \quad (\because \vec{a} \cdot \vec{b} = 0)$$

Which is a contradiction so option (c) is correct.

Let option (d) is correct



$$\Rightarrow \vec{d} = \pm \vec{a} \text{ and } \vec{c} = \pm \vec{b}$$

$$\text{As } (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) = 1$$

$$\Rightarrow (\vec{a} \times \vec{b}) \cdot (\vec{b} \times \vec{a}) = \pm 1$$

Which is a contradiction so option (d) is incorrect.

**Alternate** Option (c) and (d) may be observed from given in figure.

552 (b)

$$(\hat{i} \times \hat{j}) \cdot \vec{c} \leq |\hat{i} \times \hat{j}| |\vec{c}| \cos \frac{\pi}{6}$$

$$\Rightarrow -\frac{\sqrt{3}}{2} \leq (\hat{i} \times \hat{j}) \cdot \vec{c} \leq \frac{\sqrt{3}}{2}$$

553 (b)

It is given that  $\hat{a}$  and  $\hat{b}$  are mutually perpendicular unit vectors. Therefore,  $\hat{a}, \hat{b}$  and  $\hat{a} \times \hat{b}$  are non-coplanar vectors.

$$\therefore [\hat{a} \hat{b} \hat{a} \times \hat{b}] \neq 0$$

If the vectors  $\vec{\alpha} = x\hat{a} + y\hat{b} + z(\hat{a} \times \hat{b}), \vec{\beta} = \hat{a} + (\hat{a} \times \hat{b})$

and,  $\vec{\gamma} = z\hat{a} + y\hat{b} + y(\hat{a} \times \hat{b})$  are coplanar, then

$$[\vec{\alpha} \vec{\beta} \vec{\gamma}] = 0$$

$$\Rightarrow \begin{vmatrix} x & x & z \\ 1 & 0 & 1 \\ z & z & y \end{vmatrix} [\hat{a} \hat{b} \hat{a} \times \hat{b}] = 0$$

$$\Rightarrow \begin{vmatrix} x & x & z \\ 1 & 0 & 1 \\ z & z & y \end{vmatrix} = 0 \quad [\because [\hat{a} \hat{b} \hat{a} \times \hat{b}] \neq 0]$$

$$\Rightarrow x(0 - z) - x(y - z) + z(z - 0) = 0$$

$$\Rightarrow -xz - yx + xz + z^2 = 0$$

$$\Rightarrow z^2 = xy$$

$\Rightarrow z$  is the geometric mean of  $x$  and  $y$

554 (d)

Given,  $\vec{a} = (1, p, 1), \vec{b} = (q, 2, 2)$

$$\vec{a} \cdot \vec{b} = r \text{ and } \vec{a} \times \vec{b} = (0, -3, -3)$$

$$\text{Now, } \vec{a} \cdot \vec{b} = (\hat{i} + p\hat{j} + \hat{k}) \cdot (q\hat{i} + 2\hat{j} + 2\hat{k})$$

$$\Rightarrow q + 2p + 2 = r \quad [\text{given}] \dots (i)$$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & p & 1 \\ q & 2 & 2 \end{vmatrix}$$

$$\Rightarrow (2p - 2)\hat{i} + (q - 2)\hat{j} + (2 - pq)\hat{k}$$

$$= \{0\hat{i} + (-3)\hat{j} + (3)\hat{k} \quad [\text{given}]$$

$$\Rightarrow 2p - 2 = 0; q - 2 = -3; 2 - pq = 3$$

$$\Rightarrow p = 1, q = -1$$

From Eqs. (i),

$$-1 + 2 + 2 = r$$

$$= r = 3$$

555 (c)

We have,

$$(3\hat{i} - 2\hat{j} + \hat{k}) \cdot (2\hat{i} + \hat{j} - 4\hat{k}) = 0$$

So, the triangle is right angled

556 (a)

$$\text{Since, } 2|\hat{i} + x\hat{j} + 3\hat{k}| = |4\hat{i} + (4x - 2)\hat{j} + 2\hat{k}|$$

$$\Rightarrow 2\sqrt{1 + x^2 + 9} = \sqrt{4^2 + (4x - 2)^2 + 2^2}$$

$$\Rightarrow 12x^2 - 16x - 16 = 0$$

$$\Rightarrow (3x + 2)(x - 2) = 0$$

$$\Rightarrow x = 2, -\frac{2}{3}$$

559 (b)

$\therefore \vec{a}, \vec{b}$ , and  $\vec{c}$  are the  $p$ th,  $q$ th,  $n$ th terms of an HP respectively.

$$\frac{1}{a} = A + (p - 1)D, \frac{1}{b} = A + (q - 1)D \text{ and } \frac{1}{c}$$

$$= A + (r - 1)D$$

$$\therefore q - r = \frac{c - b}{bcD}, r - p = \frac{a - c}{acD}$$

$$\text{And } q - r = \frac{b - a}{abd}$$

$$\Rightarrow \frac{(q-r)}{a} + \frac{(r-p)}{b} + \frac{(p-q)}{c} = 0$$

$$\Rightarrow \vec{u} \cdot \vec{v} = 0$$

560 (d)

Given edges are

$$\vec{a} = \hat{i} - \hat{k}, \vec{b} = \lambda \hat{i} + \hat{j} + (1 - \lambda)\hat{k}$$

$$\text{and } \vec{c} = \mu \hat{i} + \lambda \hat{j} + (1 + \lambda - \mu)\hat{k}$$

$\therefore$  Volume of parallelepiped

$$= [\vec{a} \vec{b} \vec{c}]$$

$$= \begin{vmatrix} 1 & 0 & -1 \\ \lambda & 1 & 1 - \lambda \\ \mu & \lambda & 1 + \lambda - \mu \end{vmatrix}$$

$$= 1(1 + \lambda - \mu - \lambda + \lambda^2) - 0 - 1(\lambda^2 - \mu)$$

$$= 1 + \lambda^2 - \mu - \lambda^2 + \mu = 1$$

Hence, volume depends on neither  $\lambda$  nor  $\mu$ .

561 (a)

$$\vec{c} \cdot (\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})$$

$$= \vec{c} \cdot (\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b})$$


$$= \vec{c} \cdot \vec{b} \times \vec{a}$$

562 (c)

$$\vec{AC} - \vec{BD}$$

$$= (\vec{AB} + \vec{BC}) - (\vec{BA} + \vec{AD})$$

$$= \vec{AB} + \vec{BC} + \vec{AB} - \vec{AD} = 2\vec{AB}$$



563 (c)

We have,

$$\vec{a} + \vec{b} + \vec{c} = 0$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 0$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 = 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$= 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$$

$$= -\frac{3}{2} [\because |\vec{a}| = |\vec{b}| = |\vec{c}| = 1]$$

565 (a)

Given that,  $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = 5\hat{i} - 3\hat{j} + \hat{k}$

The projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$

$$= \frac{(2\hat{i} + \hat{j} + 2\hat{k}) \cdot (5\hat{i} - 3\hat{j} + \hat{k})}{\sqrt{(2)^2 + (1)^2 + (2)^2}}$$

$$= \frac{10 - 3 + 2}{\sqrt{9}} = \frac{9}{3} = 3$$

566 (a)

Total force,

$$\vec{F} = 3 \left( \frac{6\hat{i} + 2\hat{j} + 3\hat{k}}{7} \right) + 4 \left( \frac{3\hat{i} - 2\hat{j} + 6\hat{k}}{7} \right)$$

$$= \frac{(30\hat{i} - 2\hat{j} + 33\hat{k})}{7}$$

$$\therefore \vec{d} = 4\hat{i} + 3\hat{j} + \hat{k} - (2\hat{i} + 2\hat{j} - \hat{k})$$

$$= 2\hat{i} + \hat{j} + 2\hat{k}$$

$\therefore$  Work done  $W = \vec{F} \cdot \vec{d}$

$$= \left( \frac{30\hat{i} - 2\hat{j} + 33\hat{k}}{7} \right) \cdot (2\hat{i} + \hat{j} + 2\hat{k})$$

$$= \frac{60 - 2 + 66}{7} = \frac{124}{7}$$

567 (b)

$$[\vec{a} \vec{b} + \vec{c} \vec{a} + \vec{b} + \vec{c}] = \vec{a} \cdot \{(\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})\}$$

$$= \vec{a} \cdot \{\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b}\}$$

$$= \vec{a} \cdot \{\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} - \vec{b} \times \vec{c}\}$$

$$= [\vec{a} \vec{b} \vec{a}] + [\vec{a} \vec{c} \vec{a}] = 0$$